

# HILGARDIA

*A Journal of Agricultural Science Published by  
the California Agricultural Experiment Station*

---

VOLUME 17

APRIL, 1947

NUMBER 9

---

## CONTENTS

# DISTRIBUTION OF THE NATIVE GRASSES OF CALIFORNIA

ALAN A. BEETLE

---

UNIVERSITY OF CALIFORNIA • BERKELEY, CALIFORNIA

DISTRIBUTION OF THE NATIVE GRASSES  
OF CALIFORNIA<sup>1</sup>ALAN A. BEETLE<sup>2</sup>

THE GRASSES, supplemented by certain legumes, form the principal basis for range wealth. The natural forage value of the Gramineae as a whole makes an intensive study of their characteristics important, for the broader the knowledge concerning them the more readily may any problem be met. The following paper presents a picture of the current distributions of grasses in California, together with evidences of their floral origins by migration from other regions.

Vegetation has many characteristics which are not always apparent at first glance. For instance, certain elements of the vegetation are native in their location, some are native elsewhere and have only recently been introduced. Some are old species often representative of a primitive condition in their genus, still others appear to be recently evolved. Some of the migrants arrived in California from the north during glacial periods, some crossed the ocean, and others came from the south during interglacial periods. Some plants are distributionally restricted for a number of reasons, including: (1) specialization as to habitat or environmental repression, as the species of vernal pools; (2) recent origin (plants sometimes referred to as neoendemics or initiates), as the endemic varieties of *Distichlis spicata*; (3) ancient origin (paleoendemics or relics); and (4) genotypic specialization (genetic endemics). The limiting factors controlling distribution of grasses are then not always easily recognized, but some attempt has been made to appraise these historical and ecological factors which contribute to an understanding of the present botanical composition of the California rangeland.

Introduced grasses are frequently mentioned in order to complete the overall picture or because of contrast in distributional behavior. Principal attention is paid, however, to the native species, the limits of whose areas are stabilized by climatic and edaphic features interacting over a long period of time. Distributions of 184 California grasses are shown in the maps on pages 319–339. Not all the native species have been mapped but those omitted would not contribute anything new or disturbing. A few introduced species have been mapped where their area of adaptability seemed limited and already occupied.

<sup>1</sup> Received for publication June 20, 1946.<sup>2</sup> Assistant Professor of Agronomy and Assistant Agronomist in the Experiment Station; resigned September 5, 1946.

The first comprehensive list of the grasses of California was made by Bolander (1864-65).<sup>3</sup> At that time he knew 112 species of grasses growing without cultivation, of which 31 were introductions. Soon thereafter Thurber (1880) listed 175 graminaceous species for the state. Now some 400 grasses are known, about one fourth of which are introductions. For convenience in handling the material presented in this study, California has been divided into areas on the basis of recurrent distributions. Neither these areas nor the habitats into which they may be subdivided are wholly contiguous in all their parts. Complex in the same way as biological units, distribution patterns are recognizable but not always disjunct. There are exceptions and overlappings.

## THE NORTHERN FLORAL UNITS IN CALIFORNIA

The northern floral units in California occupy the mountain ranges of northern California southward in the Coast Ranges to Monterey Bay and southward in the Sierra Nevada to Tulare County; the mountain peaks of southern California, particularly the San Bernardino and San Jacinto mountains. This delimitation is arrived at after studying the distributions of the native grasses of northern California. Although not all grasses discussed under this heading occupy this whole area they all fall within its limits and are not found outside it. This area is by far the richest in California in numbers of Gramineae. The grasses are consistently of such typically north-temperate genera as *Melica*, *Festuca*, *Poa*, and *Glyceria*. They have all come to California from the north. One half or better are of the tribe Festuceae, about one fourth are of the Agrostideae, one eighth are of the Aveneae, and the remaining one eighth represent all other tribes.

The inclusion of northern California and the coastal pine forests with the Sierra Nevada as a homogeneous floral unit is substantiated by Cooke (1940, 1941) who, in two lists containing 487 species, varieties, subspecies, and forms, could find "no endemic plants"; most of the plants were "fairly common" though this uniform picture may have been disturbed partially by Swallen's subsequent description (1941) of *Glyceria cookei* (map 89). The endemic nature of *Panicum shastense* (map 93) is open to question and depends upon the interpretation of specific limits in *Panicum*.

The pine woods offer the only relief for the agrostologist from the host of introductions that have so altered the aspect of the California flora. Here, alone, native grasses are encountered to the near exclusion of all aliens. Their history is undoubtedly connected with that of the coniferous forests. If the fossil records of these conifers could be coordinated with the known ranges of the grass species and their nearest relatives, considerable light should be shed on the origin of many California grasses.

**Northern Grasses That Reach Their Southern Limit in California** (maps 1 to 27).—The general predominance of species of or from northern floras that are characteristic of northern California is emphasized when one picks up a flora of Nova Scotia (Dore and Roland, 1941-42) or of Alaska (Scribner and Merrill, 1910) and finds listed in either not only genera but species typical of the northern California floral unit—for example, *Agro-*

<sup>3</sup> See "Bibliography" for complete citations, referred to in the text by author and date of publication.

*pyron trachycaulum*, *Alopecurus aequalis* (map 60), *Cinna latifolia* (map 36), *Calamagrostis canadensis* (map 42), *Deschampsia caespitosa* (map 2), *Koeleria cristata* (map 58), *Trisetum spicatum* (map 35), *Poa palustris*, and *Festuca ovina* (map 55). Of these species, three—*Trisetum spicatum*, *Deschampsia caespitosa*, and *Koeleria cristata*—reappear in South America, as do *Phleum alpinum* (map 5), *Deschampsia danthonioides* (map 66), *D. elongata* (map 67), and a near relative of *Melica subulata* (map 9).

Of similar wide distribution northward are *Poa pratensis* (map 1), *Agrostis hiemalis* (map 3), *A. exarata* (map 17), *Glyceria pauciflora* (map 7), *G. borealis* (map 11), *G. striata* (map 22), *Festuca rubra* (map 8), *F. subulata* (map 15), *Beckmannia syzigachne* (map 13), *Puccinellia distans* (map 23), and *Deschampsia atropurpurea* (map 26). These grasses are never found in the Great Valley. They are, however, of wide distribution within the limits defined above—that is, they cover a considerable altitudinal range, often occurring in both the Coast Ranges and the Sierra Nevada. In the Coast Ranges they are not found south of Monterey Bay and in southern Sierra Nevada they are not found between Tulare County and the San Bernardino Mountains.

Of somewhat narrower distribution northward but of similar occurrence in California are *Glyceria elata* (map 4), *Trisetum cernuum* var. *canescens* (map 6), *Poa cusickii* (map 10), *Festuca idahoensis* (map 14), *Melica spectabilis* (map 19), *Festuca occidentalis* (map 21), *Puccinellia lemmoni* (map 25), and *Calamagrostis rubescens* (map 27).

Even more restricted but still within the same pattern are *Melica fugax* (map 12), *Festuca subuliflora* (map 20), and *Scribneria bolanderi* (map 24).

The continuous chain of the Sierran peaks is finally broken in Tulare County. A considerable element of the grass flora has, however, been able to bridge the barrier presented by the intervening low elevations and reappears on the San Jacinto, San Bernardino, and other peaks of southern California. The majority of these are found in the San Bernardino; fewer are known from the other peaks. Species exemplifying this hiatus include, in part:

|   |   |
|---|---|
| <i>Agrostis exarata</i> (map 17)        | <i>Muhlenbergia richardsonis</i> (map 40) |
| <i>Phleum alpinum</i> (map 5)           | <i>Festuca ovina</i> (map 55)             |
| <i>Agrostis idahoensis</i> (map 39)     | <i>Muhlenbergia andina</i> (map 46)       |
| <i>Deschampsia caespitosa</i> (map 2)   | <i>Festuca viridula</i> (map 48)          |
| <i>Muhlenbergia filiformis</i> (map 37) | <i>Glyceria elata</i> (map 4)             |
| <i>Poa bolanderi</i> (map 38)           |   |

Here they join a small group of grasses including *Muhlenbergia californica* (map 108), *Poa atropurpurea*, and *P. longiligula* (known also from Rock Creek Lake Basin, Inyo County), which are isolated in range but related to the Sierran species. They have not, on the other hand, been able to bridge the gap between these mountains and those of Mexico. True, *Agrostis exarata* is found in Mexico, but its migrational path was not south in the California Sierra Nevada but rather southward in the Rocky Mountain chain through New Mexico and western Texas.

Few annuals are really characteristic of this zone, and these usually occur only at the lower elevations; they are far outranked by the large number of perennials. Some of the introduced annuals listed under open range and



savanna, because of their wide tolerance and ready adaptability, are easily found here, but so are *Aira praecox* and *Festuca myuros*, which occur only within the area defined. The natives include *Festuca microstachys*, *Muhlenbergia filiformis*, and *Poa bolanderi*. *Muhlenbergia filiformis*, usually listed as a perennial, is treated here as an annual (or at least a very short-lived perennial) because it occurs on sandy stream banks and open washes associated with other very characteristic and typically tiny, annual species including *Eleocharis bella*, *Cyperus aristatus*, and many others.

The mountain meadows contain an impressive list of economically important introduced perennials:

*Agropyron cristatum*  
*Agropyron repens*  
*Agrostis alba*  
*Anthoxanthum odoratum*  
*Arrhenatherum elatius*  
*Bromus inermis*

*Dactylis glomerata*  
*Festuca elatior*  
*Phleum pratense*  
*Poa bulbosa*  
*Poa compressa*  
*Poa pratensis*

Except for sweet vernal grass (*Anthoxanthum odoratum*), a filler usually planted only for the fragrance it gives to hay, all the above introductions are of considerable economic importance. There can be no doubt that meadows in which they are well established are a greater economic resource to their owners than are mountain meadows, usually dominated by *Carex*, where the native vegetation is undisturbed. It must be remembered, however, in attempting to establish these grasses, to which areas of the state they are adaptable. It is not at all surprising that Sampson (1944) had no success in his attempts to reseed burned areas in the chaparral belt with six of the above listed species.

*Agrostis alba* and *Poa pratensis* have long been known in commercial seed mixtures. They were first brought into cultivation in the Old World and were among the first plants sown for hay in this country by settlers, the origin of the seed being the markets of the Old World. It has then usually been assumed that these grasses were introduced in all their occurrence in this country. Present-day study of ecotypes suggests that these species may have had a panboreal distribution and may have arrived in North America long before the white man. Subsequent crossing of the introduced with the native strains would have complicated the study of the origin of these grasses.

**Grasses of High Elevations in the Sierra Nevada** (maps 28 to 54).—Many grasses which are common at high elevations (about 7,000 to 9,000 feet) in the Sierra Nevada are not found elsewhere in California, except in the northern Inner Coast Range. In the Sierra Nevada they follow the mountains to Tulare County and then frequently reappear in the San Bernardino Mountains. They are not known in the Great Valley. A few that fit this California pattern are of a very widespread distribution northward—for example, *Trisetum spicatum* (map 35), *Cinna latifolia* (map 36), *Calamagrostis canadensis* (map 42), *Bromus ciliatus* (map 47), and *Danthonia intermedia* (map 51). Most of these grasses are more characteristic of a more restricted area—from British Columbia to Montana southward to California and New Mexico—for example, *Agrostis thurberiana* (map 28), *A. idahoensis* (map 39), *A. rossae* (map 43), *Bromus suksdorfii* (map 45), *Melica stricta* (map 29), *M. bulbosa* (map 44), *M. aristata* (map 30), *Poa pringlei* (map 31), *P. epilis*

(map 32), *P. bolanderi* (map 38), *Glyceria erecta* (map 33), *Danthonia unispicata* (map 34), *Muhlenbergia filiformis* (map 37), *M. andina* (map 46), *M. montana* (map 53), *Festuca viridula* (map 48), *Trisetum wolffii* (map 50), and *Stipa webberi*. Only a few are endemic to the Sierra Nevada—for example, *Trisetum cernuum* var. *projectum* (map 54), *Oryzopsis kingii*, and *Calamagrostis breweri* (map 41).

**Alpine Grasses** (maps 55 to 60).—There are only six truly alpine grasses in California, namely *Alopecurus aequalis* (map 60), *A. geniculatus* (map 59), *Glyceria californica* (map 57), *Koeleria cristata* (map 58), *Poa rupicola* (map 56), and *Festuca ovina* (map 55). The small number of alpine grasses is remarkable and attests the youthfulness of the present alpine peaks. The species are specialized representatives, closely related to other members of their genera found on the lower slopes. None is endemic to the Sierra Nevada. Although isolated in the Sierran part of their range, specific differences have not been found in the Sierran material of *Koeleria cristata*, *Alopecurus geniculatus*, and *A. aequalis*.

Russell (1926) in his discussion of California climates states that "a narrow strip in the High Sierra . . . [is] mapped as Tundra Climate. . . This treeless alpine area differs sufficiently from the Microthermal area to justify its being placed in another type and there is little doubt that it meets the Köppen requirement of having all monthly means below 50°F." The true alpine grasses of California are closely confined to this narrow strip, at least in their high Sierran phase.

**Grasses of the Northern Coast Ranges and Low to Middle Elevations in the Sierra Nevada** (maps 61 to 81).—Still following the pattern of the grasses of northern California—that is, in the Coast Ranges to Monterey Bay, in the Sierra Nevada to Tulare County, and reappearing on the peaks of some southern California ranges—are a group whose center of abundance is in the northern Coast Ranges but which occur also in the Sierra Nevada, usually at low elevations. These grasses are again of northern origin, and in most cases represent different species of the same genera already mentioned. They tend to have narrower distributions, mostly within the limits of British Columbia to Idaho and southward to California—for example, *Melica geyeri* (map 61), *M. harfordii* (map 62), *M. californica* (map 63), *Festuca elmeri* (map 64), *Poa rhizomata* (map 65), *Panicum occidentale* (map 69), *Panicum pacificum* (map 70), *Stipa lemmoni* (map 72), *Festuca californica* (map 74), *Poa howellii* (map 75), *Bromus orcuttianus* (map 79), *B. vulgaris* (map 81), and *B. laevipes* (map 80). Only two—*B. grandis* (map 73) and *Melica torreyana* (map 76)—are actually confined to California.

**Grasses of the Northern Coast Ranges** (maps 82 to 99).—This final group occupying northern California contains species mostly endemic to California and usually of very restricted range. They are, again, all found northward from Monterey Bay, and thus form a strong contrast to the Coast Ranges from Monterey Bay southward to San Diego County wherein there is not known a single endemic of narrow or restricted range.

Characteristic of the Outer Coast Range and extending into Oregon are *Deschampsia caespitosa* var. *holciformis* (map 90), *Hierochloa occidentalis* (map 91), *Phalaris californica* (map 92), and *Calamagrostis nutkaensis*

(map 98). Near the coast are the very rare *Hystrix californica* and *Agrostis californica* (map 99). More obviously specialized in habitat are *Poa napensis* (map 87) from a hot spring and *Calamagrostis crassiglumis* (map 95) from sphagnum bogs. Of the three species of *Pleuropogon*—*P. refractus* (map 84), *P. davyi* (map 85), and *P. californicus* (map 83)—two are endemic to California but fit the general scheme of northward relationship, for the only remaining known species in the genus are the very restricted *P. oregonus* and the circumboreal *P. sabinii*.

## GREAT BASIN AND DESERT FLORAL UNITS IN CALIFORNIA

**Desert Grasses** (maps 100 to 126).—Species on nonalkaline soils in the desert include:

**Native perennials**

*Aristida adscensionis* (map 118)  
*Aristida californica* (map 119)  
*Aristida divaricata* (map 120)  
*Aristida fendleriana* (map 121)  
*Aristida glabrata* (map 122)  
*Aristida glauca* (map 123)  
*Aristida parishii* (map 124)  
*Aristida purpurea* (map 125)  
*Aristida wrightii* (map 126)  
*Bouteloua aristidoides* (map 114)  
*Bouteloua barbata* (map 115)  
*Bouteloua curtipendula* (map 116)  
*Bouteloua gracilis* (map 100)  
*Bouteloua rothrockii* (map 117)  
*Hilaria rigida* (map 104)  
*Panicum urvilleanum* (map 103)  
*Poa bigelovii* (map 110)  
*Triodia pulchella* (map 113)

**Native annuals**

*Cenchrus echinatus*  
*Cenchrus pauciflorus*

**Introduced annual**

*Schismus barbatus*

**Weeds of cultivated land**

*Eriochloa gracilis*  
*Eriochloa aristata*  
*Hordeum jubatum*  
*Leptochloa filiformis* (map 101)  
*Leptochloa uninervia* (map 102)

**Introduced or subtropical economic  
grasses**

*Oryzopsis miliacea*  
*Panicum miliaceum*  
*Chloris gayana*  
*Chloris virgata*  
*Cynodon dactylon*

In connection with the flora of the extreme southern desert part of California, it is of interest to review the influence of the tremendously rich Mexican flora (Conzatti, 1943; Johnston, 1943) on that of California. The Mexican flora has roughly twice the number of native species that occur in California. This would indicate that so far as the grasses are concerned the relatively more stable geologic conditions that have existed in Mexico have been more favorable for the development and persistence of species than have the fluctuations in sea level, the glaciation, and the mountain unrest that have characterized California's past. Moreover, the sea and desert barriers that have existed through much of the past have been so effective in preventing a free interchange of species that, except for desert species and a few other exceptions the two floras are entirely distinct.

Although perhaps the best indicator species are tall, stiff *Hilaria rigida* (map 104) on the flats and low, fluffy *Triodia pulchella* (map 113) on the rocky slopes, in numbers *Aristida* (nine species) (maps 118 to 126) and *Bouteloua* (five species) (maps 100, 114 to 117) dominate the desert grass flora. This flora, in contrast to all the others found in California, has its strongest ties eastward and southward. *Bouteloua*, *Aristida*, and *Muhlenbergia* are well represented in the Mexican flora, each having one or two species in common with South American floras and in South America a second center of diversification. Clements (1934) postulates the "transformation of southeastern California from bunch-grass prairie to desert as a consequence of climatic changes" and describes *Hilaria rigida* as "derived directly from the adaptation of *Hilaria jamesii* (map 140) to a drying climate."



The close ties between the temperate floras of North and South America, already shown in the Sierran and Great Valley floras, are further demonstrated by the presence of these desert elements common to both hemispheres:

*Aristida*: compare the distributions of *A. ternipes* Cav. and *A. capillacea* Lam.  
*Bouteloua curtipendula* (Griffiths, 1912)  
*Cenchrus echinatus* (Hitchcock and Chase, 1920)  
*Cenchrus pauciflorus* (Hitchcock and Chase, 1920)  
*Distichlis spicata* (Hitchcock, 1927; Beetle, 1943a)  
*Muhlenbergia asperifolia* (Hitchcock, 1927; Parodi, 1928; Beetle, 1943c)  
*Panicum urvilleanum* (Hitchcock, 1935)  
*Sporobolus argutus* (Hitchcock, 1927)  
*Stipa speciosa* (Hitchcock, 1927)

In direct contrast to this North and South American relationship between the species of *Bouteloua*, *Muhlenbergia*, *Aristida*, and other genera is the situation in *Hierochloa*. The seven South American species of this genus (Parodi, 1941) show no immediate relationship to those of North America. They are concentrated in Magellanic floras (where *Muhlenbergia*, *Bouteloua*, and *Aristida* are entirely absent) and are linked to New Zealand by *Hierochloa redolens* ("Habita en el Ecuador, en el sur de Chile, y Argentina y en Nueva Zelandia"). Here is the strongest possible evidence in support of panantartic dispersals.

Species of desert alkaline flats include:

*Distichlis spicata* var. *divaricata* (map 111)  
*Puccinellia parishii* (map 112)  
*Sporobolus airoides*  
*Sporobolus contractus* (map 106)  
*Sporobolus cryptandrus*

**Great Basin Grasses.**—The Great Basin has contributed a large number of grasses to the borders of California—that is, to the areas that lie eastward or southward of the Sierran crest, which has been almost without exception an effective barrier to their further dispersal. These Great Basin grasses include a number of alkali-tolerant species, *Poa ampla* (map 131), *Poa nevadensis* (map 132), *Puccinellia nuttalliana* (map 134), *Distichlis spicata* var. *stricta* (map 133), *Spartina gracilis* (map 137), and *Muhlenbergia asperifolia* (map 139). The other, alkali-intolerant species include a small but diverse Great Basin (and Colorado Desert) element:

|  |                                   |
|--|-----------------------------------|
| <i>Agropyron smithii</i> (map 143)     | <i>Poa fendleriana</i> (map 129)  |
| <i>Agropyron spicatum</i> (map 144)    | <i>Poa leptocoma</i> (map 130)    |
| <i>Blepharidachne kingii</i> (map 127) | <i>Stipa comata</i>               |
| † <i>Elymus cinereus</i> (map 136)     | <i>Stipa speciosa</i> (map 142)   |
| <i>Festuca kingii</i> (map 128)        | <i>Stipa williamsii</i> (map 141) |
| <i>Hilaria jamesii</i> (map 140)       | <i>Triodia mutica</i> (map 135)   |
| <i>Oryzopsis hymenoides</i> (map 138)  |                                   |


These Great Basin species occupy the area of transition between the mountains and the desert. The group meets the desert grasses in central San Bernardino County but in general does not overlap their distribution.

## DISTRIBUTION MAPS FOR CALIFORNIA GRASSES

IN THE MAPS on the following pages, grasses with similar distributions are grouped together. Most of the grasses included are native species, but not all native species discussed in this paper are mapped. Those omitted would not affect the conclusions of the paper. The few introduced species mapped are ones that seem to have fully occupied the area to which they are adapted—that is, ones whose distribution has become relatively stable.




NORTHERN GRASSES THAT REACH THEIR SOUTHERN LIMIT IN CALIFORNIA




Map 1

*Poa pratensis*



Map 2

*Deschampsia caespitosa*




Map 3

*Agrostis hiemalis*




Map 4

*Glyceria elata*




Map 5

*Phleum alpinum*




Map 6

*Trisetum cernuum* var. *canescens*




Map 7

*Glyceria pauciflora*



Map 8

*Festuca rubra*




Map 9

*Melica subulata*




# NORTHERN GRASSES THAT REACH THEIR SOUTHERN LIMIT IN CALIFORNIA



Map 10

*Poa cusickii*



Map 11

*Glyceria borealis*



Map 12

*Melica fugax*



Map 13

*Beckmannia syzigachne*



Map 14

*Festuca idahoensis*




Map 15

*Festuca subulata*



Map 16

*Deschampsia caespitosa* var. *arctica*



Map 17

*Agrostis exarata*



Map 18

*Trisetum cernuum*

NORTHERN GRASSES THAT REACH THEIR SOUTHERN LIMIT IN CALIFORNIA




Map 19

*Melica spectabilis*




Map 20

*Festuca subuliflora*




Map 21

*Festuca occidentalis*




Map 22

*Glyceria striata*




Map 23

*Puccinellia distans*




Map 24

*Scribneria bolanderi*



Map 25

*Puccinellia lemmoni*



Map 26

*Deschampsia atropurpurea*



Map 27

*Calamagrostis rubescens*

GRASSES OF HIGH ELEVATIONS IN THE SIERRA NEVADA




*Agrostis thurberiana*




*Melica stricta*




*Melica aristata*



*Poa pringlei*




*Poa epilys*




*Glyceria erecta*



*Danthonia unispicata*




*Trisetum spicatum*



*Cinna latifolia*

# GRASSES OF HIGH ELEVATIONS IN THE SIERRA NEVADA




Map 37

*Muhlenbergia filiformis*




Map 38

*Poa bolanderi*



Map 39

*Agrostis idahoensis*




Map 40

*Muhlenbergia richardsonis*




Map 41

*Calamagrostis breweri*



Map 42

*Calamagrostis canadensis*



Map 43

*Agrostis rossae*



Map 44

*Melica bulbosa*



Map 45

*Bromus suksdorfii*




# GRASSES OF HIGH ELEVATIONS IN THE SIERRA NEVADA



Map 46

*Muhlenbergia andina*




Map 47

*Bromus ciliatus*




Map 48

*Festuca viridula*




Map 49

*Poa gracillima*




Map 50

*Trisetum wolffii*



Map 51

*Danthonia intermedia*



Map 52

*Poa suksdorfii*



Map 53

*Muhlenbergia montana*



Map 54

*Trisetum cernuum* var. *proectum*

# ALPINE GRASSES



Map 55

*Festuca ovina*




Map 56

*Poa rupicola*



Map 57

*Glyceria californica*




Map 58

*Koeleria cristata*



Map 59


*Alopecurus geniculatus*



Map 60


*Alopecurus aequalis*

## GRASSES OF THE NORTHERN COAST RANGES AND LOW TO MIDDLE ELEVATIONS IN THE SIERRA NEVADA




Map 61

*Melica geyeri*



Map 62


*Melica barfordii*



Map 63

*Melica californica*


GRASSES OF THE NORTHERN COAST RANGES AND LOW TO  
MIDDLE ELEVATIONS IN THE SIERRA NEVADA




*Festuca elmeri*




*Poa rhizomata*



*Deschampsia danthonioides*



*Deschampsia elongata*




*Danthonia californica*



*Panicum occidentale*



*Panicum pacificum*




*Agrostis diegoensis*



*Stipa lemmoni*

GRASSES OF THE NORTHERN COAST RANGES AND LOW TO  
MIDDLE ELEVATIONS IN THE SIERRA NEVADA




*Bromus grandis*




*Festuca californica*




*Poa bowellii*




*Melica torreyana*



*Bromus racemosus*



*Bromus commutatus*



*Bromus orcuttianus*



*Bromus laevipes*



*Bromus vulgaris*



# GRASSES OF THE NORTHERN COAST RANGES




Map 82

*Poa trivialis*




Map 83

*Pleuropogon californicus*



Map 84

*Pleuropogon refractus*




Map 85

*Pleuropogon davyi*



Map 86

*Glyceria leptostachya*



Map 87

*Poa napensis*



Map 88

*Glyceria occidentalis*



Map 89

*Glyceria cookei*



Map 90


*Deschampsia caespitosa*  
var. *bolciformis*

# GRASSES OF THE NORTHERN COAST RANGES



Map 91

*Hierochloa occidentalis*




Map 92

*Phalaris californica*



Map 93

*Panicum shastense*



Map 94

*Panicum scribnerianum*



Map 95

*Calamagrostis crassiglumis*




Map 96

*Calamagrostis foliosa*



Map 97

*Calamagrostis koelerioides*



Map 98

*Calamagrostis nutkaensis*



Map 99

*Agrostis californica*

# DESERT GRASSES



Map 100

*Bouteloua gracilis*



Map 101

*Leptocloa filiformis*



Map 102

*Leptocloa uninervia*



Map 103

*Panicum urvilleanum*



Map 104

*Hilaria rigida*



Map 105

*Muhlenbergia porteri*



Map 106

*Sporobolus contractus*



Map 107

*Stipa speciosa*



Map 108


*Muhlenbergia californica*

# DESERT GRASSES



Map 109

*Melica frutescens*



Map 110

*Poa bigelovii*




Map 111

*Distichlis spicata* var. *divaricata*



Map 112

*Puccinellia parishii*



Map 113

*Triodia pulchella*



Map 114

*Bouteloua aristidoides*



Map 115

*Bouteloua barbata*



Map 116

*Bouteloua curtipendula*



Map 117

*Bouteloua rothrockii*

# DESERT GRASSES




Map 118

*Aristida adscensionis*



Map 119

*Aristida californica*



Map 120

*Aristida divaricata*



Map 121

*Aristida fendleriana*




Map 122

*Aristida glabrata*



Map 123

*Aristida glauca*




Map 124

*Aristida parishii*



Map 125


*Aristida purpurea*



Map 126


*Aristida wrightii*

# GREAT BASIN GRASSES



Map 127

*Blepharidachne kingii*




Map 128

*Festuca kingii*




Map 129

*Poa fendleriana*



Map 130

*Poa leptocoma*



Map 131

*Poa ampla*




Map 132

*Poa nevadensis*



Map 133

*Distichlis spicata* var. *stricta*



Map 134

*Puccinellia nuttalliana*



Map 135

*Triodia mutica*

# GREAT BASIN GRASSES



Map 136

*Elymus cinereus*




Map 137

*Spartina gracilis*




Map 138

*Oryzopsis hymenoides*



Map 139

*Muhlenbergia asperifolia*



Map 140

*Hilaria jamesii*




Map 141

*Stipa williamsii*




Map 142

*Stipa speciosa*



Map 143


*Agropyron smithii*



Map 144

*Agropyron spicatum*

# COASTAL GRASSES




Map 145

*Poa macrantha*




Map 146

*Poa douglasii*



Map 147

*Poa unilateralis*




Map 148

*Puccinellia grandis*




Map 149

*Elymus mollis*



Map 150

*Elymus vanconverensis*




Map 151

*Elymus virescens*



Map 152

*Agrostis blasdalei*



Map 153

*Agropyron arenicola*




# COASTAL GRASSES



Map 154

*Phragmites communis*



Map 155

*Monanthochloe littoralis*



Map 156

*Distichlis spicata* var. *stolonifera*



Map 157

*Dissantheium californicum*



Map 158

*Lepturus cylindricus*



Map 159

*Pholiurus incurvus*



Map 160

*Spartina foliosa*



Map 161


*Ammophila arenaria*

GRASSES OF THE SOUTHERN COAST RANGES AND GREAT VALLEY



Map 162

*Mublenbergia microsperma*




Map 163

*Mublenbergia rigens*



Map 164

*Stipa cernua*



Map 165

*Stipa pulchra*




Map 166

*Stipa lepida*



Map 167

*Melica imperfecta*



Map 168

*Koeleria cristata*



Map 169


*Stipa coronata*



Map 170

*Elymus condensatus*

# GRASSES OF THE GREAT VALLEY



Map 171

*Orcuttia greenei*




Map 172

*Orcuttia tenuis*




Map 173

*Orcuttia pilosa*




Map 174

*Orcuttia californica*



Map 175

*Neostaphia colusana*




Map 176

*Distichlis spicata* var. *nana*




Map 177

*Puccinellia simplex*



Map 178

*Eragrostis hypnoides*



Map 179

*Eragrostis orcuttiana*


# GRASSES OF THE GREAT VALLEY




Map 180  
*Deschampsia danthonioides* var. *gracilis*




Map 181  
*Leptochloa fascicularis*



Map 182  
*Aristida hamulosa*



Map 183  
*Aristida oligantha*



Map 184  
*Alopecurus saccatus*



## THE CENTRAL VALLEY AND COASTAL FLORAL UNITS IN CALIFORNIA

### ANALYSIS BY DISTRIBUTION

**Coastal Grasses** (maps 145 to 161).—Many of the coastal grasses are typical of the northern grasses discussed earlier—that is, they occur only north of Monterey Bay and belong to genera whose distributions are north temperate. They tend to be concentrated at such points along the coast as Humboldt Bay, Point Arena, Point Reyes, and the Monterey Peninsula. This is partly, but not wholly, due to restrictions of habitat. The disrupted ranges of the very rare *Agrostis blasdalei* (map 152) and *Agropyron arenicola* (map 153) illustrate this. Other representatives of northern genera that are restricted to coastal habitats include two closely related *Poa* species—*P. macrantha* (map 145) and *P. douglasii* (map 146)—the giant ryes, *Elymus mollis* (map 149) and *E. vancouverensis* (map 150), and the rare and little-known *Puccinellia grandis* (map 148).

Grasses which occur south of Monterey Bay (often as well as northward) are more typically of south-temperate genera and are often introduced; for example, *Lepturus cylindricus* (map 158), *Pholiurus incurvus* (map 159) and *Ammophila arenaria* (map 161).

**Grasses of the Southern Coast Ranges and Great Valley** (maps 162 to 170).—The native grass flora of the Coast Ranges from Monterey County south is very meager. The most typical are a trio of *Stipa* species, *S. cernua* (map 164), *S. pulchra* (map 165), *S. lepida* (map 166), and two *Muhlenbergia* species, *M. microsperma* (map 162), and *M. rigens* (map 163). These grasses may extend northward from Monterey Bay although, with the exception of *Koeleria cristata* (map 168), they do not reach the Oregon border. The abrupt termination in northward distribution at Monterey Bay of *Muhlenbergia microsperma*, *M. rigens*, *Stipa coronata* (map 169), and *Elymus condensatus* (map 170) emphasizes the importance of the area as the most critical in the Coast Ranges; see discussion of the southern termination of northern grasses.

*Stipa cernua* and *S. pulchra* occupy both the central valley and the southern Coast Ranges without a break in their distribution. This distribution, but with hiatus involving the western half of the valley and the Inner Coast Range, is found with *Andropogon virginicus* var. *hirsutior*,<sup>4</sup> *Muhlenbergia microsperma*, *M. rigens*, *Aristida hamulosa* (map 182), *Deschampsia danthonioides* var. *gracilis* (map 180), and *Sporobolus microspermus*.

**Great Valley Grasses** (maps 171 to 184).—The four species of *Orcuttia*, *O. greenei* (map 171), *O. tenuis* (map 172), *O. pilosa* (map 173), and *O. californica* (map 174), as well as *Neostaphia colusana* (map 175) lend a strong degree of endemism to the grass flora of the Great Valley. These grasses are all rare and specialized in habitat (see discussion under "Vernal Pools," p. 347–8). The grasses of the Great Valley are not found in the surrounding hills although they may occupy various of the low valleys which are adjacent

<sup>4</sup> The California phase of *Andropogon virginicus* L. should be treated as *Andropogon virginicus* var. *hirsutior* (Hack.) Hitchc., f. *tenuispatheus* (Nash) comb. nov. (*A. glomeratus tenuispatheus* Nash, in Small, Fl. S.E. U.S. 61. 1903; *A. tenuispatheus* Nash, in Gramineae. N. Amer. Fl. 17: 113. 1912.)

to the Great Valley—for example, *Alopecurus saccatus* (map 184), *Eragrostis hypnoides* (map 178), and *E. orcuttiana* (map 179). All the species are annual except *Aristida hamulosa* (map 182) and *Distichlis spicata* var. *nana* (map 176).

#### ANALYSIS BY HABITAT

Central California, as defined here, includes the Great Valley (San Joaquin and Sacramento valleys) and the surrounding upper Sonoran zone (to an elevation of about 3,000 to 4,000 feet) which is characteristically either open grassland, parklike savanna (the scattered trees usually oaks or digger pine), or chaparral. These habitats tend to be zonal in distribution (Piemeisel and Lawson, 1937), the alkali flats of the poorly drained bottomlands being bordered by grassland, and then chaparral or savanna on the slopes.

Open grass slopes dominate this area, and their typical golden color during most of the year is produced by dry grass straw. The first fall rains germinate the seeds of the annuals and renew the vegetative growth of the perennials. Duration of the growth depends upon the season; the hills stay green for the colder, wetter months, and in early spring begin to turn as the annuals rapidly mature and dry. The last seeds of the annuals are usually shed by the middle of April. Most of the perennials mature nearly as rapidly as the annuals, with the last of them maturing by June. Prolongation of the period of grazing on green feed at this period is one of the important range problems.

The only major break in the coastal hills occurs at San Francisco Bay where the great rivers, having converged, finally enter the ocean. This break considerably affects the climate of the region far into the valley but not sufficiently to form a barrier to plant dispersal. The annual rainfall in this central portion of the Great Valley reaches an average of 15 inches, but tapers off to the north and to the south. The Coast Ranges are lowest at San Francisco Bay (and Suisun Bay), and become higher and steeper northward and southward.

Coastal California is an area of highly specialized habitats. In some places chaparral, open grassland, or oak savanna approach the ocean, but more often there is an intervening strip, usually very narrow, of either coniferous woods (redwood, Monterey cypress, or Douglas-fir), or sand dunes or salt marsh. All owe their existence to, and are modified by, the proximity of the ocean, though in different ways—the salt in the marshes, the wind against the bluffs, the fog in the redwood belt.

All the habitats mentioned may occur also on the islands off the coast of California, to which the rare *Dissanthelium californicum* (map 158) is the only grass species known to be endemic. See Eastwood (1941) for a list of the species that have been collected on the islands.

**Grasses of the Grassland and Savanna.**—Parts of this portion of California appear to have been originally as nearly a climax type of grassland as the more extensive prairies of the interior of North America. The California grasslands were then dominated by bunch grasses liberally supplemented by erect perennials and a few stoloniferous perennials and interspersed with annuals. These areas differ in two respects from similar areas elsewhere in the world: (1) the identity of the perennial species; (2) the number and importance of the annuals. In character the California grassland is related

to similar areas in eastern Oregon, eastern Washington and adjacent Idaho, and in western Montana. Together they comprise the Pacific bunch-grass area, as distinguished from the tall-grass and short-grass prairies of the Great Plains (McArdle, *et al.*, 1936). A far greater proportion of the species in the Pacific bunch-grass area, and even in the California area, trace their origins northward than is true in the tall-grass prairie (Weaver and Fitzpatrick, 1932) or the short-grass prairie, where a large southern element is usually dominant. Only *Koeleria cristata* occupies an important place in both the tall-grass and the Pacific bunch-grass floras.

The original appearance of the California grasslands is not a matter of historical record. Spanish settlers tilled the soil and grazed the hills for many years without caring to write about their surroundings in anything but the most general terms. As evidenced by adobe-brick studies (Hendry, 1931) such introductions as *Poa annua*, *Hordeum murinum*, and *Lolium multiflorum* made headway almost immediately. It is popularly supposed that the native bunch-grass perennials were far more abundant at the time and that they have since tended to disappear under the stress of cultivation, heavy grazing, and the competition of introductions. Since this area coincides with that principally occupied by California's enormous agricultural industry, such displacement is to be expected. Certainly in this portion of California no undisturbed remnants of the original flora remain (disturbance includes the effect of plant on plant as well as man on a plant community), and because of this its appearance is a matter for speculation only. Clements (1934) describes from observations made in 1918 the obliteration of "many hundred miles of a nearly continuous consociation of *Stipa pulchra*," and the establishment of "the wild oats as the one great dominant throughout." There is no evidence, however, that any grass ever recorded for California has become extinct.

The number of strong contrasts between the California grassland and that of the Great Plains eliminates the possibility of any great similarity between the two. The California range is predominantly annual and therefore strongly fluctuating, whereas the Great Plains area is predominantly perennial and therefore potentially stable. The California range developed under a natural system of year-round grazing by deer, whereas the Great Plains area developed under a natural system of rotation grazing by buffalo. In California today introductions are predominant, and the state of the virgin range can no longer be considered a clue to its potential carrying capacity. In the Great Plains area natives are predominant today, and the state of the virgin range is an index to the carrying capacity. In California controlled grazing began around 1773, but on the Great Plains controlled grazing started about 1880. The two areas differ in the identity of both the native species and the introduced, often weedy species. They differ also in the degree of variability of the factors which influence the determination of grazing capacity. The difficulties in determining the carrying capacity of a given area are multiplied proportionately to: (1) the diversity of the vegetation; (2) the variability of the rainfall; (3) the variability of the climate; (4) the variability of slope; (5) the variability of the altitude; and (6) the diversity of soil types. Even within the counties of California all of the factors are present in a degree of variability not met in whole states in the Great Plains region.



The native bunch grasses (that is, nonstoloniferous types), whose conspicuous scattered clumps give the California part of the Pacific grassland its characteristic aspect, include:

|  |                                     |
|--|-------------------------------------|
| <i>Agrostis ampla</i>                              | <i>Melica californica</i>           |
| <i>Andropogon virginicus</i> var. <i>hirsutior</i> | <i>Melica imperfecta</i>            |
| <i>Aristida hamulosa</i>                           | <i>Muhlenbergia rigens</i>          |
| <i>Bromus carinatus</i>                            | <i>Panicum pacificum</i> and allies |
| <i>Bromus laevipes</i>                             | <i>Sitanion hystrix</i>             |
| <i>Danthonia californica</i>                       | <i>Sitanion jubatum</i>             |
| <i>Elymus condensatus</i>                          | <i>Stipa cernua</i>                 |
| <i>Elymus glaucus</i>                              | <i>Stipa lepidia</i>                |
| <i>Koeleria cristata</i>                           | <i>Stipa pulchra</i>                |

Successfully introduced stoloniferous perennials in this area are few because of moisture limitations (compare *Agrostis stolonifera*, *Phalaris arundinacea*, and *P. tuberosa* var. *stenoptera*). Of the grasses in this area, only *Elymus triticoides* is strongly rhizomatous.

| Introduced perennials        | Introduced annuals          |
|------------------------------|-----------------------------|
| <i>Agrostis retrofracta</i>  | <i>Avena barbata</i>        |
| <i>Agrostis tenuis</i>       | <i>Avena fatua</i>          |
| <i>Danthonia pilosa</i>      | <i>Avena sativa</i>         |
| <i>Lolium perenne</i>        | <i>Briza maxima</i>         |
|                              | <i>Briza minor</i>          |
| Native annuals               | <i>Bromus mollis</i>        |
| <i>Agrostis exigua</i>       | <i>Bromus commutatus</i>    |
| <i>Aristida oligantha</i>    | <i>Bromus racemosus</i>     |
| <i>Bromus trinii</i>         | <i>Bromus rigidus</i>       |
| <i>Eragrostis hypnoides</i>  | <i>Bromus rubens</i>        |
| <i>Eragrostis orcuttiana</i> | <i>Elymus caput-medusae</i> |
| <i>Festuca megalura</i>      | <i>Festuca dertonensis</i>  |
| <i>Festuca microstachys</i>  | <i>Hordeum gussoneanum</i>  |
| <i>Festuca octoflora</i>     | <i>Hordeum murinum</i>      |
| <i>Phalaris angusta</i>      | <i>Lagurus ovatus</i>       |
| <i>Scribneria bolanderi</i>  | <i>Lolium multiflorum</i>   |
|                              | <i>Lolium temulentum</i>    |
|                              | <i>Phalaris canariensis</i> |
|                              | <i>Phalaris caroliniana</i> |
|                              | <i>Phalaris minor</i>       |
|                              | <i>Phalaris paradoxa</i>    |

The following introduced grasses are found in California only as weeds on disturbed or cultivated ground and usually at low elevations:

|                              |                               |
|------------------------------|-------------------------------|
| <i>Aegilops triuncialis</i>  | <i>Eleusine indica</i>        |
| <i>Avena sterilis</i>        | <i>Eragrostis cilianensis</i> |
| <i>Digitaria filiformis</i>  | <i>Eragrostis poaeoides</i>   |
| <i>Digitaria sanguinalis</i> | <i>Setaria lutescens</i>      |
| <i>Echinochloa colonum</i>   | <i>Setaria viridis</i>        |

Southward from San Francisco Bay occur spontaneously a number of introductions of subtropical origin. Sometimes they occur on the open range, in rock crevices, in chaparral, or in oak woods, seemingly without regard to the habitats defined by the native vegetation. Unlike the introductions from the Mediterranean region, which are usually very abundant, make poor forage,

and have harsh awns which cause mechanical injury to stock (for example, *Bromus* spp. and *Hordeum* spp.), these species are generally palatable and not mechanically injurious, but are much less aggressive.

## Perennials

*Andropogon saccharoides*  
*Imperata hookeri*  
*Pennisetum villosum*  
*Sorghum halepense*<sup>5</sup>  
*Tricholaena rosea*

## Annuals

*Lamarckia aurea*  
*Schismus arabicus*  
*Sorghum vulgare*

*Phalaris platensis* Henrard, an Argentine species, is reported to be very similar in characteristics to *P. lemmoni* (Henrard, 1938). The fact that *P. angusta* is considered native to both Chile and California, indicates that the genus is a real link between the floras of the two regions. Among the native California grasses of similar habitat additional links may be found in *Poa secunda* Presl, and *Bromus trinii* Desv. (Beetle, 1943b). This connection between a flora of California and one of temperate South America must be considered to have a history independent of the connections found in both the deserts and the mountains of the same regions (see p. 311 and 316). Campbell (1944) states that "species common to Chile and central southern California . . . may be explained as due to migration along the great mountain system of the Pacific Coast." It seems unlikely that the variously adapted species of grassland, desert, and mountain, all took the same route; thus desert species may have traveled a dry continental shelf similar to Lower California.

The genus *Panicum* (in number of described species one of the largest of all genera) as a whole and especially the subgenus *Dichanthelium* whose center of distribution is the coastal plain of the southeastern United States (Hitchcock and Chase, 1910), has found the mountains, deserts, and ocean surrounding California difficult barriers to traverse. Three perennial species (*P. scribnerianum*, *P. pacificum*, and *P. occidentale*), none of them common, are found at scattered localities, usually in at least semishade. A fourth, *P. thermale*, is known from the vicinity of the hot springs of Sonoma County. Thus the southeastern United States, recognized as a center of origin and dispersal for many nonboreal plants and animals of the greater part of the eastern United States (Adams, 1902), has made only a minor, occasional, accidental contribution to the flora of California; this holds true for the flora of California in general as well as for the grasses. It is apparent that the ocean to the west, the desert to the south, and the mountains to the east have long been barriers to the immigration of species to California from other floras. Only the north-to-south immigration route along the Coast Ranges and the Sierra Nevada into Oregon and Washington has been responsible for any appreciable contribution to the California grass flora from the Tehachapi Mountains northward.

<sup>5</sup> At first glance, the California ranchers who have had trouble with Johnson grass may well be startled to find it listed as "less aggressive." Reference here is made, however, to the occurrence of these grasses on the dry, open range, a very different habitat from cultivated fields or irrigated pastures. In this connection it is interesting to note the apparently greater drought resistance of the bunch-grass type of perennial over stoloniferous or rhizomatous types. This accounts for the general rarity of introduced perennials on the range, as well as the general prevalence of introduced perennials in lawns and irrigated pastures.

**Grasses of the Alkaline Flats.**—Species of alkaline flats include:

## Native perennials

*Distichlis spicata* var. *nana*  
*Distichlis spicata* var. *stricta*  
*Puccinellia distans*  
*Puccinellia lemmoni*  
*Sporobolus airoides*

## Native annuals

*Hordeum pusillum*  
*Leptochloa fascicularis*  
*Puccinellia simplex*

The presence of any of the above is an indication of alkaline soil though *Distichlis spicata* var. *stricta* and *Puccinellia simplex* have the strongest tolerance. Alkaline soils are characterized by poor drainage, high evaporation, and an accumulation of soluble salts. These are not necessarily salts of alkaline reaction but include chlorides, sulfates, and carbonates and bicarbonates of sodium, potassium, calcium, and magnesium. Northward in the Great Valley the alkaline soils are all on the west side, but southward they are more common and occur on both sides of the valley.

With the exception of *Leptochloa fascicularis*, the species listed above represent typically north-temperate genera. Judging both from their centers of variation and the location of related species, the species of wide range listed above are probably immigrants to California. These include *Puccinellia distans*, *Hordeum pusillum*, and *Leptochloa fascicularis* (and probably also *Agropyron smithii*, which is of limited occurrence in northeastern California). This does not imply that they are not native in their occurrence, but simply that the impetus which resulted in their formation was elsewhere and that they subsequently found California a favorable environment. The remaining elements, on the other hand—varieties of *Distichlis spicata* and the species of *Puccinellia* which are narrow in range—may have been developed *in situ*. This appears to be particularly true of *Distichlis*. Here the origin of *Distichlis* from *Uniola*, or at least some ancestor with perfect florets, seems evident; dioeciousness is too rare in the Festuceae to be considered primitive where corroborative evidence is wholly lacking. Within *Distichlis* the species of larger plant body (for example, *D. texana*) are more characteristic of the Festuceae and are less specialized as to habitat; they might be regarded as precursors of the halophytic *D. spicata*, which has achieved a particularly wide range because of its adaptation to alkaline habitats.

**Grasses of Aquatic Habitats.**—The true aquatics include:

## Native grasses

*Alopecurus carolinianus*  
*Alopecurus howellii*  
*Alopecurus geniculatus*  
*Alopecurus saccatus*  
*Glyceria leptostachya*  
*Glyceria occidentalis*  
*Glyceria pauciflora*  
*Leersia oryzoides*  
*Paspalum distichum*  
*Phragmites communis*

## Introduced perennials

*Alopecurus pratensis*  
*Arundo donax*  
*Paspalum dilatatum*  
*Paspalum urvillei*  
*Setaria geniculata*

## Introduced annuals

*Agrostis semiverticillata*  
*Echinochloa crusgalli*  
*Polypogon interruptus*  
*Polypogon maritimus*  
*Polypogon monspeliensis*

All of the native perennial aquatics listed are of wide distribution, at least on the North American continent, except *Alopecurus howellii*, *A. saccatus*,

and *Glyceria leptostachya*, which are confined to the northwestern corner of the United States from southern Washington to northern California. Both *Alopecurus* and *Glyceria* are north-temperate genera, and their California distributions fit the picture of predominantly northern origins proposed for species of other habitats.

**Grasses of the Vernal Pools.**—In vernal pools are found :

Introduced annuals

*Crypsis aculeata*

*Heleochoa schoenoides*

Native annuals

*Agrostis microphylla* and its varieties

*Deschampsia danthonioides* var.  
*gracilis*

*Neostaphia colusana*

*Orcuttia californica*

*Orcuttia greenei*

*Orcuttia pilosa*

*Orcuttia tenuis*

*Phalaris lemmoni*

*Pleuropogon californicus*

The vernal pool represents an unusual habitat requiring considerable adaptation on the part of the plants inhabiting it. Water collects in pockets during the rainy season and, because of the high water table and poor drainage, remains as a small lake until late in the spring. When the water finally disappears as a result of evaporation, the annuals germinate (usually at a time when the surrounding vegetation has fully matured). Although their seeds have germinated in an aquatic habitat, these annuals must grow and mature in the hottest summer temperatures on dry, cracked soils. An odorous viscid pubescence reduces their transpiration rate. All are intolerant of salinity. None are perennial.

Both *Deschampsia danthonioides* var. *gracilis* and *Crypsis aculeata* are common; *Phalaris lemmoni* is frequent. The rest of the plants are rare, their habitat being limited and frequently disturbed by cultivation.

*Orcuttia*, *Neostaphia*, and *Pleuropogon*, all related according to Hoover (1941), are the rarest and most unusual genera of California grasses. Obscure as to relationship and highly specialized in habitat, the origin of the species of these three genera represents a difficult and as yet unsolved problem. The relationship between *Orcuttia* and *Neostaphia*, viscid annuals of unpleasant odor, is easier to see than any relationship between these two and *Pleuropogon*. *Pleuropogon*, a genus of five species and one variety (Benson, 1941), in its northern origin through *P. sabinii*, its generally perennial habit, and its aquatic habitat, not to mention the general morphological characters of inflorescence and spikelets, suggests a close relationship to *Glyceria*. The endemic species of *Pleuropogon*, *Orcuttia*, and *Neostaphia* agree with the Great Valley endemics of other families in belonging to distinctly California genera.<sup>6</sup> Hoover considers them to be endemics of recent (that is, derived) origin. They would then be called neoendemics, or initiates. This appears to be borne out by the fact that the valley floor was submerged in recent geological time. On the other hand, the isolated position of these genera (that is, the absence in the flora of close relatives, or links to other more common

<sup>6</sup> Hoover, R. F. Endemism in the flora of the Great Valley of California. A thesis submitted in partial fulfillment of requirements for the degree of Doctor of Philosophy, University of California. 175 p. 1937.

genera) suggests a long history. It may well be that these plants are restricted in range, not because they are very young nor because they are very old, but because in evolution they have become too selective as to habitat.

**Chaparral Grasses.**<sup>7</sup>—The most characteristic grass species of the chaparral belt include:

Southern

*Melica frutescens*  
*Muhlenbergia microsperma*  
*Stipa parishii*  
*Stipa coronata*  
*Stipa lepida*

More northern

*Calamagrostis densa*  
*Agrostis diegoensis*  
*Melica imperfecta*  
*Melica torreyana*

Only in the southern part of the state have grass species very successfully responded to the influence of the chaparral and become so restricted as seldom to be found outside the belt. *Stipa coronata* and *S. lepida* are common and conspicuous elements; the others are somewhat rare. Apparently these species originated in, or near, their present localities, for they have no near relatives in distant floras. According to Clements (1934) the grassland is accomplishing a "gradual encroachment" on the chaparral, but the two vegetational zones have existed "since the last great pulsation of the Pleistocene," as evidenced "by the universal absence of chaparral relicts."

Northward the pure chaparral has no characteristic grass species. Here are found *Calamagrostis densa* (obviously a narrower adaptation of *C. koelerioides*) and occasionally *Melica torreyana* and *Agrostis diegoensis*. Where there are especially dense stands of hard chaparral (for example, pure stands of *Adenostema* or *Arctostaphylos*) not only the grasses but nearly all other herbaceous species are generally forced out by the competition. Where the chaparral tends to be "soft," composed of mixtures of *Arctostaphylos*, *Quercus*, *Rhamnus*, *Ceanothus*, and similar plants, perennial species are frequent, but are species characteristic of oak woods or open ranges, such as *Stipa pulchra*, *S. californica*, *Melica californica*, *Sitanion hystrix*, *Poa secunda*, *Bromus carinatus*, *Elymus glaucus*, and *Koeleria cristata*. Introduced weedy annuals spring up whenever sufficient moisture is available (*Bromus rubens*, *B. mollis*, *Avena barbata*, *Gastridium ventricosum*, and *Aira caryophyllea*) but introduced perennials are absent. Not one of the twenty-one species listed by Sampson (1944) as "common to chaparral areas" is confined in its distribution to that area, but are the characteristic species of the oak woods and open grasslands.

Cooke (1940) in discussing *Stipa californica* in the Mount Shasta area, says: [It is] generally confined to the chaparral but also present throughout the transition zone and scattered through the Canadian zone. If the chaparral should be discussed as a grass association it could be called an *Elymus glaucus-Stipa californica-Bromus carinatus* association.

**Grasses of Coastal Bluffs and Sand Dunes.**—Along the strand the characteristic species include:

Native

*Agropyron arenicola*  
*Agrostis blasdalei*  
*Agrostis californica*

Native (continued)

*Agrostis pallens*  
*Calamagrostis foliosa*  
*Elymus mollis*

<sup>7</sup> For a discussion of the ecology of and distribution of the chaparral belt, see Sampson (1944) and Jepson (1925).

## Native (continued)

*Elymus vancouverensis**Elymus virescens**Poa confinis**Poa douglasii**Poa macrantha**Poa unilateralis*

## Introduced annuals

*Holcus lanatus**Holcus mollis*

## Introduced perennials

*Agropyron junceum**Ammophila arenaria*

Like the endemics of the neighboring bog flora and unlike the endemics of the vernal pools of the Great Valley, these species are probably relics. They reached California a long time ago and have since adapted themselves to only the most specialized habitats. It is perhaps more than a coincidence that *Agrostis blasdalei* and *A. pallens*, occupying isolated coastal situations, should be the only California representatives of their section of *Agrostis*. Other members of the section are found southward in the mountains of Mexico (*A. tolucensis* H.B.K.) and the mountains of Peru (*A. bromoides* Gr.). The species related to *Poa unilateralis* are also mountain inhabiting, but are found well represented in the California Sierra Nevada. Only *Elymus virescens*, most widespread of the three listed and the commonest (California north to Alaska), appears to be of recent origin, its nearest relative being the abundant *E. glaucus* of the Great Valley and the foothills.

Cooper (1936), in a comprehensive survey of the distribution and origin of the strand and dune flora of the Pacific Coast, concludes that *Poa confinis*, *P. douglasii*, *P. macrantha*, and the two dune species of *Elymus* (lumped by Cooper as *E. arenarius*) are, together with other associated species, of "subarctic or north temperate origin." The three *Poa* species—apparently closely related, for all fall into the section *Pratenses*—are related to *P. eminens* Presl of coastal northeastern Asia and arctic America and *P. labradorica* Steud., of the coast of Labrador. *Elymus* was first considered circumboreal in distribution, the North American material often being separated as var. *villosus* (St. John, 1915), but the latest taxonomic thought not only separates the New World material (as *E. mollis*) from the Old World, *E. arenarius*, but further distinguishes on the Pacific Coast *E. vancouverensis* Vasey. Although the origins of *Agrostis pallens* and *Agropyron arenicola* are more obscure, they belong to genera of principally north-temperate development, and they offer no evidence against the general picture of a migration southward of the grass species inhabiting the California sand dunes. Further evidence for the northern origin of the grasses in the dune flora is seen in the near absence of grasses on the strand formation of the southern California coast (McKenny, 1901).

There are only two introduced perennials on the dunes, *Ammophila arenaria* and *Agropyron junceum*, both from seashore sands and dunes of Europe, the former well established over a broader range than any of the native species, the second established only locally.

**Salt-Marsh Grasses.**—The ten species found in salt marshes are:

## Native (all perennials)

*Distichlis spicata* var. *stolonifera**Hordeum nodosum* var. *boreale**Monanthochloe littoralis**Puccinellia nutkaensis**Spartina foliosa*

## Introduced

*Agrostis palustris**Lepturus cylindricus**Lolium strictum**Lolium subulatum**Pholiurus incurvus*

Only the five introduced species, all from the salt marshes of Europe, have a similar background for their appearance in the California flora. It is striking that all but *Agrostis palustris* should be annuals, whereas no native annual grasses appear in the salt marshes. The general poverty of the salt-marsh floras probably attests to their recent formation, which is further evidenced by comparatively recent radical changes in the elevation of the California land mass.

*Distichlis spicata* is of wide occurrence throughout the Americas (Beetle, 1943a). Variety *stolonifera* intergrades with the inland variety *stricta*, and these may be presumed to be closely related. Variety *stolonifera* has neither become completely adapted to the salt-marsh habitat nor lost its connection with the inland flora. This situation is similar to that found in the introduced *Agrostis palustris*, whose affinities with the common pasture grass, *A. alba* (red top), are marked.

The dioeciousness of *Monanthochloe littoralis* is not an indication of relationship with *Distichlis*, in spite of the rare occurrence of the condition in the Festuceae. *Monanthochloe* is a small genus (probably only two species) ranging across the North American subtropics, from whence it came to California.

In contrast is the probable origin of *Puccinellia grandis*, which ranges north to Alaska; there its range coincides with that of at least twelve other related species (Swallen, 1944).

Finally, *Spartina foliosa* is similar to and probably related to *S. alterniflora* of the Atlantic Coast of North America; its method of reaching the California coast is in doubt.

**Grasses of Bogs.**—*Agrostis longiligula* and *Calamagrostis bolanderi* are the only two grasses within this habitat which need special mention. The true coastal bogs, strongly peaty and containing peat moss, have a flora independent of that of the recent bogs of both the Sierra Nevada and the Coast Ranges, whose species are essentially the same as those of the pine woods which surround them. It is likely that intensive search would reveal more bogs in the coastal areas than are now known botanically. At present, knowledge of them comes for the most part from Mendocino County where the largest and most southerly (Rigg, 1933) of the bogs occurs.

Unlike the endemic species of the Great Valley to be considered of recent derivation from typically California genera, *Agrostis longiligula* and *Calamagrostis bolanderi* are representatives of cosmopolitan genera and are possibly relics. Bog societies are more common northward in North America and are closely related to similar societies in Eurasia (Transeau, 1903). According to Transeau:

Present bog habitats are continuations of similar habitats which existed in early post-glacial times, when tundra conditions and tundra vegetation were dominant . . . [and] the temperature phenomena of undrained depressions, containing deposits of peat are favorable to the preservation of these types [that is, relics].

The more recently formed bogs tend to have a characteristic flora consisting of *Calamagrostis bolanderi*, *C. breweri*, *C. inexpansa*, *Glyceria pauciflora*, and *Deschampsia caespitosa*.

## OTHER CALIFORNIA GRASSES

In addition, the grass family is represented in California by the bamboos, cereals, a few lawn grasses (*Opizia stolonifera*, *Axonopus compressus*, *Stenotaphrum secundatum*, and *Eremochloa ophiuroides*), and a few ornamentals (for example, *Cortaderia selloana* and *Ampelodesmos mauritanicus*), which are usually found only under cultivation. A partial list of introduced grasses and their histories is given by Robbins (1940); for additions see Beetle (1943, b).

## CONCLUSION

Some of the foregoing material is available in floras of California and all of it could be. The application of this enlarged focus however, prepares the way for a new viewpoint in conservation work. The practical application of the present study in the field of agronomy centers around efforts to improve rangeland, a natural resource no less important than forest or wildlife, which has been found to require a program of planned management if its value is to be maintained undiminished (Jones and Love, 1945). The natural areas defined by the known ranges of grass species are the areas within which any given program of management will be practical.

In the search for better range grasses, the regions from which introductions might be presumed to be successful could be chosen on the evidence of: (1) incidence of successful introductions in the past; and (2) failure of unsuccessful introductions in the past, regardless of whether they have been brought in by nature before man or subsequently either purposely or accidentally by man. This is the type of evidence that should control the choice of native range plants for selection improvement or the choice of species and areas in reseeding programs.

In the larger and more general ecological picture, the evidence given by the detailed study of the grass family may now be correlated with the known facts concerning the migration of whole floras. Then greater confidence may be placed in the indicator value of grasses in studies of plant succession and range management. In addition, herein is basic evidence bearing on the question of grassland as a climax or subclimax.

## ACKNOWLEDGMENTS

For helpful suggestions regarding details of this paper, the author expresses appreciation to: Dr. L. C. Constance, Mr. J. T. Howell, Dr. R. M. Love, Professor B. A. Madson, Dr. H. L. Mason, Dr. G. L. Stebbins, and Mr. R. Tofsrud.



## BIBLIOGRAPHY

ADAMS, C. C.

1902. Southeastern United States as a center of geographical distribution of flora and fauna. *Biol. Bul.* 3:115-31.

BEAL, W. J.

1891. Geographical distribution of the grasses of North America. *Amer. Assoc. Adv. Sci. Proc.* (1890) 39:312.

BETLE, A. A.

- 1943a. The North American variations of *Distichlis spicata*. *Torrey Bot. Club Bul.* 70: 638-50.  
1943b. Notes on the distribution of California grasses. *Leaflets West. Bot.* 3:258-60.  
1943c. The phytogeography of Patagonia. *Bot. Rev.* 9:667-79.  
1945. A new section *Microphyllae* in *Agrostis*. *Torrey Bot. Club Bul.* 72:541-49.  
1946a. Notes on the distribution of California grasses II. *Leaflets West. Bot.* 4:285-289.  
1946b. A new alpine *Glyceria* from California. *Madroño* 8:160-61.

BENSON, L.

1941. Taxonomic studies. *Amer. Jour. Bot.* 28:358-64.

BOLANDER, G.

1866. Grasses of the state. *California State Agr. Soc. Trans.* 1864-65:131-45.

BOYLE, W. S.

1945. A cyto-taxonomic study of the North American species of *Melica*. *Madroño* 8:1-26.

CAMPBELL, D. H.

1944. Relations of the temperate floras of North and South America. *California Acad. Sci. Proc.* 25:139-46.

CHANEY, R. W.

1934. Redwoods of the past. 7 p. Published by Save-the-Redwoods League, Berkeley, Calif.

CLEMENTS, F. E.

1934. The relict method in dynamic ecology. *Jour. Ecol.* 22:39-68.

CONZATTI, C.

1943. Poaceas. p. 111-220. *In*: Flora taxonomica Mexicana. vol. 2. Monocotiledoneas. 279 p. Porfirio Diaz B., Oaxaca, Juarez, Mex.

COOKE, W. B.

1940. Flora of Mount Shasta. *Amer. Midland Nat.* 23:497-572.  
1941. First supplement to the flora of Mount Shasta. *Amer. Midland Nat.* 26:74-84.

COOPER, W. S.

1922. The broad-sclerophyll vegetation of California: an ecological study of the chaparral and its related communities. *Carnegie Inst. Washington Pub.* 319:1-124.  
1936. Strand and dune flora of the Pacific Coast. p. 141-87. *In*: Goodspeed, T. H. *Essays in geobotany in honor of William Albert Setchell*. xxvi + 320 p. University of California Press, Berkeley, Calif.

DAVY, J. B.

1902. Stock ranges of northwestern California: notes on the grasses and forage plants and range conditions. *U. S. Dept. Agr. Bur. Plant Indus. Bul.* 12:1-81.

DORE, W. G., and A. E. ROLAND.

- 1941-42. The grasses of Nova Scotia. *Nova Scotian Inst. Sci. Proc. and Trans.* 20: 177-288.

EASTWOOD, A.

1901. General botanical features of the Coast Mountains of California. *Science* (n.s.) 14:600.  
1941. The islands of southern California and a list of the recorded plants. *Leaflets West. Bot.* 3:27-36.

FOURNIER, E.

1876. Sur la distribution géographique des Graminées Mexicaines. *Ann. Soc. Nat.* (ser. 6) 9:261-90.

GOULD, F. W.

1945. Notes on the genus *Elymus*. Madroño 8:42-47.

GRAY, A.

1846. Analogy between the flora of Japan and that of the United States. Amer. Jour. Sci. (n.s.) 2:135.  
1858-59. Observations upon the relations of the Japanese flora to that of North America and of other parts of the north temperate zone. Amer. Acad. Arts and Sci. Mem. (ser. 2) 6:377-452.

GRIFFITH, D.

1912. The grama grasses. U. S. Natl. Mus. Contrib. U. S. Natl. Herb. 14:343-428.

HENRARD, J. TH.

1938. Vicarious grasses, as demonstrated by a new species of *Phalaris* from South America. Blumea 3:168-72.

HENDRY, G. W.

1931. The adobe brick as a historical source. Agr. Hist. 5:110-27.

HITCHCOCK, A. S.

1923. Gramineae. (Revised by A. Chase.) p. 72-144. In: Jepson, W. L. (1925).  
1927. The grasses of Ecuador, Peru, and Bolivia. U. S. Natl. Mus. Contrib. U. S. Natl. Herb. 24:291-556. (See specifically p. 397.)  
1935. Manual of the grasses of the United States. U. S. Dept. Agr. Misc. Pub. 200: 1-1040.

HITCHCOCK, A. S., and AGNES CHASE.

1910. The North American species of *Panicum*. U. S. Natl. Mus. Contrib. U. S. Natl. Herb. 15:1-396.  
1920. Revisions of North American grasses. U. S. Natl. Mus. Contrib. U. S. Natl. Herb. 22:1-77. (See specifically p. 45-77.)

HOOVER, R. F.

1941. The genus *Orcuttia*. Torrey Bot. Club Bul. 68:149-56.

HOWELL, J. T.

1946. Notes on the grass family in Marin County, California. Leaflets West. Bot. 4:243-47.

JEPSON, W. L.

1925. A manual of the flowering plants of California. 1,238 p. Associated Students Store, University of California, Berkeley, Calif.

JOHNSON, B. L.

1945. Cyto-taxonomic studies in *Oryzopsis*. Bot. Gaz. 107:1-32.

JOHNSTON, I. M.

1943. Plants of Coahuila, eastern Chihuahua, and adjoining Zacatecas and Durango, II. Jour. Arnold Arb. 24:375-421.

JONES, B., and R. M. LOVE.

1945. Improving California ranges. California Agr. Exp. Cir. 129:1-48.

LAWRENCE, W. E.

1945. Some ecotypic relations of *Deschampsia caespitosa*. Amer. Jour. Bot. 32:298-314.

MCARDLE, R. E., et al.

1936. The white man's toll. U. S. — Cong. Senate Doc. 199:81-116.

McKENNEY, R. E. B.

1901. Notes on plant distribution in southern California, U. S. A. Bot. Centbl. Beihefte. Abt. B, 10:166-76.

PARODI, L. R.

1928. Sinopsis de las Gramineas Argentinas del género *Muhlenbergia*. Physis 9:205-22.  
1937. Contribucion de estudio de las Gramineas de género *Paspalum*. Rev. del Mus. de la Plata (n.s.), Sec. Bot., 1:211-50.  
1941. Revision de las Gramineas Sudamericanas del género *Hierochloe*. Rev. del Mus. de la Plata (n.s.), Sec. Bot., 3:183-212.

PIEMEISEL, R. L., and F. R. LAWSON.

1937. Types of vegetation in the San Joaquin Valley of California and their relation to the beet leafhopper. U. S. Dept. Agr. Tech. Bul. 557:1-28.

PURER, E.

1942. Plant ecology of the coastal salt marshlands of San Diego County, California. Ecol. Mono. 12:81-111.

RIGG, G. B.

1933. Notes on a sphagnum bog at Fort Bragg, California. Science (n.s.) 77:535-36.

ROBBINS, W. W.

1940. Alien plants growing without cultivation in California. California Agr. Exp. Sta. Bul. 637:1-128.

RUSSELL, R. J.

1926. Climates of California. Univ. California Pubs. Geog. 2:73-84.

SAMPSON, A. W.

1944. Plant succession on burned chaparral lands in northern California. California Agr. Exp. Sta. Bul. 685:1-144.

SAMPSON, A. W., and AGNES CHASE.

1927. Range grasses of California. California Agr. Exp. Sta. Bul. 430:1-94. (Out of print.)

SCRIBNER, F. L., and E. D. MERRILL.

1910. The grasses of Alaska. U. S. Natl. Mus. Contrib. U. S. Natl. Herb. 13:47-92.

SHANTZ, H. L., and R. L. PIEMEISEL.

1924. Indicator significance of the natural vegetation of the southwestern desert region. Jour. Agr. Res. 28:721-802.

STEBBINS, G. L., Jr., and R. M. LOVE.

- 1941a. An undescribed species of *Stipa* from California. Madroño 6:137-41.

- 1941b. A cytological study of California forage grasses. Amer. Jour. Bot. 28:371-82.

ST. JOHN, H.

1915. *Elymus arenarius* and its American representatives. Rhodora 17:98-103.

1941. The status of *Poa secunda* and of *Poa sandbergii* Gramineae in North America. In: New and noteworthy northwestern plants, part 8. Amer. Jour. Bot. 28:78-81.

SWALLEN, J. R.

1941. New United States grasses. Washington Acad. Sci. Jour. 31:348-50.

1944. The Alaskan species of *Puccinellia*. Washington Acad. Sci. Jour. 34:16-23.

THURBER, G.

1880. Gramineae. In: Watson, S. Geological survey of California. Botany, vol. 2, p. 253-328.

TRANSEAU, E. N.

1903. On the geographic distribution and ecological relations of the bog plant societies of northern North America. Bot. Gaz. 36:401-20.

WEAVER, J. E., and T. J. FITZPATRICK.

1932. Ecology and relative importance of the dominants of tall-grass prairie. Bot. Gaz. 93:113-50.

# INDEX TO GRASSES

- Aegilops triuncialis* L., 344  
*Agropyron arenicola* Davy (map 153), 341, 348, 349  
     *cristatum* (L.) Gaertn., 312  
     *junceum* (L.) Beauv., 349  
     *repens* (L.) Beauv., 312  
     *smithii* Rydb. (map 143), 316, 346  
     *spicatum* (Pursh) Scribn. and Smith (map 144), 316  
     *trachycaulum* (Link) Malte, 311  
*Agrostis alba* L., 312, 350  
     *ampla* Hitchc., 344  
     *blasdalei* Hitchc. (map 152), 341, 348, 349  
     *bromoides* L., 349  
     *californica* Trin. (map 99), 314, 348  
     *diegoensis* Vasey (map 71), 348  
     *exarata* Trin. (map 17), 311  
     *exigua* Thurb., 344  
     *hiemalis* (Walt.) B.S.P. (map 3), 311  
     *idahoensis* Nash (map 39), 311, 312  
     *longiligula* Hitchc., 350  
     *microphylla* Steud., 347  
     *pallens* Trin., 348, 349  
     *palustris* Huds., 349, 350  
     *retrofracta* Willd., 344  
     *rossae* Vasey (map 43), 312  
     *semiverticillata* (Forsk.) C. Christ., 346  
     *stolonifera* L., 344  
     *tenuis* Sibth., 344  
     *thurberiana* Hitchc. (map 28), 312  
     *tolucensis* H.B.K., 349  
*Aira caryophyllea* L., 348  
     *praecox* L., 312  
*Alopecurus aequalis* Sobol. (map 60), 311, 313  
     *carolinianus* Walt., 346  
     *geniculatus* L. (map 59), 313, 346  
     *howellii* Vasey, 346  
     *pratensis* L., 346  
     *saccatus* Vasey (map 184), 342, 346  
*Ammophila arenaria* (L.) Link (map 161), 341, 349  
*Ampelodesmos mauritanicus* (Poir.) Dur. and Schinz, 351  
*Andropogon glomeratus* (Walt) B.S.P., 341  
     *saccharoides* Swartz, 345  
     *tenuispatheus* Nash, 341  
     *virginicus* L., 341, 344  
*Anthoxanthum odoratum* L., 312  
*Aristida adscensionis* L. (map 118), 315  
     *californica* Thurb. (map 119), 315  
     *capillacea* Lam., 316  
     *divaricata* Humb. and Bonpl. (map 120), 315  
     *fendleriana* Steud. (map 121), 315  
     *glabrata* (Vasey) Hitchc. (map 122), 315  
     *glauca* (Nees) Walp. (map 123), 315  
     *hamulosa* Henr. (map 182), 341, 342, 344  
     *oligantha* Michx. (map 183), 344  
     *parishii* Hitchc. (map 124), 315  
     *purpurea* Nutt. (map 125), 315  
     *ternipes* Cav., 316  
     *wrightii* Nash (map 126), 315  
*Arrhenatherum elatius* (L.) Mert. and Koch, 312  
*Arundo donax* L., 346  
*Avena barbata* Brot., 344, 348  
     *fatua* L., 344  
     *sativa* L., 344  
     *sterilis* L., 344  
*Axonopus compressus* (Swartz) Beauv., 351  
*Beckmannia syzigachne* (Steud.) Fernald (map 13), 311  
*Blepharidachne kingii* (S. Wats.) Hack. (map 127), 316  
*Bouteloua aristoides* (H.B.K.) Griseb. (map 114), 315  
     *barbata* Lag. (map 115), 315  
     *curtipendula* (Michx.) Torr. (map 116), 315, 316  
     *gracilis* (H.B.K.) Lag. (map 100), 315  
     *rothrockii* Vasey (map 117), 315  
*Briza maxima* L., 344  
     *minor* L., 344  
*Bromus carinatus* Hook. and Arn., 344, 348  
     *ciliatus* L. (map 47), 312  
     *commutatus* Schrad. (map 78), 344  
     *grandis* (Shear) Woot. and Standl. (map 73), 313  
     *inermis* Leyss., 312  
     *laevipes* Shear (map 80), 313, 344  
     *mollis* L., 344, 348  
     *oreuttianus* Vasey (map 79), 313  
     *racemosus* L. (map 77), 344  
     *rigidus* Roth, 344  
     *rubens* L., 344, 348  
     *suksdorfii* Vasey (map 45), 312  
     *trinii* Desv., 344, 345  
     *vulgaris* (Hook.) Shear, 313  
*Calamagrostis bolanderi* Thurb., 350  
     *breweri* Thurb. (map 41), 313, 350  
     *canadensis* (Michx.) Beauv. (map 42), 311, 312  
     *crassiglumis* Thurb. (map 95), 314  
     *densa* Vasey, 348  
     *foliosa* Kearney (map 96), 348  
     *inexpansa* A. Gray, 350  
     *koelerioides* Vasey (map 97), 348  
     *nutkaensis* (Presl) Steud. (map 98), 313  
     *rubescens* Buckl. (map 27), 311  
*Cenchrus echinatus* L., 315, 316  
     *pauciflorus* Benth., 315, 316  
*Chloris gayana* Kunth, 315  
     *virgata* Swartz, 315  
*Cinna latifolia* (Trevir.) Griseb. (map 36), 311, 312  
*Cortaderia selloana* (Schult.) Aschers. and Graebn., 351  
*Crypsis aculeata* (L.) Ait., 347  
*Cynodon dactylon* (L.) Pers., 315  
*Dactylis glomerata* L., 312  
*Danthonia californica* Boland. (map 68), 344  
     *intermedia* Vasey (map 51), 312  
     *pilosa* R. Br., 344  
     *unispicata* (Thurb.) Munro (map 34), 313

- Deschampsia atropurpurea* (Wahl.) Scheele (map 26), 311  
*caespitosa* (L.) Beauv. (maps 2, 16, 90), 311, 313, 350  
*danthonioides* (Trin.) Munro (maps 66, 180), 311, 341, 347  
*elongata* (Hook.) Munro, 311  
*Digitaria filiformis* (L.) Koel., 344  
*sanguinalis* (L.) Scop., 344  
*Dissanthelium californicum* (Nutt.) Benth. (map 157), 342  
*Distichlis spicata* (L.) Greene (maps 111, 133, 156, 176), 309, 316, 342, 346, 349, 350  
*texana* (Vasey) Scribn., 346  
*Echinochloa colonum* (L.) Link, 344  
*crusgalli* (L.) Beauv., 346  
*Eleusine indica* (L.) Gaertn., 344  
*Elymus arenarius* (Abrom.) St. John, 349  
*caput-medusae* L., 344  
*cinereus* Scribn. and Merr. (map 136), 316  
*condensatus* Presl (map 170), 341, 344  
*glaucus* Buckl., 344, 348, 349  
*mollis* Trin. (map 149), 341, 348, 349  
*triticoideus* Buckl., 344  
*vancouverensis* Vasey (map 150), 341, 349  
*virescens* Piper (map 151), 349  
*Eragrostis cilianensis* (All.) Link, 344  
*hypnoides* (Lam.) B.S.P. (map 178), 342, 344  
*oreuttiana* Vasey (map 179), 342, 344  
*poaeoides* (L.) Beauv., 344  
*Eremochloa ophiuroides* (Munro) Hack., 351  
*Eriochloa aristata* Vasey, 315  
*gracilis* (Fourn.) Hitchc., 315  
*Festuca californica* Vasey (map 74), 313  
*dertonensis* (All.) Aschers. and Graebn., 344  
*elatior* L., 312  
*elmeri* Scribn. and Merr. (map 64), 313  
*idahoensis* Elmer (map 14), 311  
*kingii* Cassidy (map 128), 316  
*megalura* Nutt., 344  
*microstachys* Nutt., 312, 344  
*myuros* L., 312  
*occidentalis* Hook. (map 21), 311  
*octoflora* Walt., 344  
*ovina* L. (map 55), 311, 313  
*rubra* L. (map 8), 311  
*subulata* Trin. (map 15), 311  
*subuliflora* Scribn. (map 20), 311  
*viridula* Vasey (map 48), 311, 313  
*Gastridium ventricosum* (Gouan) Schinz and Thell., 348  
*Glyceria borealis* (Nash) Batchelder (map 11), 311  
*californica* Beetle (map 57), 313  
*cookei* Swallen (map 89), 310  
*elata* (Nash) Hitchc. (map 4), 311  
*erecta* Hitchc. (map 33), 313  
*leptostachya* Buckl. (map 86), 346, 347  
*occidentalis* (Piper) J. C. Nels. (map 88), 346  
*pauciflora* Presl (map 7), 311, 346, 350  
*striata* (Lam.) Hitchc. (map 22), 311  
*Heleochloa schoenoides* (L.) Host, 347  
*Hierochloa occidentalis* Buckl. (map 91), 313  
*redolens* (Vahl) R. and S., 316  
*Hilaria jamesii* (Torr.) Benth. (map 140), 315, 316  
*rigida* (Thurb.) Benth. (map 104), 315  
*Holcus lanatus* L., 349  
*mollis* L., 349  
*Hordeum gussoneanum* Parl., 344  
*jubatum* L., 315  
*murinum* L., 343, 344  
*nodosum* L., 349  
*pusillum* Nutt., 346  
*Hystrix californica* (Boland.) Kuntze, 314  
*Imperata hookeri* Rupr., 345  
*Koeleria cristata* (L.) Pers. (map 58), 311, 313, 341, 343, 344, 348  
*Lagurus ovatus* L., 344  
*Lamarckia aurea* (L.) Moench, 345  
*Leersia oryzoides* (L.) Swartz, 346  
*Leptochloa fascicularis* (Lam.) A. Gray (map 181), 346  
*filiformis* (Lam.) Beauv. (map 101), 315  
*uninervia* (Presl) Hitchc. and Chase (map 102), 315  
*Lepturus cylindricus* (Willd.) Trin. (map 158), 341, 349  
*Lolium multiflorum* Lam., 343, 344  
*perenne* L., 344  
*strictum* Presl, 349  
*subulatum* Visiani, 349  
*temulentum* L., 344  
*Melica aristata* Thurb. (map 30), 312  
*bubosa* Geyer (map 44), 312  
*californica* Scribn. (map 63), 313, 344, 348  
*frutescens* Scribn. (map 109), 348  
*fugax* Boland. (map 12), 311  
*geyeri* Munro (map 61), 313  
*harfordii* Boland. (map 62), 313  
*imperfecta* Trin. (map 167), 344, 348  
*spectabilis* Scribn. (map 19), 311  
*stricta* Boland. (map 29), 312  
*subulata* (Griseb.) Scribn. (map 9), 311  
*torreyana* Scribn. (map 76), 313, 348  
*Monanthochloa littoralis* Engelm. (map 155), 349, 350  
*Muhlenbergia andina* (Nutt.) Hitchc., 311, 313  
*asperifolia* (Nees and May.) Parodi (map 139), 316  
*californica* Vasey (map 108), 311  
*filiformis* (Thurb.) Rydb. (map 37), 311, 312, 313  
*microsperma* (DC.) Kunth (map 162), 341, 348  
*montana* (Nutt.) Hitchc. (map 53), 313  
*richardsonis* Rydb. (map 40), 311  
*rigens* (Benth.) Hitchc. (map 163), 341, 344  
*Neostaphia colusana* Davy (map 175), 341, 347  
*Opizia stolonifera* Presl, 351  
*Oreuttia californica* Vasey (map 174), 341  
*greenei* Vasey (map 171), 341, 347  
*pilosa* Hoover (map 173), 341, 347  
*tenuis* Hitchc. (map 172), 341, 347

- Oryzopsis hymenoides* (Roem. and Schult.)  
   Ricker (map 138), 316  
   kingii (Boland.) Beal, 313  
   miliacea (L.) Benth. and Hook., 315  
*Panicum miliaceum* L., 315  
   occidentale Scribn. (map 69), 313, 345  
   pacificum Hitchc. and Chase (map 70),  
     313, 344, 345  
   scribnerianum Nash (map 94), 345  
   shastense Scribn. and Merr. (map 93),  
     310  
   thermale Boland, 345  
   urvilleanum Kunth (map 103), 315, 316  
*Paspalum dilatatum* Poir., 346  
   distichum L., 346  
   urvillei Steud., 346  
*Pennisetum villosum* R. Br., 345  
*Phalaris angusta* Nees, 344, 345  
   arundinacea L., 344  
   californica Hook. and Arn. (map 92), 313  
   canariensis L., 344  
   caroliniana Walt., 344  
   lemmoni Vasey, 345, 347  
   minor Retz., 344  
   paradoxa L., 344  
   platensis (Arech.) Parodi, 345  
   tuberosa L., 344  
*Phleum alpinum* L. (map 5), 311  
   pratense L., 312  
*Pholurus incurvus* (L.) Schinz and Thell.  
   (map 159), 341, 349  
*Phragmites communis* Trin. (map 154), 346  
*Pleuropogon californicus* (Nees) Benth.  
   (map 83), 314, 347  
   davyi Benson (map 85), 314  
   oregonus Chase, 314  
   refractus (A. Gray) Benth. (map 84), 314  
   sabinii R. Br., 314, 347  
*Poa ampla* Merr. (map 131), 316  
   annua L., 343  
   atropurpurea Scribn., 311  
   bigelovii Vasey and Scribn. (map 110),  
     315  
   bolanderi Vasey (map 38), 311, 312, 313  
   bulbosa L., 312  
   compressa L., 312  
   confinis Vasey, 349  
   cusickii Vasey (map 10), 311  
   douglasii Nees (map 146), 341, 349  
   eminens Presl, 349  
   epilis Scribn. (map 32), 312  
   fendleriana (Steud.) Vasey (map 129),  
     316  
   gracillima Vasey (map 49)  
   howellii Vasey and Scribn. (map 75), 313  
   labradorica Fernald, 349  
   leptocoma Trin. (map 130), 316  
   longiligula Scribn. and Williams, 311  
   macrantha Vasey (map 145), 341, 349  
   napensis Beetle (map 87), 314  
   nevadensis Vasey (map 132), 316  
   palustris L., 311  
   pratensis L. (map 1), 311, 312  
   pringlei Scribn. (map 31), 312  
   rhizomata Hitchc. (map 65), 313  
   rupicola Nash (map 56), 313  
   secunda Presl, 345, 348  
   suksdorfii Vasey (map 52)  
   unilateralis Scribn. (map 147), 349  
*Polypogon interruptus* Hack., 346  
   maritimus Willd., 346  
   monspeliensis (L.) Desf., 346  
*Puccinellia distans* (L.) Parl. (map 23),  
   311, 346  
   grandis Swallen (map 148), 341, 350  
   lemmoni (Vasey) Scribn. (map 25), 311,  
     346  
   nutkaensis (Presl) Fern. and Weath., 349  
   nuttalliana (Schult.) Hitchc. (map 134),  
     316  
   parishii Hitchc. (map 112), 316  
   simplex Scribn. (map 177), 346  
*Schismus arabicus* Nees, 345  
   barbatus (L.) Chase, 315  
*Scribneria bolanderi* (Thurb.) Hack. (map  
   24), 311, 344  
*Setaria geniculata* (Lam.) Beauv., 346  
   lutescens (Weigel) F. T. Hubb., 344  
   viridis (L.) Beauv., 344  
*Sitanion hystrix* (Nutt.) J. G. Smith, 344,  
   348  
   jubatum J. G. Smith, 344  
*Sorghum halepense* (L.) Pers., 345  
   vulgare Pers., 345  
*Spartina alterniflora* Loisel., 350  
   foliosa Trin. (map 160), 349, 350  
   gracilis Trin. (map 137), 316  
*Sporobolus airoides* (Torr.) Torr., 316, 346  
   argutus (Nees) Kunth, 316  
   contractus Hitchc. (map 106), 316  
   cryptandrus (Torr.) A. Gray, 316  
   microsperma (Lag.) Hitchc., 341  
*Stenotaphrum secundatum* (Walt.) Kuntze,  
   351  
*Stipa californica* Merr. and Davy, 348  
   cernua Stebbins and Love (map 164), 341,  
     344  
   comata Trin. and Rupr., 316  
   coronata Thurb. (map 169), 341, 348  
   lemmoni (Vasey) Scribn. (map 72), 313  
   lepida Hitchc. (map 166), 341, 344, 348  
   parishii Vasey, 348  
   pulchra Hitchc. (map 165), 341, 343, 344,  
     348  
   speciosa Trin. and Rupr. (map 107), 316  
   webberi (Thurb.) Johnston, 313  
   williamsii Scribn. (map 141), 316  
*Tricholaena rosea* Nees, 345  
*Triodia mutica* (Torr.) Scribn. (map 135),  
   316  
   pulchella H.B.K. (map 113), 315  
*Trisetum cernuum* Trin. (maps 6, 18, 54),  
   311, 313  
   spicatum (L.) Richt. (map 35), 311, 312  
   wolffii Vasey (map 50), 313