

# HILGARDIA

*A Journal of Agricultural Science Published by  
the California Agricultural Experiment Station*

---

VOLUME 19

APRIL, 1949

NUMBER 4

---

**LACTOBACILLUS TRICHODES NOV. SPEC.,  
A BACTERIUM CAUSING SPOILAGE  
IN APPETIZER AND DESSERT WINES**

**JOHN C. M. FORNACHON, HOWARD C. DOUGLAS,  
and REESE H. VAUGHN**

**THE TAXONOMY OF LACTOBACILLUS  
HILGARDII AND RELATED  
HETEROFERMENTATIVE LACTOBACILLI**

**REESE H. VAUGHN, HOWARD C. DOUGLAS,  
and JOHN C. M. FORNACHON**

**UNIVERSITY OF CALIFORNIA • BERKELEY, CALIFORNIA**



## THE PAPERS IN THIS ISSUE

	PAGE
<i>Lactobacillus trichodes</i> nov. spec., a Bacterium Causing Spoilage in Appetizer and Dessert Wines . . . . .	129

A new species of *Lactobacillus* which has been observed in wines of high alcohol content for many years is described and named. The name *L. trichodes* is proposed because of the marked tendency to grow as long intertwined chains and filaments resembling a mass of hair.

Taxonomy of <i>Lactobacillus hilgardii</i> and Related Heterofermentative Lactobacilli . . . . .	133
---	-----

*Lactobacillus hilgardii* is redescribed and compared with other heterofermentative lactobacilli. The criteria of optimum and maximum temperature relations, fermentation of pentoses and other carbohydrates, rate of growth, type of growth, growth requirements, cell morphology, and the effect of pH on fermentation indicate that this and other "inactive" lactobacilli—*L. fructivorans*, *L. trichodes*, and *Betabacterium caucasicum*—are well-defined species. The name *Lactobacillus desidiosus* is proposed to replace *Betabacterium caucasicum*, in order to conform to accepted nomenclature.

### ERRATUM:

In *Hilgardia* Volume 19, Number 4, the first five lines under "Distinguishing Characteristics," page 135, should read:

The characteristics of *Lactobacillus hilgardii* which differentiate it from the other heterofermentative species investigated are shown in table 1. It ferments *d*-xylose, galactose, sucrose, maltose, and *l*-malic acid; of these, *L. fructivorans* ferments only *l*-malic acid (weakly), *Betabacterium caucasicum* only galactose (weakly), and *L. trichodes* only sucrose and maltose (both weakly).

# THE TAXONOMY OF *LACTOBACILLUS HILGARDII* AND RELATED HETEROFERMENTATIVE *LACTOBACILLI*<sup>1</sup>

REESE H. VAUGHN,<sup>2</sup> HOWARD C. DOUGLAS,  
and JOHN C. M. FORNACHON<sup>3</sup>

SOME heterofermentative bacteria of the genus *Lactobacillus* differ from the better-known species in that they decompose fewer organic compounds, are more difficult to cultivate, and grow more slowly. Pederson (1938)<sup>4</sup> has referred to these lactobacilli as the "inactive" group. Species which may be included in this group are *Betabacterium caucasicum* Orla-Jensen (1919), *Lactobacillus fructivorans* Charlton, Nelson, and Werkman (1934), and *Lactobacillus hilgardii* Douglas and Cruess (1936). Certain strains of other species reported by Pederson (1938) and the two groups described by Fornachon (1943) as *Lactobacillus* Type I (named *Lactobacillus trichodes* by Fornachon, Douglas, and Vaughn, 1949) and *Lactobacillus* Type II also belong to the inactive group. Their taxonomy is confused.

There are some obvious reasons for the taxonomic confusion. The inactive group as a whole is not well known, either because the bacteria are not prevalent in nature or because their more exacting growth characteristics do not favor isolation from sources containing other lactobacilli having less fastidious requirements. Furthermore, the original descriptions of some of the species are not adequate to allow for comparison with previously described species. Regardless of the manner in which the taxonomy of this "inactive" group is finally treated, it is fundamental that the individual species comprising the group must have adequate descriptions. The following information is presented to establish the identity of *Lactobacillus hilgardii*, for which an adequate description is lacking. (See Pederson, 1939, 1948.) With this identity established, the taxonomy of the "inactive" group of lactobacilli is discussed.

## CHARACTERISTICS OF *LACTOBACILLUS HILGARDII*

Since the original type culture of *Lactobacillus hilgardii* had been lost, it was necessary to isolate new cultures similar to or identical with the original strain. The new isolates were obtained from California table wines. The ten cultures are identical with or closely resemble the type species, as will be seen in the following description.

The new isolates of *Lactobacillus hilgardii* were compared with the type culture of *L. fructivorans*, originally supplied by Pederson; two strains of *Betabacterium caucasicum*, one isolated by Vaughn and one received from

<sup>1</sup> Received for publication January 6, 1949.

<sup>2</sup> Associate Professor of Food Technology and Associate Bacteriologist in the California Agricultural Experiment Station.

<sup>3</sup> Assistant Professor of Microbiology, University of Washington, Seattle, Washington. Formerly Research Assistant in Fruit Products, California Agricultural Experiment Station.

<sup>4</sup> Officer, Council for Scientific and Industrial Research, Australia. Formerly Associate in the California Agricultural Experiment Station.

<sup>5</sup> See "Literature Cited" for citations, referred to in the text by author and date.

Tittsler; five new isolates of *L. trichodes*; and ten cultures of *L. brevis*.<sup>6</sup> The cultures of the last two species were isolated from California wines. In addition, comparisons were also made with 30 cultures of *L. brevis* obtained from fermenting olive brines.

## NEW DESCRIPTION OF LACTOBACILLUS HILGARDII (DOUGLAS AND CRUESS, 1936)

### Morphology (at 30° C)

*Form and arrangement*: Rods with rounded ends which occur singly, in short chains and frequently in long filaments. Individual cells in stained preparations measure from 0.5 to 0.8 by 2.0 to 4.0 microns. Individual filaments may measure as much as 15 or more microns in length.

*Spore formation*: Spore formation has not been observed.

*Motility*: Neither motile cells nor flagella have been observed.

*Staining reactions*: The cells are Gram-positive in young cultures. In older cultures the cells may become Gram-negative and show granulation.

### Cultural Characteristics (at 30° C)

*Nutrient agar slant*: Growth, if any, faint.

*Tryptone yeast extract glucose agar, tryptone tomato juice glucose agar, or yeast infusion glucose agar slants*: Growth moderate, raised, hyaline to white and cream-colored with entire edges.

*Nutrient broth*. Growth, if any, faint.

*Tryptone yeast extract glucose broth, tryptone tomato juice glucose broth, or yeast infusion glucose broth*: Moderate turbidity is formed in 3 to 4 days, which clears after a few more days and leaves a granular, flocculent sediment. No pellicle or ring of growth is observed.

*Gelatine stab (at 20° C)*: Growth, if any, faint. No liquefaction after 60 days' incubation.

*Litmus milk*: No change.

### Biochemical Characteristics (at 30° C)

*Catalase*: This enzyme is not produced.

*Fermentation of glucose*: The principal end-products are inactive lactic acid, acetic acid, carbon dioxide, ethyl alcohol, and glycerol.

*Fermentation of fructose*: The fermentation of fructose is characterized by the formation of mannitol in addition to the products mentioned for glucose.

*Fermentation of d-xylose*: The end-products of *d*-xylose fermentation are inactive lactic acid and acetic acid.

*Utilization of carbohydrates*: Only *d*-xylose and fructose are consistently fermented in suitable basal media (tryptone or yeast infusion broths) adjusted to pH 6.8 to 7.0. The results with glucose and *l*-malic acid are erratic. All other sugars, alcohols, and organic acids tested in the same media with the same initial pH range are not attacked. (The basal media and substrate solu-

<sup>6</sup> The assistance of Mr. J. Richard Gililand in isolating some of these cultures is gratefully acknowledged. We also express our indebtedness to Dr. Carl S. Pederson and Dr. Ralph P. Tittsler for cultures, and to Dr. R. E. Buchanan for assistance with nomenclature.



tions were sterilized separately and mixed aseptically just prior to inoculation, or mixed and sterilized by filtration.)

When, however, the same compounds are tested in the same basal media adjusted to an initial pH in the range 4.5 to 5.5, *d*-xylose, glucose, fructose, galactose, maltose, lactose, sucrose, and raffinose are utilized, as indicated by significant acid production from these substrates. In addition, citric and malic acids are always decomposed. Utilization of other sugars, alcohols, glucosides, or organic acids tested was not detected, which is in agreement with the work of Fornachon, Douglas, and Vaughn (1940).

Neither indole nor hydrogen sulfide were produced, nor was nitrate reduced in any of the media used, although glucose was added to insure growth of the cultures.

### Requirements for Growth

*Oxygen relations:* Facultative. Initial isolation and culture is facilitated by the presence of carbon dioxide.

*Temperature requirements:* The optimum temperature for maximum acid formation from glucose in 7 days ranges from 28° to 34° C. The minimum temperature for growth in glucose media is approximately 15° and the maximum temperature is in the range of 40° to 43°.

*pH range:* The optimum initial pH value is variable. For maximum cell production and decomposition of glucose by growing cultures, the optimum initial pH range is from 4.5 to 5.5. For maximum decomposition of citric and malic acids, the optimum initial pH range is between 4.5 and 5.0. Growth has been observed in media with initial pH values of 3.8 and 8.0.

*Alcohol tolerance:* The limiting concentration of alcohol for growth of cultures in wine is between 15 and 18 per cent by volume.

*Salt tolerance:* The limiting concentration of salt (sodium chloride) for growth of cultures in suitable media is between 5 and 6 per cent (grams NaCl per 100 ml of medium).

*Suitable media:* The best medium for growth and maintenance of cultures of *Lactobacillus hilgardii* is liver infusion broth with liver particles. Cultures grown in this medium and held at 0° C remain viable for at least one year. Liver infusion broth with or without liver particles probably is best for preparation of cultures to be used as inocula. For growth of isolated colonies or mass cultures, liver infusion agar, tryptone glucose agar with tomato juice, or yeast infusion or autolysate glucose agar are satisfactory.

### Distinguishing Characteristics

The characteristics of *Lactobacillus hilgardii* which differentiate it from the *d*-xylose, galactose, sucrose, maltose, and *l*-malic acid; of these, *L. fructivorans* ferments only *l*-malic acid (weakly), *Betabacterium caucasicum* only galactose (weakly), and *L. trichodes* only sucrose and maltose (both weakly). *L. hilgardii* does not ferment *l*-arabinose or mannose, whereas *L. brevis* ferments them strongly and *B. caucasicum* ferments mannose weakly. Morphological differences also are useful for distinguishing *L. hilgardii* from *L. brevis* and *B. caucasicum*. It will be noted that *L. hilgardii* and *Lactobacillus* Type II are similar. We believe that they probably are identical species.

Table 1  
DIFFERENTIATION OF "INACTIVE" GAS-FORMING (HETEROFERMENTATIVE) SPECIES OF LACTOBACILLUS

Species and number of cultures	Morphology		Filamentation	Optimum conditions for glucose fermentation		Fermentation* of												Alcohol tolerance,† per cent	
				Temperature range, °C	Initial pH range														
	Cell size, μ	Cell arrangement		Frequent	30-35	4.5-5.5	d-Xylose	l-Arabinose	Glucose	Fructose	Mannose	Galactose	Sucrose	Maltose	Lactose	Raffinose	Glycerol		Mannitol
<i>L. hilgardii</i> , 10 cultures.....	0.5-0.8 × 2.0-4.0	Single and short chains	Frequent	30-35	4.5-5.5	++	-	±	++	-	+	+	+	±	±	-	-	++	±
<i>L. fructivorans</i> , 1 culture.....	0.5-0.8 × 1.5-4.0	Chains of rods	Frequent	25-30	4.5-5.5	-	-	+	++	-	-	-	-	-	-	-	-	±	-
<i>L. trichodes</i> , 5 cultures.....	0.4-0.6 × 2.0-4.0	Long chains	Marked; intertwining	25-30	4.5-5.5	-	-	++	++	-	-	±	-	-	-	-	-	-	-
<i>Lactobacillus</i> , Type II,† 4 cultures.....	0.6-0.8 × 2.0-4.0	Single, pairs, and chains	Frequent	30-35	4.5-5.5	++	-	++	++	-	+	+	+	-	-	-	-	+	±
<i>Betabacterium caucasicum</i> , 2 cultures.....	0.7-1.0 × 2.0-4.0	Single rods	Frequent	25-30	5.0-7.0	-	++	++	++	±	++	-	-	-	-	-	-	-	-
<i>L. brevis</i> , 40 cultures.....	0.7-1.0 × 2.0-4.0	Single and pairs	Occasional	30-35	4.5-7.5	++	++	++	++	+	++	+	++	+	±	±	±	++	++

\* — No activity; ±, weak activity; + moderate activity; ++, maximum activity. All fermentations were conducted at the optimum initial pH range for each culture.  
† Alcohol tolerance (per cent by volume) on basis of growth in wine with 1 ml yeast autolysate per 100 ml of wine.  
‡ Data for *Lactobacillus* Type II taken from Fornachon (1943).

## TAXONOMIC RELATIONSHIPS OF SPECIES OF THE "INACTIVE" GROUP

The description of *Lactobacillus hilgardii* makes it possible to consider the relationships of the "inactive" species to each other and to other species which have been described in the literature. Pederson (1938) considered *Betabacterium caucasicum*, *L. fructivorans*, and "a culture labeled *Bacterium mannitopeum*" as resembling the *L. brevis* group because the cultures had similar optimum temperatures and fermented *l*-arabinose. Although Pederson indicated the need for further study, this raises the question of whether *L. hilgardii* also resembles the *L. brevis* group and whether the inactive cultures investigated here are merely strains of *L. brevis*. Moreover, Orla-Jensen *et al.* (1947) have recently expressed their willingness to make the name *Betabacterium caucasicum* a synonym of *Betabacterium pentoaceticum*. We contend, however, that there are several well-defined species of "inactive" heterofermentative lactobacilli.

The fermentation of pentoses and optimum temperature relations are used as primary characters for separation of the heterofermentative species of *Lactobacillus*. If these are valid criteria for taxonomic differentiation, as maintained by Orla-Jensen (1919, 1943) and Pederson (1938, 1939, 1948), then none of the "inactive" species are more closely related to one another or to *L. brevis* than are the other commonly recognized species *L. buchneri*, *L. fermenti*, and *L. pastorianus* to one another or to *L. brevis*.

It is true that the "inactive" species do resemble *Lactobacillus brevis* when optimum and maximum temperature relations are considered. (They all have optimum temperatures ranging between 25° and 35° C and maximum temperatures of about 40° C or even less.) However, separation of the "inactive" species from *L. brevis* is not difficult when pentose and other carbohydrate fermentations are considered, as has been shown in table 1. Other criteria which serve to separate the "inactive" species from *L. brevis* include differences in rate of growth, type of growth, growth requirements, cell morphology, and the effect of pH on fermentation (table 1).

The general growth characteristics of *Lactobacillus hilgardii*, *L. fructivorans*, *L. trichodes*, and *Betabacterium caucasicum* as compared with *L. brevis* are important for differential purposes. The "inactive" species all grow slowly even under optimum conditions. Visible signs of growth appear only after from 4 or 5 days to as long as 2 weeks. These species also tend to produce a flocculent sediment composed of chains of cells and filaments in the depths of liquid cultures, so that the supernatant liquid is left permanently clear or almost clear for some time. Cultures of *L. brevis*, on the other hand, produce abundant growth in 2 days or less; quickly grow throughout the liquid; and produce a minimum of filaments.

The "inactive" species also are much more exacting in their requirements for growth. Initial isolation and culture is facilitated by the presence of carbon dioxide; otherwise purification and maintenance of cultures is difficult. Growth of all the "inactive" species is stimulated markedly by the addition of tomato juice or yeast autolysate to the culture media; yeast autolysate is



required for growth of *Lactobacillus trichodes*. *L. brevis* also responds to these conditions but less markedly, and the conditions are not mandatory.

The effect of the initial pH value of the medium on growth and fermentative activity, if properly assessed, also serves to differentiate the "inactive" species from the *Lactobacillus brevis* group. This effect has been previously stressed (Fornachon, Douglas, and Vaughn, 1940).

*Lactobacillus hilgardii*, *L. fructivorans*, and *L. trichodes* bear some physiological resemblance to *Bacterium gracile* (*L. gracilis*) Müller-Thurgau (1908). However, on the basis of morphological study of available cultures, Charlton, Nelson, and Werkman (1934) concluded that *Bacterium gracile* was a member of the genus *Leuconostoc* rather than *Lactobacillus*.<sup>7</sup> Pederson (1939) expressed a similar opinion. If this is true, there is obviously no possibility that *L. hilgardii*, *L. fructivorans*, or *L. trichodes* corresponds with *Bacterium gracile*. We believe therefore that *L. hilgardii*,<sup>8</sup> *L. fructivorans*, *L. trichodes*, and *Betabacterium caucasicum* are well-defined species of heterofermentative lactobacilli.

We propose to transfer the species *Betabacterium caucasicum* to the genus *Lactobacillus* because the latter is more widely accepted as the genus name for the heterofermentative lactobacilli than *Betabacterium*, in this country at least. The specific name *caucasicus* is preëempted for the type species of *Lactobacillus*. Hence the name *Lactobacillus desidiosus* (from Latin *desidiosus*, inactive, indolent) is proposed to replace the name *Betabacterium caucasicum*.

Synonyms of *Lactobacillus desidiosus* are *Betabacterium caucasicum* Orla-Jensen, The Lactic Acid Bacteria, 1919, 175; and *Betabacterium pentoaceticum* Orla-Jensen, Orla-Jensen and Kjaer, Antonie van Leeuwenhoek, 12, 1947, 112; the latter in part only. *Lactobacillus pentoaceticus* Fred, Peterson, and Davenport, J. Biol. Chem., 39, 1919, 358; Peterson and Fred, J. Biol. Chem., 41, 1920, 431; and Fred, Peterson, and Anderson, J. Biol. Chem., 48, 1921, 385, is described as actively fermenting both arabinose and xylose. It is therefore considered a synonym of *Lactobacillus brevis* (Orla-Jensen) Bergey *et al.*, by Pederson.

<sup>7</sup> Credence for this conclusion is strengthened by comparison of photographs published by Müller-Thurgau (1908) and Müller-Thurgau and Osterwalder (1913, 1918) with those of Charlton, Nelson, and Werkman (1934).

<sup>8</sup> It has been claimed through an error (Cruess, 1943) that Vaughn and Douglas considered *L. hilgardii* to be very similar to *L. plantarum*. The fallacy has persisted (Cruess, 1947; Olsen, 1948). The heterofermentative nature of *L. hilgardii* was obvious in the first incomplete description and in Fornachon, Douglas, and Vaughn (1940).

## LITERATURE CITED

CHARLTON, D. B., M. E. NELSON, and C. H. WERKMAN.

1934. Physiology of *Lactobacillus fructivorans* sp. nov. isolated from spoiled salad dressing. Iowa State Col. Jour. Sci. 9:1-11.

CRUESS, W. V.

1943. The role of microörganisms and enzymes in wine making. *Advances in Enzymology* 3:349-86.

1947. The principles and practice of wine making. 2d ed. 476 p. Avi Pub. Co., New York, N.Y.

DOUGLAS, H. C., and W. V. CRUESS.

1936. A *Lactobacillus* from California wine: *Lactobacillus hilgardii*. *Food Research* 1:113-19.

FORNACHON, J. C. M.

1943. Bacterial spoilage of fortified wines. 126 p. Australian Wine Board, Adelaide.

FORNACHON, J. C. M., HOWARD C. DOUGLAS, and REESE H. VAUGHN.

1940. The pH requirements of some heterofermentative species of *Lactobacillus*. *Jour. Bact.* 40:649-55.

1949. *Lactobacillus trichodes* nov. spec., a bacterium causing spoilage in appetizer and dessert wines. *Hilgardia* 19(4):129-32.

FRED, E. B., W. H. PETERSON, and AUDREY DAVENPORT.

1919. Acid fermentation of xylose. *Jour. Biol. Chem.* 39:347-84.

FRED, E. B., W. H. PETERSON, and J. A. ANDERSON.

1921. The characteristics of certain pentose-destroying bacteria, especially as concerns their action on arabinose and xylose. *Jour. Biol. Chem.* 48:385-412.

MÜLLER-THURGAU, H.

1908. Bakterienblasen (Bacteriocysten) *Centr. Bakt., II*, 20:353-400, 449-68.

MÜLLER-THURGAU, H., and A. OSTERWALDER.

1913. Die Bakterien im Wein und Obstwein und die dadurch verursachten Veränderungen. *Centr. Bakt., II*, 36:129-338.

1918. Weitere Beiträge zur Kenntnis der Mannitbakterien im Wein. *Centr. Bakt. II*, 48:1-35.

OLSEN, ERIK.

1948. Studies of bacteria in Danish fruit wines. *Antonie van Leeuwenhoek* 14:1-28.

ORLA-JENSEN, S.

1919. The lactic acid bacteria D. klg. D. *Vidensk., selskab skr. naturv. og math. A fel. series* 8, 5:81-196.

1943. The lactic acid bacteria. Die echten Milchsäurebakterien. *Ergänzungsband. D. klg. D. Vidensk., selskab biol. skr.* 2(3):1-145.

ORLA-JENSEN, S., ANNA D. ORLA-JENSEN, and AGNETE KJAER.

1947. On the ensiling of lucerne by means of lactic acid fermentation. *Antonie van Leeuwenhoek* 12:97-114.

PEDERSON, C. S.

1938. The gas-producing species of the genus *Lactobacillus*. *Jour. Bact.* 35:95-108.

1939. *Bergey's manual of determinative bacteriology*. 5th ed., p. 362-78. Williams and Wilkins Co., Baltimore, Md.

1948. *Bergey's manual of determinative bacteriology*. 6th ed., p. 349-63. Williams and Wilkins Co., Baltimore, Md.

PETERSON, W. H., and E. B. FRED.

1920. Fermentation of fructose by *Lactobacillus pentoaceticus*, n. sp. *Jour. Biol. Chem.* 41:431-50.





The journal *Hilgardia* is published at irregular intervals, in volumes of about 600 pages. The number of issues per volume varies.

Subscriptions are not sold. The periodical is sent as published only to libraries, or to institutions in foreign countries having publications to offer in exchange.

You may obtain a single copy of any issue free, as long as the supply lasts; please request by volume and issue number from:

Publications Office  
College of Agriculture  
Berkeley 4, California

The limit to nonresidents of California is 10 separate issues on a single order. A list of the issues still available will be sent on request.

