

**DETERMINING THE PRIMARY DRIVERS OF LIKING TO PREDICT CONSUMER  
ACCEPTANCE OF FRESH NECTARINES AND PEACHES**

**DRIVERS OF LIKING OF FRESH NECTARINES AND PEACHES**

Claudia Delgado<sup>a\*</sup>, Gayle M. Crisosto<sup>a</sup>, Hildegard Heymann<sup>b</sup>, and Carlos H Crisosto<sup>a</sup>

<sup>a</sup> Department of Plant Sciences, University of California, Davis, CA 95616, USA

<sup>b</sup> Department of Viticulture and Enology, University of California, Davis, CA 95616, USA

**Corresponding author:**

\*Claudia Delgado,

University of California, Davis,

Department of Plant Sciences

One Shields Avenue, Davis, CA 95616

Email: [cdelgado@ucdavis.edu](mailto:cdelgado@ucdavis.edu)

Tel.: +1 530-752 9356 Fax: +1 530 752 8502

New address:

Chromocell Corporation

685 U.S. Highway One, North Brunswick, NJ 08902

Email: [Claudia.Delgado@Chromocell.com](mailto:Claudia.Delgado@Chromocell.com)

Tel.: +1 732-565-1113 ext. 7418 Fax: +1 732-565-1183

## 1 **ABSTRACT**

2 A generic descriptive analysis using 11 judges provided 16 sensory attributes that described the  
3 aroma, flavor, and texture characteristics of seven nectarine and peach cultivars selected for their  
4 predominant sensory attributes. Simultaneously, the ‘in-store’ acceptability of these cultivars  
5 was evaluated by 120 consumers from northern California. The relationships among  
6 instrumental measurements (flesh firmness, ripe soluble solids concentration\_(RSSC), and ripe  
7 titratable acidity (RTA)), sensory panel descriptors, and consumer hedonic responses were  
8 studied. In these cultivars, RSSC was the only instrumental measurement significantly related to  
9 overall liking. Cultivars with medium acidity and/or flavor-aroma were liked “very much”, and  
10 consumer willingness to pay more was correlated with overall liking without regard to cultivar.  
11 External preference mapping revealed three clusters that were associated with ethnicity and  
12 consumer preferences within each cluster. Sweetness was the main driver of liking for two  
13 consumer clusters; however, for the third cluster, the perception of fruit aromas described as  
14 grassy/green fruit and pit aromas were the main drivers of liking. There was a high correlation  
15 between instrumental measurements and their sensory perception; however, the sensory attribute  
16 measurements explained cultivar characteristics better than instrumental measurements alone.  
17 Sweetness correlated positively with overall liking and consumer acceptance.

## 18 **PRACTICAL APPLICATIONS:**

19 The main objective of this study was to identify drivers of liking for fresh peaches and nectarines  
20 in order to understand consumer preferences for these fruits. This information can be used by  
21 postharvest researchers to evaluate the potential of new postharvest technologies and consumer

22 acceptance and for plant breeders to develop new cultivars with desirable sensory attributes  
23 driven by the consumer.

24

25 **KEYWORDS:**

26 Instrumental quality; Descriptive analysis; Fruit sensory attributes; Consumer acceptance;

27 External preference mapping, L-PLS regression.

## 28 INTRODUCTION

29         Despite increasing knowledge of the health benefits of eating fruits and vegetables,  
30 consumption of some tree fruit commodities in the United States, such as peaches, nectarines,  
31 and plums, has been static. Since 1980, consumption has averaged 5.5 pounds per capita per  
32 year (USDA Economic Research Data, 2009). A recent consumer quality survey of 1,552  
33 consumers (Sterling-Rice Group, 2006) corroborates previous survey results (Bruhn 1995)  
34 concluding that lack of flavor and chilling injury symptoms are still the main barriers restricting  
35 California peach, nectarine, and plum purchasing in the U.S. market. Some researchers have  
36 tried to predict tree fruit consumer acceptance and/or preference using fruit physicochemical  
37 quality measurements at harvest, such as soluble solids concentration (SSC) for overall  
38 sweetness, penetration force for firmness-texture, and titratable acidity (TA) for sourness (Crisosto  
39 *et al.* 2003). A similar approach was used in Italy (Esti *et al.* 1997) and Slovenia (Colaric *et al.*  
40 2005) to predict consumer quality. However, most of these studies did not attempt to relate these  
41 measurements to consumer responses. Other groups have attempted to evaluate the efficacy of  
42 such physicochemical measurements to explain consumer responses to apples (Hoehn *et al.*  
43 2003) pears (Predieri and Gatti 2009), pineapples (Schulbach *et al.* 2007), mangos (Malundo *et*  
44 *al.* 2001), blueberries (Saftner *et al.* 2008), and oranges (Obenland *et al.* 2009). In some  
45 instances, fruit physicochemical measurements were related to consumers' hedonic ratings or  
46 acceptance percentages (Crisosto, Crisosto and Bowerman 2003; Crisosto and Crisosto 2001;  
47 Crisosto and Crisosto 2005; Crisosto *et al.* 2004; Guerra *et al.* 2009; Gunness *et al.* 2009).  
48 Despite general agreement that measured soluble sugars and/or organic acid concentrations are  
49 key components in predicting consumer acceptability of fresh fruits, other fruit quality  
50 characteristics also affect liking. In peaches, fruit firmness, color, and aroma were important

51 characteristics consumers used to evaluate fruit quality when selecting fruit to purchase (Bruhn  
52 1995; Bruhn *et al.* 1991). These physicochemical measurements and sensory techniques can be  
53 used by the industry to evaluate the potential effect of new postharvest technologies on consumer  
54 acceptance; by shippers to evaluate current postharvest practices; by retail managers to validate  
55 their handling practices; and by plant breeders to develop new cultivars with desirable sensory  
56 attributes. Although significant correlations have been reported between physicochemical  
57 parameters measured instrumentally and sensory properties/hedonic scores (Colaric *et al.* 2005;  
58 Crisosto and Crisosto 2005; Rossiter *et al.* 2000), these correlations usually do not predict  
59 consumer behavior well. Despite the expense of consumer tests, they are more effective in  
60 predicting consumer behavior (Bett 2002; Harker *et al.* 2008; Saftner *et al.* 2008). In an ideal  
61 situation, researchers should simultaneously use physical instrumental fruit quality  
62 measurements and sensory methodology to evaluate consumer responses; however, because of  
63 budget constraints, fast rotation, intensive preparation, and/or limited quantities of fruit available,  
64 it is not always possible to conduct sensory evaluations. Descriptive analysis will provide with  
65 the characterization of the most important attributes for the fruit cultivars, while consumer tests  
66 will indicate how much those fruit cultivars are liked. External preference mapping is a well  
67 known technique used in the sensory field to relate sensory and consumer data with the purpose  
68 of identifying the drivers of liking (Lawless and Heymann, 2010, Yenken *et al.* 2011). L-PLS  
69 analysis is a relatively new technique based on partial least square regression (PLS) used to  
70 characterize three different datasets: the sensory attributes [X], consumers liking [Y] and  
71 consumers information [Z] to identify demographic differences among consumers (Lengard and  
72 Kermit, 2006).

73           Several researchers have explored the relationship between instrumental measurements,  
74 sensory properties, and consumer perception in other fruit commodities for example: apples  
75 (Dailliant-Spinnler *et al.* 1996; Harker *et al.* 2003; Harker *et al.* 2008; Harker *et al.* 2002; Kühn  
76 and Thybo 2001; Oraguzie *et al.* 2009), tomatoes (Causse *et al.* 2010; Lee *et al.* 1999; Sinesio *et*  
77 *al.* 2010), and strawberries (Ares *et al.* 2009). We believe that the recent release and marketing  
78 of tree fruit cultivars with different flavors and the establishment of ripening protocols (Crisosto  
79 1999) justify the expense of developing drivers of liking for tree fruit. Thus, the main goal of  
80 this study was to identify drivers of liking for fresh nectarines and peaches that predict consumer  
81 acceptance and/or preferences for these fruits during postharvest handling.

82

## 83 **MATERIALS AND METHODS**

### 84 **Cultivar selection and fruit preparation**

85           Seven peach and nectarine cultivars were selected for this study for their commercial  
86 importance, differences in titratable acidity, flavor, and aroma, and similar melting flesh texture  
87 after ripening (Table 1). ‘August Pearl’ is a low acid, white flesh nectarine; ‘Fire Sweet’, a  
88 medium acid, flavorful, yellow flesh nectarine; ‘August Bright’, a high acid, yellow flesh  
89 nectarine; ‘Autumn Snow’, a low acid, white flesh peach; ‘Ryan Sun’, a medium acid, yellow  
90 flesh peach with balanced sensory attributes; ‘O’Henry’, a medium acid, flavorful, yellow flesh  
91 peach; and ‘Summer Lady’, a medium acid, sweet, yellow flesh peach. The cultivars were  
92 selected based on their previously determined sensory attributes (Crisosto and Crisosto 2005;  
93 Crisosto *et al.* 2006; Crisosto *et al.* 1998). For each cultivar, fruit were harvested at peak size  
94 and California Well-mature for that cultivar from commercial orchards in Fresno Co., CA, then  
95 held at 0 °C (85% RH) for up to 10 d, except for ‘Summer Lady’, which was held at 5 °C (85%

96 RH) prior to ripening to induce onset of chilling injury (Crisosto and Labavitch 2002). The same  
97 fruit from each cultivar was used for both descriptive analysis and the consumer study.

98

### 99 **Instrumental fruit quality measurements**

100 Fruit were ripened in a temperature-controlled room at 20 °C (85% RH) until a  
101 subsample reached a flesh firmness of  $\leq 17.8$  N as described (Crisosto 1999). On the day of the  
102 descriptive analysis session or ‘in-store’ consumer study, a 2-cm diameter piece of skin was  
103 removed from one cheek of each ripened fruit of the cultivar to be tested and the flesh firmness  
104 (penetration force) was measured with a UC firmness tester (Western Industrial Supply, San  
105 Francisco, CA) equipped with an 8 mm tip. If the fruit was ripe ( $\leq 17.8$  N flesh firmness), a  
106 numerical code was written on the tip of the fruit and the flesh firmness recorded. A sample  
107 consisted of one longitudinal slice cut from the stem end to the blossom end on the cheek  
108 opposite that on which flesh firmness was measured (Crisosto and Crisosto 2005). In addition, a  
109 longitudinal wedge was removed from the same area as the flesh firmness measurement, placed  
110 between two layers of cheesecloth, and the juice expressed for subsequent soluble solids  
111 concentration (SSC) and titratable acidity (TA) measurements. The SSC of the juice was  
112 measured with a digital temperature-compensated refractometer (model PR-32 $\alpha$ , Atago Co.,  
113 Tokyo, Japan). TA was measured with an automatic titrator (TitraLab®850, Radiometer  
114 Analytical, Copenhagen, Denmark) and expressed as percent malic acid.

115

### 116 **‘In-Store’ consumer study**

117 One hundred twenty consumers who reported eating fresh nectarines/peaches participated  
118 in the study. The experiment was conducted at a major supermarket in Davis, California. Each

119 consumer evaluated seven samples; the experimental design was a Williams Latin square design  
120 provided by the FIZZ software. One fruit sample per cultivar was evaluated by each consumer  
121 using a written questionnaire. For each nectarine or peach sample, consumers expressed their  
122 overall liking using the 9-point hedonic scale (Peryam and Pilgrim 1957). Consumer acceptance  
123 was calculated as the percentage of respondents who liked the sample, with scores >5.  
124 Consumer dislike of a sample was calculated as the percentage with scores <5. At the  
125 supermarket, the samples were prepared in the produce room out of sight of the testing area as  
126 described (Crisosto and Crisosto 2005).

127

### 128 **Generic descriptive analysis**

129 A generic descriptive analysis (Lawless and Heymann 2010) was used to identify sensory  
130 descriptors for fresh nectarines and peaches. The panel consisted of 11 judges (six women and  
131 five men) with an average age of 32 years. Each judge completed eight training sessions. The  
132 first two sessions covered the development of the language; four sessions were intended to  
133 achieve concept alignment, provide references, eliminate similar terms or ambiguities, and  
134 perfect use of the scale; and the last two sessions evaluated the judges' agreement and  
135 understanding of the attributes. FIZZ software (Biosystèmes) was used to build an automated  
136 session. Sixteen attributes were defined by the panel using standards (Table 2) and evaluated  
137 using a continuous, unstructured 10-cm line scale anchored at the ends by low and high intensity,  
138 except for firmness, which was anchored by soft and hard, and crunchy, which was anchored by  
139 not and very. Samples were evaluated in triplicate with one single fruit used for each judge. The  
140 order of presentation of the samples was randomized using a Latin square design provided by the  
141 FIZZ software.

142

### 143 **Statistical analysis**

144       The majority of the statistical analyses were executed using SAS version 9.1 (SAS  
145 Institute, Cary, NC). To understand the relationships between physicochemical measurements,  
146 sensory attributes, and consumer hedonic ratings, univariate analysis (correlation, analysis of  
147 variance, and Fisher's LSD multiple mean comparisons) and multivariate analysis (canonical  
148 variate analysis (CVA), MANOVA, and preference mapping) were performed. Market clusters  
149 were determined using external preference mapping and cluster analysis. Cluster analysis was  
150 performed with XL-Stat Version 2009.3.02 . The Unscrambler version 9.8 was used to perform  
151 block partial least square regression (L-PLS) analysis.

152

## 153 **RESULTS AND DISCUSSION**

### 154 **Instrumental fruit quality measurements**

155       RSSC varied from 10 to 13.4% and RTA ranged between 0.21 and 0.77% in the ripe fruit  
156 (Table 1). In general, peaches had lower RSSC than nectarines and RTA varied among cultivars.  
157 Among the nectarines, white-fleshed 'August Pearl' and yellow-fleshed 'Fire Sweet' had low  
158 RTAs (~0.30%), while yellow-fleshed 'August Bright' had a high RTA (0.77%). 'Autumn  
159 Snow', a white-fleshed peach, had the lowest RTA (0.21%) and the yellow-fleshed peaches  
160 'O'Henry', 'Summer Lady', and 'Ryan Sun' had medium RTAs (0.50%). Our previous 10 years  
161 of surveys indicated that RSSC is more variable than RTA for a given cultivar over years or  
162 locations. Orchard management and environmental conditions have a strong effect on RSSC but  
163 less on RTA. We observed larger changes in fruit TA than in SSC during ripening on and off the  
164 tree. RTA measurements reported here are similar to those measured in previous surveys. RTA

165 values reported here for ‘August Bright’ nectarine (0.77%) and ‘Summer Lady’, ‘O’Henry,’ and  
166 ‘Ryan Sun’ (~0.50%) peaches were somewhat lower than previously reported for mature fruit  
167 (~0.60 to 0.80%). These differences in RTA are explained by loss of fruit acidity during  
168 ripening.

169

### 170 **‘In-Store’ consumer study**

171 For this consumer population, there were no significant differences in the distribution of  
172 female and male ages ( $p>0.05$ , chi-square test): 58% were female and 42% male. The average  
173 age was 33 years with a standard deviation of 17.5 years. Of the total population, 41% identified  
174 themselves as White-Caucasian, 36% as Asian-Asian American, 18% as Hispanic or Latino, 2%  
175 American Indian or Alaska Native, 1% Black-African American, 12% Mixed or other, and 9%  
176 preferred not to report their ethnicity. The consumption rate of nectarines and peaches for this  
177 consumer population was approximately equally distributed among once a month (23%  
178 nectarines, 24% peaches); two to three times a month (28% nectarines, 23% peaches); once a  
179 week (21% nectarines, 21% peaches), and two to four times a week (12% nectarines, 18%  
180 peaches). These consumption rates were low for when nectarines and peaches are in season; this  
181 supports reports that indicate a static consumption for nectarines and peaches since 1980 (USDA  
182 Economic Research Data, 2009).

183 The 120 consumers differed in their preferences for the seven cultivars (ANOVA  
184  $p<0.05$ ). Pearson’s correlations ( $p \leq 0.05$ ) between overall liking for each cultivar and  
185 instrumental quality measurements of ripe soluble solids concentration (RSSC), ripe titratable  
186 acidity (0.20 to 0.80% RTA), and flesh firmness (6.5 to 20.2 N) were not significant ( $p>0.05$ ),  
187 except for RSSC ( $R^2= 90.2$ ). Overall liking increased significantly from like slightly to like

188 moderately (positive slope) as RSSC increased from 10.0 to 14.0%. Even though the  
189 relationship between RTA and overall liking was not significant, it had a negative slope,  
190 suggesting that cultivars with high acidity were less preferred. The significant negative effect of  
191 high RTA on overall liking has been reported previously on fruit with RTA higher than 0.80 to  
192 1.00% and RSSC lower than 12.0% (Crisosto and Crisosto 2001; Crisosto and Crisosto 2005).  
193 The RTA for the tested cultivars ranged from 0.21 to 0.77%, which may explain the lack of  
194 significant correlation between hedonic scores and RTA measurements. The relationship  
195 between overall liking and flesh firmness had a flat slope and was not significant in any cultivar.  
196 This lack of relationship differs from other fruit commodities such as apples, where texture  
197 change is one of the most important drivers of liking (Harker *et al*, 2008). However, all fruit  
198 tested here was ripened to a firmness penetration force of 6.3 to 17.8 N, considered “ready to  
199 eat” with maximum sensory potential based on our previous work (Crisosto and Crisosto 2005).

200         Since acceptance of these cultivars measured as a degree of liking was not affected by  
201 changes in RTA, and RSSC was the only significant instrumental measurement affecting overall  
202 liking of nectarines and peaches, a detailed statistical analysis between RSSC and degree of  
203 liking was pursued. In general, degree of liking increased as RSSC increased and then reached a  
204 plateau (Table 3). Among the nectarine cultivars, RSSC did not affect degree of liking within  
205 each cultivar and consumer acceptance ranged from 65 to 91% (Table 3). Nectarines with  
206 predominant sensory characteristics of low acidity and/or flavor/aroma had high consumer  
207 acceptance percentages (72 to 91%) and were liked “moderately” to “very much” (6.6 to 7.7).  
208 Nectarines with high acidity were liked less (5.7 to 6.7) and less accepted (65 to 82%). In the  
209 nectarine cultivar with high acidity (~0.80%), acceptance increased and rejection decreased for  
210 fruit with RSSC  $\geq$  12.0%. In the peach with low acidity, RSSC from 10.4 to 14.5% did not

211 significantly affect degree of liking or acceptance. Rejection was around 17% and acceptance  
212 ranged from 67 to 84%. In peaches with predominant sensory characteristics of flavor or  
213 medium acidity, fruit with RSSC < 9.0% had a low degree of liking (~4.5) and acceptance (25 to  
214 36%). For fruit with RSSC  $\geq$  9.0%, degree of liking increased for peaches with high flavor and  
215 for peaches with medium acidity reached a plateau above 9.0% RSSC. In this small population  
216 of consumers that tasted nectarines and peaches with predominant flavor and high RSSC, degree  
217 of liking and acceptance was very high (91%). This data also suggests that nectarines or peaches  
218 with very low acidity may have a low potential consumer acceptance; however, this is affected  
219 by ethnicity (Crisosto and Crisosto 2002). It is important to point out that perception of flavor in  
220 peaches decreased and “off flavor” increased during cold storage as a consequence of chilling  
221 injury (Crisosto and Labavitch 2002; Infante *et al.* 2009). In most cultivars, this flavor loss is  
222 faster when fruit is stored at 5 °C than at 0 °C. In this study, all cultivars were handled rapidly to  
223 avoid any onset of chilling injury except for ‘Summer Lady’, in which onset of loss of flavor or  
224 “off flavor” development may have occurred.

225 A further detailed analysis of nectarine and peach cultivars by hedonic scale categories  
226 for purchase intent, price expectation, second consumption, and RSSC was conducted (Tables 4,  
227 5). In general, consumers were willing to pay more for fruit with a higher hedonic score; this  
228 trend was independent of the nectarine (Table 4) or peach (Table 5) cultivar. The same trends  
229 occurred for purchase intent and willingness to consume the fruit for a second time. Other  
230 researchers have found a correlation between overall liking and the price consumers would be  
231 willing to pay for specialty food such as extra virgin olive oil (Delgado and Guinard 2011;  
232 Stefani *et al.* 2006) and consumers agreed to pay more when they liked cheeses (Napolitano *et*  
233 *al.* 2010). These ‘in-store’ consumer test results agreed with previous studies (Crisosto *et al.*

234 2006), in which peaches and/or nectarines with predominant sensory attributes such as flavor  
235 and/or aroma had a slightly higher consumer acceptance (~10%) than the standard ones. These  
236 results confirmed our previous sensory study and demonstrate that tree fruit degree of liking is  
237 associated with buying habits and even willingness to pay more. These results justify changes in  
238 orchard management to produce more fruit with high RSSC and selection of cultivars with  
239 predominant sensory attributes by plant breeders during cultivar development and growers for  
240 future plantings (Crisosto *et al.* 1997).

241

### 242 **Generic descriptive analysis**

243 Sixteen attributes were defined by the judges to describe the sensory characteristics of the  
244 seven nectarine and peach cultivars (Table 2). These attributes were evaluated through a three-  
245 way ANOVA (judges, cultivars, replications, and all two-way interactions). The ANOVA F-  
246 ratios confirmed that the panel performance was satisfactory (data not shown). The replication  
247 effects were not significant ( $p>0.05$ ) for the majority of attributes evaluated, except for firmness,  
248 crunchy, juicy, and melting. Given the complexity of fresh nectarines and peaches, this  
249 difference may be due more to variation in the fruit than to variation among judges. This  
250 explanation was confirmed because the replication per cultivar interaction was significant,  
251 indicating that there was some variation in the fruit that is reflected in the replication effect.  
252 Chilling injury symptoms such as mealy-woolly texture develop in specific areas in the fruit.  
253 Despite ‘Summer Lady’ peaches having the highest mealy texture score, ‘Summer Lady’ also  
254 had the highest overall, grassy/green fruit, and pit aromas; bitter taste; and melting and fibrous  
255 textures. This cultivar also had moderate sweetness and sourness, and the least floral aroma,  
256 firmness, and crunchy attributes among the cultivars. ‘August Pearl’ nectarine had the highest

257 floral aroma (Table 6). There were no significant differences in the sweetness of ‘Autumn  
258 Snow’ peach and ‘August Pearl’ and ‘Fire Sweet’ nectarines, which had the highest sweetness  
259 scores. The sourest cultivar was ‘August Bright’ nectarine, followed by ‘Ryan Sun’ and  
260 ‘O’Henry’ peaches; the descriptive panel was able to detect some minimal bitterness in these  
261 cultivars, highest for ‘Ryan Sun’, ‘Summer Lady’, and ‘O’Henry’ peaches. ‘August Bright’ and  
262 ‘Fire Sweet’ nectarines had the firmest flesh. Even though care was taken to follow  
263 recommended postharvest ripening practices for stone fruit, these cultivars still may behave  
264 differently during ripening and exhibit slight differences in texture. Differences in texture  
265 perception have also been observed in blueberries at different ripeness (Saftner *et al.* 2008). The  
266 three nectarine cultivars exhibited no significant differences in crunchy texture, which is  
267 somewhat expected because the nectarine cultivars were also the firmest cultivars. Other authors  
268 have found an association between firmness and crunchiness in processed tomatoes (Lee *et al.*  
269 1999) and kiwifruit (Stec *et al.* 1989). The highest means for mealy, melting, and fibrous  
270 textures were found in ‘Summer Lady’ peach, which makes sense because this cultivar was held  
271 at a different temperature than the rest (5 °C) to induce mealy texture. We believe that the low  
272 floral and high grassy/green fruit and pit aromas, combined with high mealiness and low  
273 firmness and crunchy textures detected in ‘Summer Lady’ were the onset of chilling injury  
274 symptoms detected by the judges.

275 A canonical variate analysis (CVA) was conducted to understand sensory similarities and  
276 differences among the seven cultivars and characterize their significant sensory attributes (Fig.  
277 1). The variance explained corresponds to 64.5% on the x axis and 22.3% on the y axis. There  
278 was a significant simultaneous effect among all attributes and cultivars (MANOVA, Wilks’  
279 Lambda,  $F = 9.06$ ;  $df 72$ ,  $p < 0.05$ ). Sour, sweet, floral aroma, overall aroma, and mealy were

280 the main attributes associated with the first dimension (CV1), while firmness, grassy/green fruit  
281 aroma and bitterness were the attributes related to the second dimension (CV2). Pit aroma,  
282 fibrous, and melting have short vectors, an indication of low discrimination for these attributes.  
283 ‘Autumn Snow’ peach was defined as sweet and fibrous, with some grassy/green fruit aroma,  
284 and not sour or mealy. ‘August Pearl’ and ‘Fire Sweet’ nectarines were sweet, firm, crunchy,  
285 and not sour, with a floral aroma. ‘Ryan Sun’ and ‘O’Henry’ peaches and ‘August Bright’  
286 nectarine were defined mainly by their sour taste; the main difference among them was in  
287 firmness. ‘August Bright’ was firmer, crunchier, and also had some floral aroma lacking in  
288 ‘Ryan Sun’ and ‘O’Henry’. ‘Summer Lady’ did not cluster with any other cultivar and was  
289 characterized as melting and mealy, with the highest overall, grassy/green fruit, and pit aromas.

290 The relationship among sweetness, sourness, and firmness, measured both instrumentally  
291 and by a descriptive analysis panel, was studied using principal component analysis (PCA). A  
292 correlation matrix was used to explain the relationship between the instrumental quality  
293 measurements and the descriptive panel sensory attributes measurements (Fig. 2). The total  
294 variance explained was 46.4% on the x axis and 28.4% on the y axis. The PCA analysis  
295 demonstrated that the sensory attributes sourness, firmness, and sweetness were highly correlated  
296 with the instrumental measurements (RTA, firmness, and RSSC). The length of the vector  
297 indicates the discrimination among samples provided by a particular attribute; for example,  
298 firmness as rated by the descriptive panel and firmness as measured instrumentally have exactly  
299 the same length vector, indicating that both methods provide similar discrimination among  
300 samples. However, sweetness and sourness as rated by the descriptive panel had slightly longer  
301 vectors than those generated by RSSC and RTA; this may be an indication that sensory methods  
302 provide better discrimination and characterization among samples for these attributes.

303

304 External preference mapping was selected to study the relationship between the sensory  
305 properties and the hedonic responses; cluster analysis (Wards Method, Euclidean distance)  
306 revealed three segments among the consumer population. Consumers in cluster 1 (n=52)  
307 preferred ‘O’Henry’ peaches and ‘August Bright’ nectarines. Cluster 2 (n=54) and cluster 3  
308 (n=14) were similar: consumers in both liked ‘August Pearl’ and ‘Fire Sweet’ nectarines and  
309 ‘Autumn Snow’ peaches. However, while consumers in cluster 2 accepted cultivars that were  
310 mainly characterized by sourness, consumers in cluster 3 did not like these sour cultivars at all.  
311 Verification of the differences among clusters was accomplished by one-way ANOVA applied to  
312 overall liking on each cluster per cultivar, with the exception of ‘Fire Sweet’ nectarine that had  
313 no difference in liking among the three clusters. Overall liking was significantly different for  
314 each cluster for the rest of the cultivars, so even though the number of consumers in cluster 3 is  
315 small, it is important to keep this cluster. ‘August Bright’ nectarines were liked equally by  
316 clusters 1 and 2, but were disliked by cluster 3. ‘August Pearl’ nectarines received the lowest  
317 average hedonic score from cluster 1, but fared better with clusters 2 and 3. ‘O’Henry’ peaches  
318 were preferred by consumers in cluster 1 only, while ‘Ryan Sun’ was liked only by consumers in  
319 cluster 2. Consumers in cluster 3 provided the highest hedonic score for ‘Autumn Snow’  
320 peaches, while cluster 1 provided the highest hedonic score for ‘Summer Lady’ peaches (Fig. 3).  
321 The external preference mapping provided the main drivers of liking and which cultivars were  
322 preferred most by consumers. Sweetness was the main driver of liking for clusters 2 and 3,  
323 while cluster 1 preferences were mainly driven by the aroma composition (overall, grassy/green  
324 fruit and pit aromas). The fact that sweetness was a key driver of liking agrees with other fruit  
325 commodities such as apples (Dailliant-Spinnler *et al.* 1996; Thybo *et al.* 2004), strawberries

326 (Lado *et al.* 2010), pineapples (Schulbach *et al.* 2007), tomatoes (Causse *et al.* 2010), and fruit-  
327 based products such as apple juice (Rødbotten *et al.* 2009) or pear fruit leathers (Huang and  
328 Hsieh 2005). The perception of grassy/green fruit and pit aromas that were detected in ‘Summer  
329 Lady’ stored at 5 °C to induce chilling injury could be the first signs of chill injury development;  
330 it has been suggested that specific volatiles can be used to detect onset of mealiness (Crisosto  
331 and Labavitch 2002). Block-Partial least square regression, (L-PLS) was used to study the  
332 relationship between the sensory descriptors given by the panel and the hedonic responses of  
333 consumers. L-PLS demonstrated differences in demographics for the clusters (Fig. 4). There  
334 were some differences in preferences according to gender; male consumers preferred more firm,  
335 crunchy cultivars and females tended to like the nectarine ‘August Bright’ more. Ethnicity had a  
336 strong influence on preferences within each cluster; for example, preferences in cluster 1 were  
337 mainly associated with White-Caucasian ethnicity that assigned more importance to overall  
338 aroma. Consumers in clusters 2 and 3 were mainly Asian-Asian Americans who preferred sweet  
339 nectarines and peaches.

340

## 341 **CONCLUSIONS**

342 There were strong correlations between the instrumental measurements of penetration  
343 firmness, ripe soluble solids concentration (RSSC), and ripe titratable acidity (RTA), and their  
344 respective sensory panel descriptors of firmness-texture, sweet, and sour. The sensory  
345 descriptors explained cultivar differences better than instrumental measurements alone.

346 RSSC (sweetness predictor) was the only instrumental measurement that was highly  
347 associated with overall liking by consumers. The expected price that consumers were willing to

348 pay and purchase intent increased with the overall degree of liking and was not affected by  
349 cultivar.

350 Sweetness perception was the main driver of liking for two consumer clusters; however,  
351 for the third cluster the sensory attributes of grassy/green fruit and pit aromas were the main  
352 drivers of liking.

353 **ACKNOWLEDGEMENTS**

354           The authors would like to acknowledge financial support from USDA NIFA grant #  
355 2009-51181-05783, “Increasing consumption of specialty crops by enhancing their quality and  
356 safety” and the California Tree Fruit Agreement for making this work possible. Thanks to Dr.  
357 Helene Hopfer, Melanie Lai, Catherine Hsu, Lauren Ginn, Paulo Wu Liu, and John Zuanyuan  
358 Chan for their assistance on this work.

359

360

361 **REFERENCES**

- 362
- 363 ARES, G., BARRIOS, S., LAREO, C. and LEMA, P. 2009. Development of a sensory quality  
364 index for strawberries based on correlation between sensory data and consumer  
365 perception. *Postharvest Biology and Technology* 52, 97-102.
- 366
- 367 BETT, K. 2002. Evaluating sensory quality of fresh-cut fruits and vegetables. In *Fresh-Cut*  
368 *Fruits and Vegetables*, (O. Lamikanra, ed.) pp. 427-438, CRC Press, Boca Raton, Florida
- 369
- 370 BRUHN, C.M. 1995. Consumer and retailer satisfaction with the quality and size of california  
371 peaches and nectarines. *Journal of Food Quality* 18, 241-256.
- 372
- 373 BRUHN, C.M., FELDMAN, N., GARLITZ, C., HARWOOD, J., IVANS, E., MARSHALL, M.,  
374 RILEY, A., THURBER, D. and WILLIAMSON, E. 1991. Consumer perceptions of  
375 quality: apricots, cantaloupes, peaches, pears, strawberries, and tomatoes. *Journal of Food*  
376 *Quality* 14, 187-195.
- 377
- 378 CAUSSE, M., FRIGUET, C., COIRET, C., LÉPICIER, M., NAVEZ, B., LEE, M.,  
379 HOLTHUYSEN, N., SINESIO, F., MONETA, E. and GRANDILLO, S. 2010. Consumer  
380 Preferences for Fresh Tomato at the European Scale: A Common Segmentation on Taste  
381 and Firmness. *Journal of Food Science* 75, S531-S541.
- 382
- 383 COLARIC, M., VEBERIC, R., STAMPAR, F. and HUDINA, M. 2005. Evaluation of peach and  
384 nectarine fruit quality and correlations between sensory and chemical attributes. *Journal*  
385 *of the Science of Food and Agriculture* 85, 2611-2616.
- 386
- 387 CRISOSTO, C.H. 1999. Optimum procedures for ripening stone fruit. In: *Management of Fruit,*  
388 *Ripening.* Postharvest Horticulture Series No. 9. University of California, Davis, CA, pp.  
389 28-30.
- 390
- 391 CRISOSTO, C.H., CRISOSTO, G. and BOWERMAN, E. 2003. Understanding consumer  
392 acceptance of peach, nectarine and plum cultivars. *Acta Hort. (ISHS)* 115-119.
- 393
- 394 CRISOSTO, C.H. and CRISOSTO, G.M. 2001. Understanding consumer acceptance of early  
395 harvested 'Hayward' kiwifruit. *Postharvest Biology and Technology* 22, 205-213.
- 396
- 397 CRISOSTO, C.H. and CRISOSTO, G.M. 2002. Understanding American and Chinese consumer  
398 acceptance of 'Redglobe' table grapes. *Postharvest Biology and Technology* 24, 155-162.
- 399
- 400 CRISOSTO, C.H. and CRISOSTO, G.M. 2005. Relationship between ripe soluble solids  
401 concentration (RSSC) and consumer acceptance of high and low acid melting flesh peach  
402 and nectarine (*Prunus persica* (L.) Batsch) cultivars. *Postharvest Biology and Technology*  
403 38, 239-246.
- 404 CRISOSTO, C.H., CRISOSTO, G.M., ECHEVERRIA, G. and PUY, J. 2006. Segregation of  
405 peach and nectarine (*Prunus persica* (L.) Batsch) cultivars according to their organoleptic  
406 characteristics. *Postharvest Biology and Technology* 39, 10-18.

- 407 CRISOSTO, C.H., GARNER, D., CRISOSTO, G.M. and BOWERMAN, E. 2004. Increasing  
408 'Blackamber' plum (*Prunus salicina* Lindell) consumer acceptance. *Postharvest Biology*  
409 *and Technology* 34, 237-244.
- 410  
411 CRISOSTO, C.H., JOHNSON, R.S., DAY, K.R. and DEJONG, T. 1997. Orchard factors  
412 affecting postharvest stone fruit quality. *HortScience*, 820-823.
- 413  
414 CRISOSTO, C.H. and LABAVITCH, J.M. 2002. Developing a quantitative method to evaluate  
415 peach (*Prunus persica*) flesh mealiness. *Postharvest Biology and Technology*, 151-158.
- 416  
417 CRISOSTO, G.M., CRISOSTO, C.H. and WATKINS, M. 1998. Chemical and organoleptic  
418 description of white flesh nectarines and peaches. *Acta Hort. (ISHS)* 465, 497-506.
- 419  
420 DAILLANT-SPINLER, B., MACFIE, H.J.H., BEYTS, P.K. and HEDDERLEY, D. 1996.  
421 Relationships between perceived sensory properties and major preference directions of 12  
422 varieties of apples from the Southern Hemisphere. *Food Quality and Preference* 7, 113-  
423 126.
- 424  
425 DELGADO, C. and GUINARD, J.-X. 2011. How do consumer hedonic ratings for extra virgin  
426 olive oil relate to quality ratings by experts and descriptive analysis ratings? *Food Quality*  
427 *and Preference* 22, 213-225.
- 428  
429 ESTI, M., MESSIA, M.C., SINESIO, F., NICOTRA, A., CONTE, L., LA NOTTE, E. and  
430 PALLESCHI, G. 1997. Quality evaluation of peaches and nectarines by electrochemical  
431 and multivariate analyses: relationships between analytical measurements and sensory  
432 attributes. *Food Chemistry* 60, 659-666.
- 433  
434 GUERRA, M., SANZ, M.A. and CASQUERO, P.A. 2009. Influence of Harvest Dates on  
435 Quality, Storage Capacity and Sensory Attributes of European Plum cv. Green Gage.  
436 *Food Science and Technology International* 15, 527-534.
- 437  
438 GUNNESS, P., KRAVCHUK, O., NOTTINGHAM, S.M., D'ARCY, B.R. and GIDLEY, M.J.  
439 2009. Sensory analysis of individual strawberry fruit and comparison with instrumental  
440 analysis. *Postharvest Biology and Technology* 52, 164-172.
- 441  
442 HARKER, F.R., GUNSON, F.A. and JAEGER, S.R. 2003. The case for fruit quality: an  
443 interpretive review of consumer attitudes, and preferences for apples. *Postharvest*  
444 *Biology and Technology* 28, 333-347.
- 445  
446 HARKER, F.R., KUPFERMAN, E.M., MARIN, A.B., GUNSON, F.A. and TRIGGS, C.M.  
447 2008. Eating quality standards for apples based on consumer preferences. *Postharvest*  
448 *Biology and Technology* 50, 70-78.
- 449 HARKER, F.R., MAINDONALD, J., MURRAY, S.H., GUNSON, F.A., HALLETT, I.C. and  
450 WALKER, S.B. 2002. Sensory interpretation of instrumental measurements 1: texture of  
451 apple fruit. *Postharvest Biology and Technology* 24, 225-239.
- 452

- 453 HOEHN, E., GASSER, F., GUGGENBÜHL, B. and KÜNSCH, U. 2003. Efficacy of  
454 instrumental measurements for determination of minimum requirements of firmness,  
455 soluble solids, and acidity of several apple varieties in comparison to consumer  
456 expectations. *Postharvest Biology and Technology* 27, 27-37.  
457
- 458 HUANG, X. and HSIEH, F.-H. 2005. Physical Properties, Sensory Attributes, and Consumer  
459 Preference of Pear Fruit Leather. *Journal of Food Science* 70, E177-E186.  
460
- 461 INFANTE, R., MENESES, C. and CRISOSTO, C.H. 2009. Preconditioning treatment maintains  
462 taste characteristic perception of ripe 'September Sun' peach following cold storage.  
463 *International Journal of Food Science & Technology* 44, 1011-1016.  
464
- 465 KÜHN, B.F. and THYBO, A.K. 2001. The influence of sensory and physiochemical quality on  
466 Danish children's preferences for apples. *Food Quality and Preference* 12, 543-550.  
467
- 468 LADO, J., VICENTE, E., MANZZIONI, A. and ARES, G. 2010. Application of a check-all-that-  
469 apply question for the evaluation of strawberry cultivars from a breeding program.  
470 *Journal of the Science of Food and Agriculture* 90, 2268-2275.  
471
- 472 LAWLESS, H.T. and HEYMANN, H. 2010. *Sensory Evaluation of Food*, Second Ed., Springer,  
473 NY.  
474
- 475 LEE, S.Y., LUNA-GUZMÁN, I., CHANG, S., BARRETT, D.M. and GUINARD, J.X. 1999.  
476 Relating descriptive analysis and instrumental texture data of processed diced tomatoes.  
477 *Food Quality and Preference* 10, 447-455.  
478
- 479 LENGARD, V. and KERMIT, M. 2006. 3-Way and 3-block PLS regressions in consumer  
480 preference analysis. *Food Quality and Preference* 17, 234-242  
481
- 482 MALUNDO, T.M.M., SHEWFELT, R.L., WARE, G.O. and BALDWIN, A.E.A. 2001. An  
483 alternative method for relating consumer and descriptive data used to identify critical  
484 flavor properties of mango (*Mangifera indica l.*). *Journal of Sensory Studies* 16, 199-214.  
485
- 486 NAPOLITANO, F., BRAGHIERI, A., PIASENTIER, E., FAVOTTO, S., NASPETTI, S. and  
487 ZANOLI, R. 2010. Cheese liking and consumer willingness to pay as affected by  
488 information about organic production. *Journal of Dairy Research* 77, 280-286.  
489
- 490 OBENLAND, D., COLLIN, S., MACKAY, B., SIEVERT, J., FJELD, K. and ARPAIA, M.L.  
491 2009. Determinants of flavor acceptability during the maturation of navel oranges.  
492 *Postharvest Biology and Technology* 52, 156-163.  
493
- 494 ORAGUZIE, N., ALSPACH, P., VOLZ, R., WHITWORTH, C., RANATUNGA, C.,  
495 WESKETT, R. and HARKER, R. 2009. Postharvest assessment of fruit quality  
496 parameters in apple using both instruments and an expert panel. *Postharvest Biology and*  
497 *Technology* 52, 279-287.  
498

- 499 PERYAM, D.R. and PILGRIM, F.J. 1957. Hedonic scale method of measuring food preferences.  
500 Food Technology *11*, 9-14.  
501
- 502 PREDIERI, S. and GATTI, E. 2009. Effects of cold storage and shelf-life on sensory quality and  
503 consumer acceptance of 'Abate Fetel' pears. Postharvest Biology and Technology *51*,  
504 342-348.  
505
- 506 RØDBOTTEN, M., MARTINSEN, B.K., BORGE, G.I., MORTVEDT, H.S., KNUTSEN, S.H.,  
507 LEA, P. and NÆS, T. 2009. A cross-cultural study of preference for apple juice with  
508 different sugar and acid contents. Food Quality and Preference *20*, 277-284.  
509
- 510 ROSSITER, K.L., YOUNG, H., WALKER, S.B., MILLER, M. and DAWSON, D.M. 2000. The  
511 effects of sugars and acids on consumer acceptability of kiwifruit. Journal of Sensory  
512 Studies *15*, 241-250.  
513
- 514 SAFTNER, R., POLASHOCK, J., EHLENFELDT, M. and VINYARD, B. 2008. Instrumental  
515 and sensory quality characteristics of blueberry fruit from twelve cultivars. Postharvest  
516 Biology and Technology *49*, 19-26.  
517
- 518 SCHULBACH, K.F., PORTIER, K.M. and SIMS, C.A. 2007. Evaluation of overall acceptability  
519 of fresh pineapple using the regression tree approach. Journal of Food Quality *30*, 993-  
520 1008.  
521
- 522 SINESIO, F., CAMMARERI, M., MONETA, E., NAVEZ, B., PEPARAIIO, M., CAUSSE, M.  
523 and GRANDILLO, S. 2010. Sensory Quality of Fresh French and Dutch Market  
524 Tomatoes: A Preference Mapping Study with Italian Consumers. Journal of Food Science  
525 *75*, S55-S67.  
526
- 527 STEC, M.G.H., HODGSON, J.A., MACRAE, E.A. and TRIGGS, C.M. 1989. Role of fruit  
528 firmness in the sensory evaluation of kiwifruit (*Actinidia deliciosa* cv Hayward). Journal  
529 of the Science of Food and Agriculture *47*, 417-433.  
530
- 531 STEFANI, G., ROMANO, D. and CAVICCHI, A. 2006. Consumer expectations, liking and  
532 willingness to pay for specialty foods: Do sensory characteristics tell the whole story?  
533 Food Quality and Preference *17*, 53-62.  
534
- 535 THYBO, A.K., KÜHN, B.F. and MARTENS, H. 2004. Explaining Danish children's preferences  
536 for apples using instrumental, sensory and demographic/behavioural data. Food Quality  
537 and Preference *15*, 53-63  
538
- 539 YENKET, R., CHAMBERS IV, E. and ADHIKARI, K. 2011. A comparison of seven preference  
540 mapping techniques using four software programs. Journal of Sensory Studies *26*, 135-  
541 150.  
542

543 TABLE 1. CODES, MEANS (X) AND STANDARD DEVIATIONS (S.D.) OF FLESH  
 544 FIRMNESS, RIPE SOLUBLE SOLIDS CONCENTRATION (RSSC) AND RIPE  
 545 TITRATABLE ACIDITY (RTA) FOR NECTARINE AND PEACH CULTIVARS  
 546 EVALUATED IN THE DESCRIPTIVE ANALYSIS AND CONSUMER STUDY.  
 547

Fruit	Cultivar	Type*	Cultivar Code	Flesh Color	Firmness		RSSC		RTA	
					(N)	(%)	(%)	(%)		
					X	S.D.	X	S.D.	X	S.D.
Nectarine	Fire	N-F	FS	Yellow	14.6	0.8	13.4	1.2	0.31	0.04
	Sweet									
	August	N-HA	AB	Yellow	12.8	0.7	11.8	1.2	0.77	0.08
	Bright									
	August	N-LA	AP	White	18.0	1.0	13.1	1.2	0.24	0.04
	Pearl									
	O'Henry	P-F	OH	Yellow	15.0	0.8	10.0	1.1	0.49	0.07
Peach	Ryan	P-MA	RS	Yellow	17.7	1.2	10.8	1.4	0.48	0.07
	Sun									
	Autumn	P-LA	AS	White	17.5	1.5	12.6	1.2	0.21	0.04
	Snow									
	Summer	P-SM	SL	Yellow	6.3	0.5	12.0	1.1	0.51	0.07
	Lady									

548

549 \* N-F = nectarine flavorful, N-HA = nectarine high acidity, N-LA = nectarine low acidity, P-F =  
 550 peach flavorful, P-MA = peach medium acidity, P-LA = peach low acidity, P-SM = peach  
 551 slightly mealy.  
 552

553 TABLE 2. SENSORY ATTRIBUTE DEFINITIONS AND STANDARDS USED FOR  
 554 TRAINING THE DESCRIPTIVE ANALYSIS PANEL TO EVALUATE NECTARINE AND  
 555 PEACH CULTIVARS.  
 556

Attributes	Description	Standard
Overall aroma	Intensity of aroma (whole aroma)	Verbal description
Floral	Smell of flowers	Nectarine cultivars
Almond	Smell of almonds	1% Artificial flavor (McCormick) over 50 g of peach/nectarine paste*
Grassy/green fruit	Smell of grass associated with unripe fruit	0.01 grams over 50 g of peach/nectarine paste*
Pit	Woody aroma associated with fruit with traces of pit	10 pits were removed from the fruit and used as a standard
Overall flavor	Intensity of flavor (whole flavor)	Verbal description
Sweet	Sweet taste, example sucrose solution	Sucrose solutions 0- 50 g/L in spring water Fruit with different soluble solids concentrations ranging from 8-16%
Sour	Sour taste, example citric acid solution	Citric acid solutions 0 – 1.5 g/L in spring water Fruit with different titratable acidities ranging from 0.2% to 0.8%
Bitter	Bitter taste, example like coffee	Caffeine solutions 0.7 g/L – 2.5 g/L
Firmness	Flesh only: measured at the first bite ranging from soft to hard	Fruit with different firmness, ranging from 4.4 to 44.4 N
Crunchy	Flesh only: making a crunching sound when chewed or pressed	Verbal description
Juicy	Flesh only: amount of liquid	Verbal description
Mealy	Gritty, sandy texture, dry not juicy	Verbal description Fruit with different degrees of mealiness
Melting	How easy to fracture into mouth (high melting examples ice cream, chocolate)	Verbal description and use of ice cream and chocolate to explain concept
Smooth	Texture of the fruit related to having a continuous even surface	Verbal description
Fibrous	After first two bites, amount of fibers in the sample	Verbal description

557

558 \* A paste (50/50) peach/nectarine was prepared as base.

559

560 TABLE 3. DEGREE OF LIKING AND CONSUMER ACCEPTANCE ACCORDING TO RIPE  
 561 SOLUBLE SOLIDS CONCENTRATION (RSSC) CLASSES FOR DIFFERENT NECTARINE  
 562 AND PEACH CULTIVARS AND THEIR CORRESPONDING INSTRUMENTAL  
 563 MEASUREMENTS.  
 564

Cultivar Type*	RSSC class (%)	n	Liking (1-9) <sup>***</sup>	Acceptance (%)	Neutral (%)	Rejection (%)	RSSC (%)	RTA (%)
N-LA	11-11.9	19	7.2 a <sup>***</sup>	84	5	11	11.6	0.26
	12-12.9	41	7.0 a	83	7	10	12.5	0.24
	13-13.9	39	7.1 a	82	15	3	13.4	0.24
	14->15	19	7.3 a	89	11	0	14.6	0.23
	RSSC class	n	Liking	Acceptance	Neutral	Rejection	RSSC	RTA
N-F	9.8-11.9	12	6.6 a	75	17	8	11.3	0.31
	12-12.9	25	6.6 a	72	16	12	12.5	0.31
	13-13.9	48	6.7 a	83	6	10	13.5	0.31
	14-14.9	24	7.0 a	88	4	8	14.2	0.31
	15->16	11	7.7 a	91	0	9	15.7	0.29
	RSSC class	n	Liking	Acceptance	Neutral	Rejection	RSSC	RTA
N-HA	10-10.9	31	5.7 a	65	3	32	10.4	0.75
	11-11.9	43	6.0 a	74	0	26	11.4	0.78
	12-12.9	28	6.7 a	82	0	18	12.3	0.80
	13->16	18	6.5 a	78	6	17	13.6	0.80
	RSSC class	n	Liking	Acceptance	Neutral	Rejection	RSSC	RTA
P-LA	<10-10.9	9	5.6 a	67	11	22	10.4	0.20
	11-11.9	34	6.4 a	71	12	18	11.6	0.20
	12-12.9	34	6.5 a	68	12	21	12.5	0.20
	13-13.9	31	6.7 a	84	6	10	13.4	0.20
	14-<16	12	6.7 a	75	8	17	14.5	0.20
	RSSC class	n	Liking	Acceptance	Neutral	Rejection	RSSC	RTA
P-F	<9	22	4.8 a	36	23	41	8.5	0.47
	9-9.9	41	5.4 ab	59	7	34	9.3	0.47
	10-10.9	33	6.2 bc	76	6	18	10.4	0.49
	11-11.9	13	5.7 abc	62	8	31	11.4	0.54
	12-<13	11	7.2 c	91	0	9	12.4	0.54
	RSSC class	n	Liking	Acceptance	Neutral	Rejection	RSSC	RTA
P-MA	<9	12	4.2 a	25	8	67	8.4	0.483
	9-9.9	25	6.0 bc	72	4	24	9.5	0.455
	10-10.9	34	5.4 ab	62	12	26	10.4	0.452
	11-11.9	28	6.3 bc	71	4	25	11.3	0.487
	12-<15	21	6.6 c	71	19	10	12.7	0.472
	RSSC class	n	Liking	Acceptance	Neutral	Rejection	RSSC	RTA
P-SM	9-9.9	5	5.6a	60	20	20	9.5	0.5

10-10.9	14	6.6a	71	7	21	10.5	0.5
11-11.9	30	6.7a	80	3	17	11.5	0.5
12-<15	71	6.5a	78	4	18	12.8	0.5

565

566 \* Cultivar type: N-F = nectarine flavorful, N-HA = nectarine high acidity, N-LA = nectarine low acidity,  
 567 P-F = peach flavorful, P-MA = peach medium acidity, P-LA = peach low acidity, P-SM = peach slightly  
 568 mealy.

569

570 \*\* Degree of liking: 1 = dislike extremely, 2 = dislike very much, 3 = dislike moderately, 4 = dislike  
 571 slightly, 5 = neither like nor dislike, 6 = like slightly, 7 = like moderately, 8 = like very much, 9 = like  
 572 extremely.

573

574 \*\*\* Same letters within the same column indicate no significant difference between means  $p < 0.05$ .

575

576 TABLE 4. PURCHASE INTENT, PRICE EXPECTATION, SECOND CONSUMPTION, AND  
 577 RIPE SOLUBLE SOLIDS CONCENTRATION (RSSC) BY DEGREE OF LIKING (1-9) FOR  
 578 NECTARINE CULTIVARS BY NORTHERN CALIFORNIA CONSUMERS.  
 579

Cultivar type*	Degree of liking** (score)	Purchase Intent***	Price**** (\$)	Second Consumption*****	RSSC (%)
N-F	1.0	1.0 a	0.00 a	1.0 a	12.7
	2.0	-	-	-	-
	3.0	2.0 a	0.40 a	2.0 a	13.1
	4.0	2.2 a	0.70 a	2.3 a	13.8
	5.0	3.1 b	0.60 a	3.0 b	12.6
	6.0	3.4 b	1.30 b	3.1 b	13.4
	7.0	3.9 c	1.30 b	3.9 c	13.3
	8.0	4.5 d	1.60 b	4.5 d	13.5
	9.0	4.8 d	1.60 b	4.9 d	14.3
	<i>P</i> -value	<0.0001	<0.0001	<0.0001	<0.0001
N-HA	1.0	1.0 a	0.00 a	1.0 a	11.5
	2.0	1.1 a	0.30 a	1.3 ab	11.8
	3.0	1.7 b	0.40 a	1.9 bc	11.6
	4.0	2.4 c	0.40 a	2.2 cd	10.8
	5.0	4.0 de	0.70 ab	3.0 de	11.8
	6.0	3.6 d	1.30 b	3.5 e	11.5
	7.0	3.8 d	1.40 b	3.9 e	11.8
	8.0	4.3 e	1.50 b	4.3 f	11.8
	9.0	4.9 f	2.50 c	4.8 g	12.5
	<i>P</i> -value	<0.0001	<0.0001	<0.0001	<0.0001
N-LA	1.0	-	-	-	-
	2.0	-	-	-	-
	3.0	2.3 ab	0.80 ab	1.9 a	11.8
	4.0	1.8 a	0.40 a	2.2 ab	12.6
	5.0	2.8 b	0.70 ab	3.0 bc	13.3
	6.0	3.1 b	1.10 b	3.5 c	13.1
	7.0	3.8 c	1.10 b	3.9 d	13.0
	8.0	4.5 d	1.70 c	4.3 e	13.1
	9.0	5.0 e	1.90 c	4.8 f	13.1
	<i>P</i> -value	<0.0001	<0.0001	<0.0001	<0.0001

580

581 \* Cultivar type: N-F = nectarine flavorful, N-HA = nectarine high acidity, N-LA = nectarine low acidity,  
 582 P-F = peach flavorful, P-MA = peach medium acidity, P-LA = peach low acidity, P-SM = peach slightly  
 583 mealy.

584  
 585 \*\* Degree of liking score: 1 = dislike extremely, 2 = dislike very much, 3 = dislike moderately, 4 = dislike  
 586 slightly, 5 = neither like nor dislike, 6 = like slightly, 7 = like moderately, 8 = like very much, 9 = like  
 587 extremely.

588  
 589 \*\*\* Purchase intent: 1 = definitely would not buy, 2 = probably would not buy, 3 = neither would not buy,  
 590 nor would buy, 4 = probably would buy, 5 = definitely would buy.

591  
592  
593  
594  
595  
596  
597  
598  
599

\*\*\* Price willing to pay per pound of the sample tasted at retail.

\*\*\*\* Second consumption: 1 = certainly will not consume this nectarine again, 2 = probably will not consume this nectarine again, 3 = not sure or undecided, 4 = probably will consume this nectarine again, 5 = certainly will consume this nectarine again.

\*\*\*\*\* Same letters within the same column indicate no significant difference between means.

600

601 TABLE 5. PURCHASE INTENT, PRICE EXPECTATION, SECOND CONSUMPTION, AND  
 602 RIPE SOLUBLE SOLIDS CONCENTRATION (RSSC) BY DEGREE OF LIKING (1-9) FOR  
 603 PEACH CULTIVARS BY NORTHERN CALIFORNIA CONSUMERS.  
 604

Cultivar Type*	Degree of liking** (score)	Purchase Intent***	Price**** (\$)	Second Consumption*****	RSSC (%)
P-F	1.0	1.0 a	0.00 a	1.0 a	9.9 ab
	2.0	1.4 a	0.10 a	1.4 a	9.9 ab
	3.0	2.0 b	0.40 a	1.7 ab	9.6 ab
	4.0	2.2 b	0.50 a	2.1 b	9.5 ab
	5.0	2.9 c	0.70 ab	3.0 c	9.3 a
	6.0	3.1 c	1.00 bc	3.0 c	9.8 ab
	7.0	3.9 d	1.40 cd	3.9 d	10.1 bc
	8.0	4.3 e	1.60 d	4.2 de	10.6 cd
	9.0	5.0 f	1.5cd	4.8 e	11.8 d
		<i>P</i> -value	<0.0001	<0.0001	<0.0001
P-MA	1.0	1.0 a	0.10 a	1.0 a	10.5 ab
	2.0	1.1 a	0.40 ab	1.6 a	9.4 a
	3.0	1.8 b	0.30 a	1.5 a	10.3 ab
	4.0	2.3 b	0.80 abc	2.4 b	10.3 ab
	5.0	2.8 c	0.80 abc	2.6 b	11.3 b
	6.0	3.3 d	0.90 bc	3.4 c	10.3 ab
	7.0	3.6 d	1.20 c	3.7 c	10.8 b
	8.0	4.6 e	1.70 d	4.4 d	11.2 b
	9.0	5.0 e	2.00 d	5.0 d	10.6 ab
		<i>P</i> -value	<0.0001	<0.0001	<0.0001
P-LA	1.0	1.0 a	-	3.0 cd	13.1
	2.0	2.0 ab	0.30 ac	1.5 ab	12.7
	3.0	1.2 a	0.20 ab	1.3 a	11.8
	4.0	1.9 a	0.60 bc	2.2 bc	12.5
	5.0	2.7 bc	0.80 bc	2.6 c	12.4
	6.0	3.0 c	0.90 c	3.2 d	12.3
	7.0	3.6 d	1.50 d	3.9 d	12.6
	8.0	4.5 e	1.30 d	4.7 e	12.5
	9.0	4.8 e	2.0 e	4.9 e	12.9
		<i>P</i> -value	<0.0001	<0.0001	<0.0001
P-SM	1.0	1.0 a	0.0 a	1.0 a	10.9
	2.0	1.0 a	0.1 ab	1.0 a	12.2
	3.0	1.8 b	0.1 ab	1.8 b	12.1
	4.0	2.4 c	0.7 bcd	2.7 c	11.7
	5.0	2.5 c	0.8 cde	3.0 cd	11.4
	6.0	3.2 d	1.0 de	3.5 d	11.9
	7.0	3.9 e	1.4 e	3.9 e	12.2
	8.0	4.4 f	1.7 f	4.5 f	12.2
	9.0	4.8 g	2.0 f	4.7 f	11.8
		<i>P</i> -value	<0.0001	<0.0001	<0.0001

606 \* Cultivar type: N-F = nectarine flavorful, N-HA = nectarine high acidity, N-LA = nectarine low acidity,  
607 P-F = peach flavorful, P-MA = peach medium acidity, P-LA = peach low acidity, P-SM = peach slightly  
608 mealy.

609  
610 \*\* Degree of liking score: 1 = dislike extremely, 2 = dislike very much, 3 = dislike moderately, 4 = dislike  
611 slightly, 5 = neither like nor dislike, 6 = like slightly, 7 = like moderately, 8 = like very much, 9 = like  
612 extremely.

613  
614 \*\*\* Purchase intent: 1 = definitely would not buy, 2 = probably would not buy, 3 = neither would not buy,  
615 nor would buy, 4 = probably would buy, 5 = definitely would buy.

616  
617 \*\*\*\* Price willing to pay per pound of the sample tasted at retail.

618  
619 \*\*\*\*\* Second consumption: 1 = certainly will not consume this nectarine again, 2 = probably will not  
620 consume this nectarine again, 3 = not sure or undecided, 4 = probably will consume this nectarine again, 5  
621 = certainly will consume this nectarine again.

622  
623 \*\*\*\*\* Same letters within the same column indicate no significant difference between means.

624

625

626 TABLE 6. OVERALL MEANS FOR THE SIGNIFICANT SENSORY ATTRIBUTES OF  
 627 AROMA, TASTE AND TEXTURE FOR SEVEN NECTARINE AND PEACH CULTIVARS.  
 628

Aroma Attributes											
Overall			Floral			Grassy/green fruit			Pit		
Cultivar type *	Mean		Cultivar type	Mean		Cultivar Type	Mean		Cultivar type	Mean	
P-SM	5.4	a**	N-LA	4.2	a	P-SM	2.5	a	P-SM	2.9	a
P-F	4.6	a	N-HA	3.1	b	P-F	2.1	a b	P-F	1.9	b
N-LA	4.5	b	P-MA	2.9	b	N-F	1.7	c b	N-HA	1.7	b
N-HA	4.4	c	P-LA	2.7	b	N-HA	1.4	c b d	N-F	1.5	b c d
P-MA	4.4	c	N-F	2.5	b	P-MA	1.3	c d	P-MA	1.3	c d
N-F	3.7	c	P-F	2.4	b	N-LA	1.0	c d	P-LA	0.9	c d
P-LA	3.3	d	P-SM	2.4	B	P-LA	0.9	d	N-LA	0.8	
LSD	0.78		LSD	0.83		LSD	0.68		LSD	0.61	

629  
630

Taste Attributes								
Sweet		Sour		Bitter				
Cultivar type	Mean	Cultivar type	Mean	Cultivar Type	Mean			
P-LA	5.3	a	N-HA	6.0	a	P-MA	1.3	a
N-LA	5.1	a	P-MA	4.6	b	P-SM	1.2	a b
N-F	4.9	a	P-F	4.6	b	P-F	1.0	a b c
P-SM	3.6	b	P-SM	3.7	c	P-LA	0.8	b c
P-MA	3.0	b c	P-LA	1.3	d	N-F	0.7	b c
N-HA	2.7	c	N-F	1.3	d	N-LA	0.6	c
P-F	2.4	c	N-LA	0.8	d	N-HA	0.5	c
LSD	0.69		LSD	0.68		LSD	0.51	

631  
632

Texture Attributes														
Firmness			Crunchy			Mealy			Melting			Fibrous		
Cultivar type	Mean		Cultivar Type	Mean		Cultivar Type	Mean		Cultivar type	Mean		Cultivar type	Mean	
N-HA	4.6	a	N-HA	3.6	a	P-SM	2.2	a	P-SM	6.8	a	P-SM	4.8	a
N-F	4.1	a b	N-LA	3.3	a	N-LA	1.3	b	P-LA	4.7	b	N-LA	4.6	a b
N-LA	3.8	c b	N-F	3.0	a	N-HA	1.1	b c	P-MA	4.4	b	P-F	4.4	a b
P-F	3.6	c b d	P-F	2.3	b	N-F	1.1	b c	N-LA	3.8	b c	P-LA	4.4	a b
P-LA	3.2	c	P-LA	1.9	b	P-LA	0.8	b c	N-F	3.8	b c	N-HA	4.3	a b c
P-MA	3.0	d	P-MA	1.9	b	P-MA	0.8	b c	P-F	3.8	b c	P-MA	3.8	b c
P-SM	1.2	e	P-SM	1.0	c	P-F	0.5		N-HA	3.4	c	N-F	3.4	c
LSD	0.63		LSD	0.68		LSD	0.68		LSD	0.97		LSD	0.95	

633

634 \* Cultivar type: N-F = nectarine flavorful, N-HA = nectarine high acidity, N-LA = nectarine low acidity,  
635 P-F = peach flavorful, P-MA = peach medium acidity, P-LA = peach low acidity, P-SM = peach slightly  
636 mealy.

637

638 \*\* Same letters within the same column indicate no significant difference between means  $p < 0.05$ .

639

640 Figure captions:

641

642 FIG. 1. CANONICAL VARIATE ANALYSIS (CVA) FOR SEVEN FRESH NECTARINE  
643 AND PEACH CULTIVARS.

644

645 Each sphere represents a cultivar\* (refer to Table 1 for cultivar descriptions). There is no  
646 significant difference between cultivars ( $p < 0.05$ ) when two spheres overlap. Attributes are  
647 represented by vectors.

648

649 \* Cultivar type (cultivar): N-F = nectarine flavorful ('Fire Sweet'), N-HA = nectarine high acidity  
650 ('August Bright'), N-LA = nectarine low acidity ('August Pearl'), P-F = peach flavorful  
651 ('O'Henry'), P-MA = peach medium acidity ('Ryan Sun'), P-LA = peach low acidity ('Autumn  
652 Snow'), P-SM = peach slightly mealy ('Summer Lady').

653

654

655 FIG. 2. PRINCIPAL COMPONENT ANALYSIS BY CORRELATION OF INSTRUMENTAL  
656 VS. DESCRIPTIVE VARIABLES.

657

658 Cultivars are represented by ellipses and attributes by vectors. Please refer to Table 1 for a full  
659 description of the samples. Cultivar type (cultivar): N-F = nectarine flavorful ('Fire Sweet'), N-  
660 HA = nectarine high acidity ('August Bright'), N-LA = nectarine low acidity ('August Pearl'), P-  
661 F = peach flavorful ('O'Henry'), P-MA = peach medium acidity ('Ryan Sun'), P-LA = peach  
662 low acidity ('Autumn Snow'), P-SM = peach slightly mealy ('Summer Lady').

663

664

665 FIG. 3. EXTERNAL PREFERENCE MAPPING INCLUDING CONSUMER SEGMENTS

666

667 Cultivars are represented by solid circles. Please refer to Table 1 for a full description of the  
668 samples. Cultivar type (cultivar): N-F = nectarine flavorful ('Fire Sweet'), N-HA = nectarine  
669 high acidity ('August Bright'), N-LA = nectarine low acidity ('August Pearl'), P-F = peach  
670 flavorful ('O'Henry'), P-MA = peach medium acidity ('Ryan Sun'), P-LA = peach low acidity  
671 ('Autumn Snow'), P-SM = peach slightly mealy ('Summer Lady').

672

673 Solid square indicates consumers belong to cluster 1 ( $n=52$ ). Consumers in cluster 2 ( $n=54$ ) are  
674 indicated by a triangle, and cluster 3 ( $n=14$ ) by a cross.

675

676 FIG. 4. BLOCK-PARTIAL LEAST SQUARE REGRESSION (L-PLS).

677

678 Descriptive analysis data: sensory attributes per cultivar [X] vs. overall liking by 120 consumers  
679 indicated by solid squares [Y] and consumers' demographics [Z]. Variables in the outer circle  
680 are more important than variables in the inner circle. Cultivar type (cultivar): N-F = nectarine  
681 flavorful ('Fire Sweet'), N-HA = nectarine high acidity ('August Bright'), N-LA = nectarine low  
682 acidity ('August Pearl'), P-F = peach flavorful ('O'Henry'), P-MA = peach medium acidity  
683 ('Ryan Sun'), P-LA = peach low acidity ('Autumn Snow'), P-SM = peach slightly mealy  
684 ('Summer Lady').

685