

Sugarbeet

Lee Panella,* Stephen R. Kaffka, Robert T. Lewellen,
J. Mitchell McGrath, Mike S. Metzger, and Carl A. Strausbaugh

ABSTRACT

Sugarbeet (*Beta vulgaris* L. subsp. *vulgaris*) is an industrial crop grown commercially as a hybrid, with sucrose refined from the root as the plant constituent of interest. Yield of refined white sugar produced per unit area is a complex trait. Sugar yield is calculated by the root weight times the proportion of root that is sucrose, minus the amount of sucrose lost during processing, often expressed as percent loss to molasses or juice purity. Yields have risen in the last 100 yr from about 10% sucrose and 22.4 Mg ha⁻¹ (10 t ac⁻¹) to 18% sucrose and more than 65 Mg ha⁻¹ (29.3 t ac⁻¹). There also has been a shift in acreage from the western United States (especially California and Colorado) to the northern Midwest (Red River Valley). Although not all of this improvement is attributable to genetic improvement in the crop, a strong collaborative effort between commercial breeders and public breeders (USDA-ARS) has increased yield potential, while improving resistance to many diseases, leading to increases in yield. This continuing collaboration between private and public breeders will be necessary to meet the challenges of climate change and the utilization of beet as a sustainable biofuel and chemical feedstock.

Abbreviations: BCT, beet curly top; BNYVV, Beet necrotic yellow vein virus; CMS, cytoplasmic male sterility; GHG, greenhouse gases; QTL, quantitative trait loci; RRV, Red River Valley.

L. Panella, USDA-ARS, NPA, Crop Research Lab., 1701 Centre Dr., Fort Collins, CO 80526.

*Corresponding author (Lee.Panella@ars.usda.gov).

S.R. Kaffka, Dep. of Plant Sciences, Univ. of California, Davis, CA 95616-8780 (srkaffka@ucdavis.edu).

R.T. Lewellen, USDA-ARS, Crop Improvement and Protection Research Unit, 1636 E. Alisal St., Salinas, CA 93905 (Robert.Lewellen@ars.usda.gov).

J.M. McGrath, USDA-ARS, Sugarbeet and Bean Research Unit, Dep. of Plant, Soil, and Microbial Sciences, 494 Plant & Soil Sci. Bldg., 1066 Bogue St., Michigan State Univ., East Lansing, MI 48824-1325 (mitchmcg@msu.edu).

M. Metzger, Minn-Dak Farmer's Cooperative, 7525 Red River Rd., Wahpeton, ND 58075 (mmetzger@mdf.coop).

C.A. Strausbaugh, USDA-ARS, Northwest Irrigation and Soils Research Lab., 3793 North 3600 East, Kimberly, ID 83341-5076 (Carl.Strausbaugh@ars.usda.gov).

doi:10.2135/cssaspecpub33.c13

Yield Gains in Major U.S. Field Crops. CSSA Special Publication 33.

Stephen Smith, Brian Diers, James Specht, and Brett Carver, editors.

© 2014. ASA, CSSA, and SSSA, 5585 Guilford Rd., Madison, WI 53711-5801, USA.

Sugarbeet is an industrial crop. Neither the seed nor the foliage is the plant constituent of commercial interest—not even the root itself, but rather the sucrose refined from the root (an enlarged hypocotyl). Yield of refined white sugar produced per unit area is a complex trait. Sugar yield is calculated by the root weight times the proportion of root that is sucrose (usually expressed as the percentage of fresh weight), minus the amount of sucrose lost during processing (often expressed as the percentage of loss to molasses or juice purity). Loss to molasses arises from cations such as Na⁺, K⁺, and amino nitrogen (R-NH₃⁺) balancing the charge of the anion sucrose at physiological pH. The cations that interfere with the extraction of sucrose are related in a large extent to the plant nutrients in the soil in which the sugarbeet is grown; however, the small amino-nitrogen compounds (e.g., betaine) can be lowered by proper nitrogen fertilization (Follett et al., 1970). All of these components are variable in a population and can be improved through selection and breeding (Biancardi et al., 2010).

Taxonomically, sugarbeet is ordered as Dicotyledoneae, Caryophyllidae (Centrospmae), Amaranthaceae (formerly Chenopodiaceae), *Beta vulgaris* L. subsp. *vulgaris* (McGrath et al., 2011). Originally, it is likely that beet was domesticated as a pot herb, and leaves first harvested from the wild progenitor, sea beet [*B. vulgaris* L. subsp. *maritima* (L.) Arcang] for food (Biancardi et al., 2012; Ford-Lloyd et al., 1975; Lange et al., 1999). By the Middle Ages there is reference to the root being used, both as a vegetable and medicinal herb (Biancardi et al., 2012; Goldman and Navazio, 2008). William Morgan provided a historical account of the development of fodder beet in 1822 (cited in Ford-Lloyd et al., 1975), which provided the gene pool for the original sugarbeet (Fischer, 1989). Beet has been grown exclusively for sucrose only since the late 1700s, when it became possible to measure the sucrose present in the beet juice (Francis, 2006; McGrath and Fugate, 2012). Commercial-scale production of sucrose from beet was hastened in Napoleonic France, when that country was blockaded from its colonies and sucrose from sugarcane (Biancardi et al., 2010). The commercial beet crop harvested is the first year's vegetative growth of a biennial plant. Beet has a complex breeding system. Flowering requires 10 to 16 wk at 4°C (i.e., vernalization), at which time the rosette "bolts" a flower stalk with indeterminate inflorescences. Flowers are perfect and wind pollinated. Cross-fertilization is enforced by a complex self-incompatibility system (Biancardi et al., 2010).

Breeding has improved sugar yield markedly over the past 150 yr, from 4 to 6% at the onset of sucrose yield selection to >18% in modern hybrids and from a national average of just more than 20 Mg ha⁻¹ in 1909 to a current average more than 60 Mg ha⁻¹. Progeny testing and recurrent selection for sucrose content were employed, likely the first scientific application of these tools in common use today (Vilmorin, 1859, cited in Gayon and Zallen, 1998; Ware, 1880). Selection diversified the sugarbeet germplasm into types with a greater percentage of extractable sucrose (i.e., Z-type sugarbeet for "Zucker," the German word for sugar) and types with increased tonnage (i.e., E-type sugarbeet for "Ernte," the German word for harvest). Sucrose accumulates in sugarbeet roots to about 75% of dry weight (15–20% fresh weight). Sucrose biosynthesis occurs by mechanisms similar to those in other plants. Sucrose is concentrated within the innermost five of 10 to 12 concentric cortical rings, around the point of maximum root girth. It accumulates in vacuoles of parenchyma cells adjacent to the vascular tissue, and

the percent sucrose is positively correlated to the amount of solar radiation captured once the leaf canopy reaches full coverage (Elliott and Weston, 1993).

Doney et al. (1981) demonstrated that roots with smaller cells give higher sucrose percentages than those with larger cells. Savitsky (1940) suggested three or four major genes control sucrose percentage in crosses of sugarbeet with divergent types such as fodder and red beet (3–12% sucrose fresh weight each). Others suggest sucrose percentage is a quantitatively controlled trait with high heritability (Culbertson, 1942; Zhao et al., 1997). Recent mapping of sucrose genes suggests at least five loci on five of the nine beet chromosomes contribute to final sucrose content in a sugarbeet × sugarbeet population (Schneider et al., 2002). Root yield, on the other hand, is generally considered a nonadditive trait, and quantitative trait loci (QTL) analyses to date have shown only two loci with significant influence over yield (Schneider et al., 2002).

The improvement in commercial cultivars in percent sucrose over the past 50 yr, although considerable, has not been as great as the increase in root yield. Over the last 100 yr of sugarbeet production, an increase in both sucrose content, and especially, root yield has been achieved through genetic improvement and better crop management (Francis, 2006; Fig. 13–1). It is difficult to estimate, however, exactly how much of that improvement is due to genetic improvement.




Fig. 13–1. Changes in the area of sugarbeet harvested are shown for the entire United States from 1909 to 2012, as well as changes over time (1924–2012) in the area harvested in the western irrigated regions (California, eastern Oregon, Idaho, Montana, Wyoming, Colorado, and western Nebraska) and the eastern rainfed regions (North Dakota, Minnesota, and Michigan). The growth in average root yield per hectare is shown for the entire United States from 1909 to 2012 (right vertical axis). Data are from USDA-NASS. *"Other States" includes Arizona, Illinois, Indiana, Iowa, Kansas, Maine, New Mexico, New York, Ohio, South Dakota, Texas, Utah, Washington, and Wisconsin.

All commercial sugarbeet cultivars currently grown in the United States are hybrid, generated using a single, vulnerable, and complex system of cytoplasmic male sterility (CMS) (Owen, 1945). Modern cultivars also are monogerm (i.e., a single flower borne per axil), which eliminates the need for labor to thin stands, as was required for open-pollinated multigerm sugarbeet cultivars (Savitsky, 1950). New characters can be difficult to introgress into the inbreds that are hybrid parents. The pollinator parent of a hybrid often is selected out of a multigerm population that was selected for disease resistance. It can be easier to fix resistance alleles in pollinators than in the CMS (female) parent lines, because of the time invested in developing a maintainer line (O-type line) and female pair. It is necessary that the O-type maintainer be homozygous recessive at both of the loci that control the restoration of sterility in the female.

Current germplasm developed in U.S. public breeding programs and released to the seed companies is the product of several phases of development: early efforts to improve disease resistance in open-pollinated cultivars, followed by conversion of open-pollinated cultivars into genetic CMS parents for hybrid cultivar production (Biancardi et al., 2005); the mining of existing crop diversity to manage emerging diseases and achieve desired agronomic traits (e.g., reducing soil tare); and more recently the discovery and deployment of novel traits from crop wild relatives and unadapted genetic resources (Panella and Lewellen, 2007). Many sugarbeet breeding activities in the public breeding sector have been focused on improving populations via mass selection of mother roots, harvested and vernalized for subsequent seed production, and releasing the improved populations from these programs in the form of enhanced germplasm (Doney, 1995).

Foundations of U.S. Germplasm

Through World War I, most U.S. seed was imported from Europe or Russia as unadapted, open-pollinated cultivars, and the U.S. beet sugar industry struggled with recurring disease problems using these materials (Lewellen, 1992; Panella and Lewellen, 2007). The USDA's Division of Sugar Plants responded by the development of a sugarbeet research program that began almost 100 yr of collaboration that has been maintained among commercial, federal, and state researchers and plant breeders. Through this collaboration there was the development and release of cultivars resistant to *Beet curly top virus* for western growing regions (Coons et al., 1931) and later, for eastern growers, of cultivars resistant to aphanomyces seedling and root diseases (caused by *Aphanomyces cochlioides* Drechs.) (Coons et al., 1954). This distinction in germplasm persists today in modern hybrids (McGrath et al., 1999). Cercospora leaf spot (caused by *Cercospora beticola* Sacc.) also was a major focus for sugarbeet germplasm enhancement, mainly in areas east of the Rocky Mountains, and the development of this germplasm has been reviewed recently (Panella and McGrath, 2010). Development of the two U.S. germplasm pools has not been summarized recently, and since these are foundational germplasm materials, their development is briefly summarized here.

Beet curly top (BCT) in sugarbeet (caused by a mixture of closely related *Curtovirus* species: *Beet curly top virus*, *Beet mild curly top virus*, and *Beet severe curly top virus*) (Stenger and McMahan, 1997; Strausbaugh et al., 2008), transmitted by the beet leafhopper, *Circulifer tenellus* (Baker), is not found in European production areas (Bennett, 1971). Beet curly top severely impacted yields just as the sugarbeet

industry was beginning to expand in the western United States (Bennett, 1971; Carsner, 1924, 1926a; Murphy, 1946). This especially was true in California and Idaho (Fig. 13–2), where the beet leafhopper over-winters in weeds in the surrounding foothills. In California between 1910 and 1930 there was a decrease in root yield, sugar yield per acre, as well as a sharp drop in the area harvested (Duffus, 1978; Wisler and Duffus, 2000) (Fig. 13–3 and 13–4). Survival of the sugarbeet industry in the western United States depended on being able to manage this devastating disease (Bennett, 1971; Panella, 2005). In the 1920s, the USDA-ARS, then called the Bureau of Plant Industry) joined the private sector sugarbeet breeding efforts to develop cultivars resistant to BCT (Bennett, 1971; Bennett and Leach, 1971; Carsner, 1926b; Coons et al., 1931; Coons, 1953; Murphy, 1942, 1946). A series of joint releases between USDA-ARS and the sugarbeet industry began with the release of 'US 1,' a curly top resistant open-pollinated cultivar (Carsner, 1933; Owen et al., 1939). US 1 primarily had been derived from selections within European cultivars, the most important of which was R & G Old Type, also known as 'KleinE' (Lewellen 1992). Further increases in resistance to BCT resulted in 'US 33' and 'US 34' selected from heavily curly top infested fields of 'US 1,' and these eventually were replaced by 'US 12' and 'US 22,' which were further improved as 'US 22/2' and 'US22/3' (Coons et al., 1955).

Aphanomyces became a serious threat to eastern U.S. sugarbeet growing areas by the early 1940s, primarily causing a serious damping-off problem (black-leg), but also a mature root rot. Both remain problems, but their occurrence is limited and sporadic, mainly being a problem in warmer, wetter soils. Bockstahler observed differential reactions to the chronic form of black-leg in 1941 (Henderson and Bockstahler, 1946). In *A. cochlidioides* infested plots in Waseca, MN, a polycross cultivar designated as '1942 Minnesota Synthetic No. 1' was created, and additional populations were selected in subsequent years (Bockstahler




Fig. 13–2. Yield fluctuations from 1905 until 1930 seen in the areas of California and Idaho most affected by beet curly top (from Carsner, 1933, p. 2).

and Reece, 1948). Most aphanomyces-resistant germplasm was derived from selections within populations polycrossed in 1945 and 1946 from 'US 215,' 'US 216,' 'US 200,' 'R-581,' 'NYA,' 'LaRoyale,' 'Cesena,' 'US 22,' 'SL 561,' 'SL 611,' 'SL 618,' 'American 0-701-0,' 'American 1,' 'American 3,' 'Minnesota Synthetic No. 1,' as well as synthetics containing Italian varieties, *Beta maritima* [sic], and sucrose and yield selections. Much of this early germplasm had its roots in European open-pollinated cultivars, such as R&G Pioneer and Old Type, or in Great Western open-pollinated cultivars, such as GW304 and GW359, derived from some of these European sources (Lewellen, 1992).

Four hundred four roots from 48 progenies of this family, as well as five related families, were intercrossed in 1947 as '48B3-00.' The direct increase of this seed was released as 'US 1177'. Fifty-four roots of '48B3-00' were reselected in 1949, and named '50B3-0,' and a direct increase was released as 'US 400' in 1954 (Coons et al., 1954), which had better aphanomyces black-leg resistance than 'US 1177'. By 1955, enough commercial seed of 'US 400' was available to plant 12,150 ha (30,000 ac). Selections of 105 roots from '50B3-0' in nurseries at East Lansing, Blissfield, MI and Waseca, MN were interpollinated to produce '53AB3-0,' which was released in 1955 as 'US 401,' with greater aphanomyces and cercospora leaf spot resistance than 'US 400,' and replaced it commercially. A second important lineage of releases from these materials differed in the method of selection. That is, 'US 400' and 'US 401' were derived from mass selection after the 1947 progeny selection, but the alternate lineage relied on selection of the best individual roots in the best families (i.e., mother root selection) from progeny of '53AB1,' which was thrice selected from the 1946 Waseca selections for the highest combined black-leg, leaf spot, sucrose, and yield. These selections included the '02 clone' selected from '53AB1-32' (self-sterile but sib-fertile), which was released subsequently as 'EL 40' (Hogaboam et al., 1982). Collaborative work for cercospora leaf spot selection was centered in Beltsville, MD, where 'SP 5460-0' was released from '53AB1-65' and 'SP5481-0' from combined progenies of '53AB1,' and likely 'SP 5822-0.' SP 5822 likely was the progenitor of 'SP 6322,' which was the original pollinator of the historically important and widely cultivated hybrid 'US H20' (Coe and Hogaboam, 1971a,b).

Rhizomania is another yield limiting virus problem worldwide (Biancardi et al., 2002), caused by the *Beet necrotic yellow vein virus* (BNYVV), which is vectored by the plasmodiophorid, *Polymyxa betae* (Rush et al., 2006; Rush, 2003). Rhizomania infection causes a reduction of up to 50% in root yield and negatively affects sugar yield and processing quality (Büttner et al., 1995). In the United States, the virus first was discovered in 1983 in California (Duffus et al., 1984; Biancardi et al., 2002). The first documented and widely used source of resistance to BNYVV was the gene, *Rz1* (Lewellen et al., 1987). *Rz1* is monogenic and although the dominant allele does not confer immunity, it greatly reduces damage and symptom expression. *Rz1* has been deployed in backcross and germplasm enhancement programs worldwide and has become the major management tool against yield loss by BNYVV (Asher et al., 2002; Francis et al., 1998; Pavli et al., 2011; Rush et al., 2006; Scholten et al., 1997).

Shortly after the discovery of *Rz1*, an additional resistance gene was found in a sea beet (*Beta vulgaris* subspecies *maritima*) accession (WB42-PI 546385), designated *Rz2* (Lewellen, 1997; Scholten et al., 1999; Whitney, 1989). Cultivars with the *Rz1* gene began showing rhizomania symptoms in some fields in the Imperial

Valley of California in the early 2000s. Extensive testing in 2004 and 2005 by USDA-ARS at Salinas, confirmed that resistance conditioned by the *Rz1* gene had been overcome (Liu et al., 2005; Liu and Lewellen, 2007; Rush et al., 2006). The *Rz2* source of resistance, alone or in tandem with *Rz1*, has since become more common in commercial cultivars. Other sources of rhizomania resistance that have been identified are *Rz3* from WB41 (Gidner et al., 2005; Lewellen et al., 2007), *Rz4* derived from a composite of about 60 sea beet accessions from Europe and the Middle East (Grimmer et al., 2007), and *Rz5* from WB258 (PI 546426) (Grimmer et al., 2008). There is debate as to whether or not these sources of resistance represent individual genes or simply alleles at the same locus, but all may be needed to maintain genetic resistance to rhizomania (McGrann et al., 2009).

The U.S. beet seed industry initially was a cooperative effort among sugar factory districts, which always had conducted official cultivar trials for their growers; public breeders (USDA-ARS), who develop enhanced germplasm; private breeders working for U.S. seed companies, often owned by the sugar processors; and approved cultivar multiplication, primarily done by the West Coast Beet Seed Co. of Salem, OR (Johnson and Burt, 1990). With the advent of hybrids, the commercial opportunities afforded by hybrid seed sales, and the reentry of large international seed companies into the U.S. market, a process which began again during the 1970s, public sector breeders have focused more on germplasm enhancement, while commercial sugarbeet seed company breeders have come to dominate cultivar development. Along with the recent assimilation of many small seed companies into larger corporations, this has led to a few companies supplying all of the seed to U.S. growers. This has been an important aspect in rising yields due to their affiliation with international seed companies, which have diverse germplasm collections and genetic marker and genetic engineering capabilities.

Some biolistic- and *Agrobacterium*-mediated transformation of sugarbeet has occurred in the public sector, but this technology is largely subsumed by the commercial sector, and the only genetically engineered trait currently commercialized is glyphosate [*N*-(phosphonomethyl)glycine] resistance for weed control. This trait initially was deregulated in 1998, but seed from that transformation event was never commercialized. The current glyphosate-resistant sugarbeet resulted from another transformation event (H7-1, U.S. Patent 7335816 B2) and was deregulated in March 2005, grown on a limited basis in Wyoming in 2007, and released across the United States in 2008. Commercial adoption rates quickly reached >98% by 1 yr after its official initial deregulation. A series of legal challenges resulted in partial re-regulation in 2010 and finally full deregulation in 2012. Although other bioengineered traits of interest are in development (varied disease resistances, increased water and nitrogen use efficiency, and winterbeet for more temperate climates), none have been commercialized.

Cultivation and Beet Production

Sugarbeet cultivation came to this country with German and Russian immigrants, who were familiar with the crop from their homelands. After the U.S. Civil War and the abolition of slavery, there was concern that the United States might not remain self-sufficient in sugar production. In addition, the potential economic impact of a northern sugar industry was also an attractive proposition in itself (Grant, 1871). Between 1838 and 1879, 14 beet sugar factories were built—all of

which failed (Theis, 1971). Finally in 1879, the factory built in Alvarado, CA in 1870 turned a profit. The industry then began to expand and by 1909, there were about 162,000 ha (400,000 ac) of sugarbeet harvested in the United States, with an average yield of about 22.4 Mg ha⁻¹ (10 t ac⁻¹), with 10% sucrose, processed in 41 factories (Harris, 1919; USDA-NASS, 2013) (Fig. 13–1). From 1909 to present (Fig. 13–1), maximum acreage (1969) was about 607,500 ha (1.5 million acres) and currently varies between 400,000 and 510,000 ha (1.00 and 1.25 million acres). Both root yield (Fig. 13–1) and the percentage of sucrose in the root have risen (Francis, 2006), as has the capacity of the 21 currently operating sugarbeet processing factories to process sugarbeet. Today, U.S. farmers and processors are producing much more sugar on less land.

By 1924, when the USDA-NASS began keeping records for most crops, sugarbeet was being cultivated in 20 states. The U.S. sugarbeet crop grew as more land was brought into cultivation in more areas; sugarbeet was cultivated at one time or another in 28 states ranging from California to New York and from Maine to the semiarid southwest. Today, sugarbeet is cultivated in 10 states that often are grouped into five production regions: the West (California), the Intermountain Region (Idaho and eastern Oregon), the Great Plains (Colorado, Wyoming, Nebraska, and Montana), the Upper Midwest or Red River Valley (Minnesota and North Dakota), and the Great Lakes (Michigan). In Fig. 13–1, the area harvested data are shown for “irrigated” (West, Intermountain, and Great Plains regions) and rainfed (Upper Midwest and Great Lakes regions) areas from these 10 states, with the historical production from all the other states combined. It is clear from the changes in the area harvested (Fig. 13–1) that production was initially highest in the western United States, where the sugarbeet crop was irrigated, and that as the area under cultivation in the Red River Valley and Michigan increased, the area cultivated for sugarbeet peaked at the end of the 1960s, when the production in states other than the current 10 states began to fall. Disease pressure in some growing areas, fluctuations in environmental conditions, and economic fluctuations in the price of sugar pushed many areas out of production. A serious lack of investment in factories to keep them efficiently operating with the latest technology was also a major factor. Production also migrated to the rainfed areas, where cost of production is lower by avoiding the need to irrigate. As the area under cultivation has dropped, yields have continued to rise, maintaining production goals (Fig. 13–1).

In the United States root yield was variable and did not increase much during the 20 yr from 1950 to 1970 (Fig. 13–1). However, since then the development and use of superior cultivars (resistance to BCT, bolting, *Rhizoctonia solani* Kühn, *Cercospora*, downy mildew [*Peronospora farinosa* (Fr.:Fr.) Fr. f. sp. *betae* Byford] resistance, and other pests and diseases), the introduction of hybrid cultivars with monogerm seed, greater use of new chemical protectants, better harvesting equipment, and more advanced fertilization management have led to a steady increase of yield in the United States. This increase was mirrored in much of Europe (Zimmermann and Zeddies, 2000). Breeders and producers in each of the U.S. growing regions share best ideas for cultural practices and continued improvements in germplasm for cultivar development; however, each growing region has a unique set of challenges that often differs from other regions in the magnitude of the problem. For instance, curly top and yellowing virus problems are mainly confined to the western U.S. growing regions, fungal leaf spot and root rots are more

problematic in the east, and rhizomania is a very serious problem in all growing regions. Thus, solutions achieved in areas of severe impact are transferred throughout the United States (and worldwide) as the situation demands.

California—The Western Region

Although the initial attempts to grow sugarbeet in the United States were made in the eastern part of the country as early as 1838 (Grant, 1871), it was in California that sugar production from beet first became successful. After a number of failures, E.H. Dyer was successful in establishing a factory that made a profit by 1879. This factory in Alvarado, CA is considered the first successful U.S. factory. In 1888, Claus Spreckels built a factory at Watsonville, CA, which produced 907 Mg (1000 t) of sugar in the first year, and by 1889, there were two sugarbeet processing factories operating in the United States, both in central California (Harris, 1919; Theis, 1971). By 1899 California was the leader in beet sugar production, with 37% of the U.S. cultivated area. There was a severe drought from 1897 to 1900, and then tough economic times, which caused dramatic fluctuations in the production and cultivation of sugarbeet in California (Souder, 1970). Nonetheless, by 1910, the area under cultivation had grown to about 36,650 ha (90,500 ac), with a yield of about 24.48 Mg ha⁻¹ (11.08 t ac⁻¹) (Fig. 13–3 and 13–4). Over the next 20 yr, the disease pressure from BCT caused a steady drop in root yield, until the first curly top-resistant cultivars became available in the early 1930s (Fig. 13–3 and 13–4). Beet curly top affects the percentage of sucrose in the root as well, causing an even more drastic drop in gross sugar yield over the area harvested (see Fig. 1 of Duffus, 1978; Wisler and Duffus, 2000). With resistant cultivars available to




Fig. 13–3. Changes in the area of sugarbeet harvested are shown for the state of California from 1924–2012. Data are from USDA-NASS.

manage BCT, and improved crop cultural practices, the industry rebounded, and the area under cultivation, root yield, and gross sugar yield grew over the next 20 yr (Fig. 13–3 and 13–4) (Duffus, 1978; Wisler and Duffus, 2000) until there were nine sugarbeet processing factories active in California (United States Beet Sugar Association, 1936).

In 1951, the virus yellows complex was reported in California for the first time (Coons and Kotila, 1951). *Virus yellows* is term used to describe a complex of yellows-inducing viruses including, *Beet western yellows virus*, *Beet chlorosis virus*, *Beet yellows virus*, *Beet mosaic virus* and other aphid-transmitted viruses. With heavy infections of more virulent strains, root yield reductions of more than 50% and losses of up to two percentage points of sucrose have been reported (Bennett et al., 1957). Therefore, although U.S. yields trended upward (Fig. 13–1), it was clear that virus yellows limited yield progress in California, much as curly top had depressed yields three decades earlier (Fig. 13–4). The cycle of infection of yellowing viruses was sustained by continual (year-round) beet crops in the areas most affected. With both fall-sown and spring-sown crops, aphid vectors could move the yellowing viruses from one crop to the next. This disease transmission cycle was broken when growers organized regionally to isolate old beet fields from young fields by “beet-free” periods in diverse regions of the state (Wisler and Duffus, 2000). With this cycle broken, the new, higher yielding hybrids; increased access to irrigation water; and economic changes, which favored sugarbeet being grown on more productive land, led to increased yields (Duffus, 1978).




Fig. 13–4. Changes in the average root yield are shown for the state of California from 1910 to 2011. Linear regression lines show yield trends that were present over different time periods. Data are from USDA-NASS, California Field Office.

In addition to the increasing area infested with BNYVV during the 1980s and early 1990s, a number of other problems affected California sugarbeet production. Changing economics of growing sugarbeet, sugarbeet cyst nematode (*Heterodera schachtii* Schmidt) pressure, and a number of whitefly (*Bermisia* spp.) vectored diseases impacted both yields and acreage (Wisler and Duffus, 2000). The less productive areas in northern California were the first to lose sugarbeet in their rotations. California had about 25% of the U.S. sugarbeet area under cultivation in the 1960s and 1970s, but by the end of the 1990s it represented only 10% of the U.S. acreage, and today there only are about 10,100 ha (25,000 ac) in the Imperial Valley, where sugarbeet is grown as a fall-planted, summer-harvested crop (Fig. 13–3) (Wisler and Duffus, 2000).

Imperial Valley of California

The Imperial Valley of California deserves a brief discussion because it is a unique growing area for sugarbeet in the United States and has the highest gross sugar yields per area in the world. Sugarbeet has been grown in the Imperial Valley since 1932, shipped by rail to factories elsewhere in California or Chandler, AZ. The currently operating processing factory in Brawley was built by Holly Sugar Corporation at the end of the World War II and opened in 1947. Shipping of a portion of the crop to other factories continued from then until the early 1990s, when rail shipment became uneconomical or factories closed. The factory in Brawley is the last one operating in California, where 24 (11 successful) factories have been constructed, operated, and eventually closed since the first successful commercial sugarbeet factory was opened in 1870 near San Jose (Harris, 1919; Souder, 1970).

The Imperial Valley is a low desert ecosystem with mild, sunny winters and hot summers. It receives the most sunlight of any location in the United States. Most soils are high in pH and calcareous, with large amounts of naturally occurring gypsum. Most Imperial Valley fields are tile drained at 1.8 m (6 ft) in depth. Subsurface drainage maintains positive salt balances in the Imperial Valley soils, which otherwise would salinize. Water delivery to the Imperial Valley and power production is provided by the Imperial Irrigation District, first organized in 1911. The All American Canal conveys approximately 3.2074 billion m³ (2.6 million ac ft³) of water per year to 172,125 ha (425,000 ac) of farmland and seven cities and towns. The canal runs along the Mexican border to the south. Water is then diverted through a series of canals northwards to towns and fields. Because the land slopes northwards toward the Salton Sea, little power is needed to move water. Imperial Irrigation District holds some of the most senior water rights to the Colorado River. The result for growers has been a relatively constant supply of water at low prices (currently ~\$20 per 1233.6 m³ [ac ft³]) in comparison to many other irrigation districts in California.

Unlike all other U.S. growing regions where sugarbeet is a spring-planted crop, the Imperial Valley sugarbeet crop is planted from the middle of September to early October, and harvest begins ~180 d later starting in early April. Harvest usually is concluded in July, but there has been a tendency in recent years to extend the harvest later in the month, often running into August. This has occurred due to increasingly larger sugarbeet yields, and at times due to operating difficulties in the factory. Harvest campaign length has tended to increase over the last 25 yr, but is not solely responsible for increased yields over that period. Examining the period from 2007, when all sugarbeet production in California was in the

Imperial Valley (Fig. 13–5) compared to the California production since 1989 (Fig. 13–6), it is clear how much higher yields in the Imperial Valley have been. In 2012 in the Imperial Valley, another world-record sugarbeet crop was harvested on a 28.35-ha (70-ac) field. That field averaged 176.31 Mg ha⁻¹ (78.71 t ac⁻¹) with 15.89% sugar (fresh weight) for a gross sugar yield of 11,356.4 kg ha⁻¹ (25,014 lb ac⁻¹) (Lil-leboe, 2012).

Yields in the Imperial Valley of California continue to increase. Sugarbeet crops increase tonnage during the spring period (April–June) at a greater rate than from establishment in fall to start of harvest in April. Dry matter more than doubles by the end of the last 4 mo of growth in late July and early August, compared to the period from establishment in September until the start of harvest in April. A number of factors have combined to contribute to larger yields: improved performance of new hybrids, including improved rhizomania resistance; reduction in losses to *Lettuce infectious yellows virus* starting in the early 1990s; better and wider selection of herbicides for the control of weeds (glyphosate-resistant cultivars had not been grown in the Imperial Valley, but they have been approved beginning in the 2013 growing season); the adoption of newer style harvesters allowing for harvest from wetter soil, permitting irrigation closer to harvest, and better management and timing of late season irrigations during the hottest weather; improved seed quality and stand establishment practices; and a partial shift to production on better quality soils in recent years due to recent increases in the profitability of sugarbeet. These factors have interacted to increase beet dry matter accumulation in the April to June period at a greater rate than observed during previous decades. During the same period, sugar concentration in roots




Fig. 13–5. Increased average root yield per hectare in the Imperial Valley of California is shown from 2007 to 2012. Data are from the California Beet Growers Association.

has remained relatively stable, so increased root yields have been primarily responsible for increased sugar yields.

The rate of increase in root yield and decrease in sugar percentage both accelerate over time during the growing season in the IV. This is due to increasing temperatures resulting in several disadvantages as the season progresses. Year-round mild to hot temperatures makes it impossible to store sugarbeet in piles, as is done in other beet-producing regions with spring sown, fall harvested crops. In the IV, logistics are managed to process sugarbeet within 24 h of harvest, with the factory running 24 h, 7 d a week from April to July. Temperatures, especially night time temperatures, and correlated respiratory sugar losses increase as the season progresses, the costs of insect pest management and correlated losses increase, larger amounts of irrigation water are needed, and damage from nematodes and root rots becomes more common, especially in July and August. To operate the factory for a long enough season, late season inefficiencies and costs (especially water use) must be tolerated. Besides future genetic improvements (including the use of transgenic traits), the most likely technology that could help reduce losses late in the season and improve overall efficiency may be the adoption of drip irrigation technology.

Intermountain Region (Idaho, Oregon)

Sugarbeet processing factories operated in the Intermountain Region (Idaho, Utah, eastern Oregon, and eastern Washington) beginning in the late 1800s, and by 1908 the Utah-Idaho Sugar Co. and the Amalgamated Sugar Company had




Fig. 13-6. Changes in the average root yield per hectare are shown for the state of California from 1989 to 2012. The line represents the linear regression through these averages. Data are from the California Beet Growers Association.

established themselves as the largest beet sugar processing companies in the region. In the early 1920s, the Utah-Idaho Sugar Co. had 20 factories in four states (ID, UT, WA, OR), and Amalgamated Sugar Co. had eight factories in Idaho and Utah. During the 1920s and 1930s, a number of factories closed or ceased operation because of drought and curly top, which caused large fluctuations in yield (Fig. 13-2) and area cultivated for sugarbeet (Fig. 13-7). Primarily due to BCT, by 1934, Amalgamated had only two operating factories. As BCT-resistant cultivars became available in the early 1930s, yields improved, and Amalgamated resumed operation of five of eight factories. In 1966, the Utah-Idaho Sugar Co. had only five sugar factories left and by the late 1970s had closed all operations. In 1994, the Amalgamated Sugar Co. became a farmer owned cooperative (renamed Snake River Sugar Co.). In 2005, the Nyssa, OR factory was closed leaving only the Nampa, Paul, and Twin Falls factories still operating in the Intermountain region (in Idaho).

Commercial sugarbeet production areas in Idaho, Oregon, and Washington are all coordinated through the Snake River Sugar Co. During 2012, there were 74,058 ha (183,000 ac) (Fig. 13-7) planted in Idaho, all of which were in the southern portion of the state, where over the last 30 yr, there has been a shift in acreage from Treasure Valley (southeast Oregon and southwestern Idaho) to Magic Valley (south-central Idaho) and eastern Idaho. In Oregon, commercial production is concentrated in the eastern portion of the state, where 4411 ha (10,900 acres) were harvested in 2012. (Most seed production for sugarbeet in the United States also occurs in Oregon, but is concentrated in the Willamette Valley, west of the Cascade mountain range.) Sugarbeet production was widespread in central




Fig. 13-7. Area planted to sugarbeet in Idaho and the Intermountain region (Idaho, Oregon, and Washington) from 1924 to 2012 based on statistics from the USDA-NASS.

Washington until a major reduction in acreage occurred in the mid-1970s with the closure of the factory at Moses Lake, WA (Fig. 13–7). The USDA-NASS survey last reported data for Washington in 2008 with only 647 ha (1600 ac). Although there still was a small acreage in Washington in 2012, it was too small to be included in the USDA-NASS survey.

Most production fields in this area are prepared by plowing with a mold-board plow, roller harrowed (perhaps more than once), and then marked and/or bedded. Some growers will “drag off” the beds at planting to expose moist soil. Planting in lower elevation areas (610 m [2000 ft]) occurs as early as the first week of March, while higher elevation (1676 m [5500 ft]) areas typically are planted in mid to late April. The crop was thinned by hand or mechanically until the early 1990s, but now is planted to stand (14 cm [5.5 in] between plants). Before 2008, weed control was accomplished through tillage and about a dozen herbicides; however, since the release of glyphosate-resistant sugarbeet, weed control has shifted to the use of glyphosate. The use of glyphosate has eliminated problems associated with herbicide toxicity to sugarbeet seedlings and weather interfering with timely herbicide application. About 4000 ha (10,000 ac) are currently planted using strip tillage (20–30 cm wide), an additional management tool afforded by glyphosate-resistant sugarbeet, which is particularly useful in windy areas with sandy ground prone to soil erosion and plant damage in the spring at planting time and crop emergence. The entire crop is irrigated, but the method of irrigation has changed considerably over the last 30 yr, with a 48% reduction in gravity irrigation and a 49% increase in sprinkler irrigation (Fig. 13–8) due to reduced




Fig. 13–8. Method of irrigation (gravity or sprinkler) on crop land in Idaho from 1984 to 2008 based on statistics from the USDA-NASS.

labor costs, higher irrigation efficiency, less soil erosion and nutrient loss, and government programs that helped offset the cost of the equipment change.

The primary disease problems in this production area include two virus diseases (BCT and rhizomania), various root rots (*R. solani* Kühn and some bacterial rots), and powdery mildew (*Erysiphe polyfoni* DC). As in the rest of the United States, rhizomania is managed through the use of resistant cultivars. The use of chemical protectants often provides the yield stability needed to introduce new genetic improvements throughout the entire commercial genepool (e.g., rhizomania resistance or glyphosate resistance), while the previous array of disease resistance is being reconstituted.

A survey in the western United States in 2006 and 2007 indicated the primary species of BCT were *Beet severe curly top virus* and *Beet mild curly top virus* (Strausbaugh et al., 2008). Since the mid 1930s host resistance has been the primary control measure for this problem, but in 2006 clothianidin ([C(E)]-N-[(2-chloro-5-thiazolyl)methyl]-N'-methyl-N'-nitroguanidine; Poncho, NipsIt), a neonicotinoid insecticide seed treatment, became available for sugarbeet (Strausbaugh et al., 2006). The clothianidin seed treatment along with thiamethoxam (3-[(2-chloro-5-thiazolyl)methyl]tetrahydro-5-methyl-N-nitro-4H-1,3,5-oxadiazin-4-imine, marketed as Cruiser) has become a valuable supplement to host resistance through control of the beet leafhopper vector (Strausbaugh et al., 2006, 2012). In Idaho, yields with neonicotinoid seed treatments averaged $72.90 \pm 3.79 \text{ Mg ha}^{-1}$ ($32.52 \pm 1.69 \text{ t ac}^{-1}$) from




Fig. 13-9. Average root yield trends over the last 30 yr for Idaho sugarbeet production based on statistics from the USDA-NASS. Some of the significant events during this timeframe that likely influenced yields were the introduction of rhizomania, which is a viral disease caused by *Beet necrotic yellow vein virus* (BNYVV), release of rhizomania resistant varieties based on the *Rz1* gene, neonicotinoid seed treatments for curly top vector and pest control (Poncho), and glyphosate resistant varieties.

2006 to 2010, a 17% increase when compared to the preceding 5 yr 60.57 ± 4.51 Mg ha⁻¹ (27.04 ± 2.01 t ac⁻¹) (Strausbaugh et al., 2012). In 2008, glyphosate-resistant cultivars also were introduced but did not seem to lead to immediate yield increases, although weather may have been a factor, as well (Fig. 13–9). Idaho yields for 2006–2007 averaged 74.09 ± 4.28 Mg ha⁻¹ (33.05 ± 1.91 t ac⁻¹) and the first 3 yr (2008–2010) with glyphosate yields averaged 72.12 ± 4.15 Mg ha⁻¹ (32.17 ± 1.85 t ac⁻¹) (Fig. 13–9). In the first commercial glyphosate-resistant cultivars, the level of general disease resistance was not as high as in the previous cultivars. For example, the curly top resistance in some of the initial glyphosate cultivars tested before 2008 was worse than the susceptible check at the later evaluation dates (C.A. Strausbaugh, 2005, data not shown). Thus, the neonicotinoid seed treatments have helped make the transition to glyphosate-resistant cultivars possible (Strausbaugh et al., 2012). Since the initial introduction of glyphosate-resistant cultivars, the level of resistance to curly top and resistance to other diseases has improved steadily. With this improved disease resistance and favorable weather, yields again have increased in Idaho during 2011 [77.11 Mg ha⁻¹ (34.4 t ac⁻¹)] and 2012 [79.13 Mg ha⁻¹ (35.3 t ac⁻¹)] (Fig. 13–9).

Another major concern in the Amalgamated production area has been various root rots, such as rhizoctonia crown and root rot (Strausbaugh et al., 2011; Strausbaugh and Gillen, 2009), bacterial root rot (*Leuconostoc mesenteroides* and others) (Strausbaugh and Gillen, 2008), and aphanomyces root rot (caused by *Aphanomyces cochlioides* Drechsler). Rhizoctonia root rot of mature roots is the primary problem, and is caused by *R. solani* AG-2-2 IIIB strains (Bolton et al., 2010; Strausbaugh et al., 2011). However, the fungus initially seems only to penetrate the outer portion of the root (~5% of the root mass) and subsequent bacterial invaders led by *Leuconostoc mesenteroides* can act synergistically to rot the majority of the root mass (Strausbaugh and Eujayl, 2012; Strausbaugh et al., 2011). The Rhizoctonia–bacterial root rot complex is worse in the warmer longer growing season portion of the production area (Strausbaugh et al., 2011). The cooler, higher elevation area in eastern Idaho has very little problem with root rot. Cultivars with resistance to *R. solani* are available in limited numbers, but banding on fungicides, crop rotation, and careful irrigation will continue to be necessary management practices (Bolton et al., 2010; Buhre et al., 2007; Kiewnick et al., 2001; Kirk et al., 2008; Rush and Winter, 1990; Strausbaugh et al., 2013a,b; Stump et al., 2004; Windels and Brantner, 2005). Aphanomyces root rot only recently has become a concern in eastern Oregon and now has been identified in a number of locations in southern Idaho. The management of these root rotting diseases will be critical for continued yield increases in this production region.

The primary pests limiting sugarbeet yield in the Amalgamated production region include: beet leafhopper (vector for curly top), root maggot [*Tetanops myopaeformis* (Roder)], wireworms [*Limonius californicus* (Mann.) and others], cutworms (*Agrotis* spp. and others), root aphid (*Pemphigus betae* Doane), black bean aphid [*Aphis fabae* (Scopoli)], beet leafminer [*Pegomya betae* (Curt.)], and spider mites (*Tetranychus urticae* Koch) (Strausbaugh et al., 2010). The neonicotinoid (clothianidin and thiamethoxam [3-[(2-chloro-5-thiazolyl)methyl]tetrahydro-5-methyl-*N*-nitro-4*H*-1,3,5-oxadiazin-4-imine]) seed treatments provide good early season control of nonchewing pests, but also will provide considerable protection from chewing pests (Strausbaugh et al., 2010). Systemic and foliar insecticides can be used to manage pest problems where the seed treatments are weak. A limited number of commercial cultivars with root aphid resistance also are

available. The loss of aldicarb (2-methyl-2-(methylthio)propanal *O*-[(methylamino) carbonyl]oxime) reduced the choice of systemic insecticides and also raised concerns over sugarbeet cyst nematode control. Thus, there is a push to increase the number of commercial cultivars with nematode tolerance available. Currently each seed company has a few cultivars available.

Early season harvest (10% of crop) usually begins in mid September in all areas. Harvest in the longer season, lower elevation areas frequently stretches into November, while other areas try to complete harvest in October. In 2012, Idaho ranked second for total yield (5.829 million tonnes [6.425 million tons]) by state, while Oregon was ranked last (379,203 Mg [418,000 tons]) (Fig. 13–7). In this production area, approximately one-third of the crop is processed directly through the factory, while another third remains in short-term storage (<90 d) under ambient conditions outdoors and the remaining one-third is placed in long-term storage (>90 d) (Peterson et al., 1984). Piles for long-term storage typically are covered and ventilated, and there also is some indoor storage. Piles are cooled with ambient air and prolonged rainy, unusually warm weather in the winter months can have serious, deleterious effects on the quality of the sugarbeet in storage.

Great Plains Region (Colorado, Montana, Nebraska, Wyoming)

The first mention of sugarbeet as a potential crop in Colorado was in the mid 1800s. Because the crop could not be brought from food to table, it had not come with the first settlers, and a certain level of economic development had to be present to provide the capital for the technology and facilities to refine sucrose from the sugarbeet. Nonetheless by the late 1800s, Colorado State Agricultural College (now Colorado State University) had shown in test plots the suitability of the crop to the soils and climate of the Great Plains Region (Hamilton, 2009). The first sugar extracting factories were built right at the beginning of the 20th century, and the Great Western Sugar Company was founded in 1905. In 1907 there were 16 factories operating in Colorado, and by 1926, 22 factories were active in Colorado (Hecker, 2004), and there were more than 81,000 ha (200,000 ac) of sugarbeet under cultivation (Fig. 13–10). Great Western Sugar Company became a major force in the sugar industry and owned 26 factories in six states (Kansas and Ohio in addition to the Great Plains states). It also had a strong research and seed cultivar development program that was very active in the Great Plains.

All sugarbeet grown in the Great Plains region is irrigated, and the climate is excellent for sugarbeet production, with long days, and cool nights due to the altitude. Nonetheless, there are a number of diseases that have had a severe effect on production in the Great Plains. Curly top certainly was a problem in some years and could depress yields across the Great Plains region, especially in the hot dry years of the Dust Bowl; even today, it still can be severe, especially during periods of early season drought. The years until the BCT-resistant cultivars were released showed lack of yield gain (Fig. 13–11) and yield fluctuations in the Great Plains but not as severe as the up and down cycles seen in California and Idaho (Fig. 13–2 and 13–4).

In the early 1920s, breeding efforts were initiated to find sources of resistance to cercospora leaf spot, which was a problem in the Great Plains, especially in the Arkansas Valley and all of eastern Colorado and Nebraska (Skuderna, 1925), where night temperatures were higher, as was the relative humidity. It also was




Fig. 13–10. Area of sugarbeet harvested in Colorado, Montana, Nebraska, and Wyoming from 1924 to 2012 based on statistics from the USDA-NASS.




Fig. 13–11. Average root yield for Colorado, Montana, Nebraska, and Wyoming from 1924 to 2012 based on statistics from the USDA-NASS.

a serious problem in Michigan and Ohio (Dahlberg et al., 1940). Resistant lines appeared beginning in 1937 with the release of 'US 217' and continued with a series of synthetic cultivars: 'US 200 × 215,' 'US 215 × 216,' 'US 216 × 225,' which were widely grown in these areas and alleviated to some extent the serious damage done by this disease (reviewed by Panella and McGrath, 2010). From the end of World War II until the early 1980s, there was a general increase in yield as hybrid sugarbeet became available and cultivars with monogerm seed were introduced (Fig. 13–11). However, during this period there was a tremendous decrease in the area cultivated in Colorado (Fig. 13–10), and although area cultivated in the other Great Plains states remained more constant, there has been a general decrease to reach today's levels in this region. In Colorado the decline in sugarbeet cultivation is directly tied to the slow demise of the Great Western Sugar Company, which was a result of the retirement of seasoned management after being sold in 1967 to a less experienced chief executive (Hamilton, 2009). The purchase by the Hunt brothers in 1974 and descent into bankruptcy in 1984 ended the Great Western Sugar Company. It was purchased by Tate & Lyle in 1985, but because growers were not paid timely in 1984 for their crop, in 1985 only about 1000 ha (2500 ac) of sugarbeet was planted. Finally, in 2002, the processing company was sold to the growers in the four state region, and it has been stabilized as the Western Sugar Cooperative (Hamilton, 2009).

The decrease in area cultivated has had severe agronomic consequences, which include the shortening of rotations, leading to increased disease pressure. This was exacerbated by drought in the late 1990s and early 2000s and the appearance of rhizomania in the early 1990s. Rhizomania was confirmed in Colorado and the Great Plains region by 1995, and it is managed almost exclusively with resistant germplasm. Along with resistance to rhizomania, came cultivars with improved resistance to sugarbeet cyst nematode, rhizoctonia root rot, fusarium wilt (caused by *Fusarium oxysporum* f. sp. *betae*), aphanomyces black root, cercospora leaf spot, BCT, and sugarbeet root aphid. During the last 10 yr, there also has been an increase in efficacy and diversity of protective chemical available for disease management. (For many soil borne pests and diseases, fumigation, especially with 1,3-Dichloropropene [Telone II], had been used in the past to prepare for planting sugarbeet. Although effective, there is a high cost to the grower, economically and environmentally.) These improvements and the introduction of glyphosate-resistant cultivars have led to the steady rise of yields over the past 10 yr in the Great Plains of about $1.90 \text{ Mg ha}^{-1} \text{ yr}^{-1}$ ($0.85 \text{ t ac}^{-1} \text{ yr}^{-1}$) (Fig. 13–12). Yields have risen from a four state average of 42.6 Mg ha^{-1} (19 t ac^{-1}) to an average of almost 67.2 Mg ha^{-1} (30 t ac^{-1}) (Fig. 13–12), and although the sucrose percentage has not risen appreciably, it has remained constant.

Upper Midwest Region (Eastern North Dakota, Western Minnesota—Red River Valley)

Sugarbeet has been raised in the Red River Valley (RRV) Region of eastern North Dakota and western Minnesota since 1918 when the first sugarbeet crop was planted on a farm near Crookston, MN. The sugarbeet raised on this small acreage (under contract from the Minnesota Sugar Company) was sent by railcar to Chaska, MN for processing. Over the next few years, the area under cultivation grew steadily (Fig. 13–13), until there was enough sugarbeet production in the RRV to justify building a processing facility near East Grand Forks, MN




Fig. 13-12. In the past 10 yr average yield in the Great Plains region (Colorado, Nebraska, Montana, and Wyoming) has risen from an average of 42.6 Mg ha⁻¹ (19 t ac⁻¹) to an average of almost 67.2 Mg ha⁻¹ (30 t ac⁻¹) at a rate of about 1.90 Mg ha⁻¹ yr⁻¹ (0.85 t ac⁻¹ yr⁻¹). Data are from USDA-NASS.




Fig. 13-13. Area planted to sugarbeet in the Upper Midwest region (Red River Valley, North Dakota and Minnesota) from 1924 to 2012 based on statistics from the USDA-NASS.

(completed in 1926). Processing plants in Crookston and Moorhead, MN soon followed in 1948 and 1954, respectively, as sugarbeet acreage continued to expand rapidly, and another factory was built in 1965 near Drayton, ND.

Since the mid 1970s, when three independent grower-owned sugar cooperatives (American Crystal Sugar Company, which purchased the cooperative at Hillsboro after 1 yr of operation; Minn-Dak Farmers Cooperative; and Southern Minnesota Beet Sugar Cooperative) were formed, the RRV has become the largest production area of sugarbeet within the United States. This particular region represented 32% of the U.S. area cultivated to sugarbeet in 1978 (170,455 ha [421,200 ac]) (Fig. 13–13) and has increased steadily over the past three decades to 276,400 ha (683,000 ac), which was 56% of U.S. area sown during the 2012 growing season (USDA-NASS, 2013). One of the main reasons for the increased acreage and continued popularity of this specialty crop is the vital role that sugarbeet production plays in the regional economy. Expansion of acreage also is due to development of long-term storage technology that allows these factories to have the longest processing campaigns in the world—200 d and sometimes more than 250 d in length. The total economic activity (direct and secondary impacts) generated by the sugar industry of the RRV was nearly \$4.9 billion in 2011 alone; or expressed alternatively, each hectare of sugarbeet planted generated about \$18,542 (\$7,507 ac^{-1}) in total local business activity (Bangsund et al., 2012).




Fig. 13–14. Average yield per harvested area in the Upper Midwest region has increased $1.08 \text{ Mg ha}^{-1} \text{ yr}^{-1}$ ($0.48 \text{ t ac}^{-1} \text{ yr}^{-1}$) in commercial fields and $0.74 \text{ Mg ha}^{-1} \text{ yr}^{-1}$ ($0.33 \text{ t ac}^{-1} \text{ yr}^{-1}$) in the official variety trials over the last 24 yr. Data are from American Crystal Sugar Company, Minn-Dak Farmers Cooperative, and Southern Minnesota Beet Sugar Cooperative.

During the last 25 yr, the sugar industry in the RRV has seen a steady upward trend in overall crop yield and quality. When looking specifically at crop yield, total yield per harvested area has increased $1.08 \text{ Mg ha}^{-1} \text{ yr}^{-1}$ ($0.48 \text{ t ac}^{-1} \text{ yr}^{-1}$) (Fig. 13–14) in commercial fields and $0.74 \text{ Mg ha}^{-1} \text{ yr}^{-1}$ ($0.33 \text{ t ac}^{-1} \text{ yr}^{-1}$) in the Official Variety Trials (replicated field trials established for the evaluation of varietal performance and approval of sale within each cooperative). Combined with an increased level of quality, these increases equate to a 183 and 148 $\text{kg ha}^{-1} \text{ yr}^{-1}$ increase in recoverable white sugar per hectare for commercially grown fields and the official variety trials, respectively (Fig. 13–15). These average annual gains are a reflection of the increased genetic potential of the commercial cultivars, and also of agronomic advancements in technology, efficiency, and production practices seen by the beet sugar industry over the last three decades.

Dynamic changes in the seed cultivars available to sugarbeet growers farming in the RRV have been one of the most important aspects of this upward trend in yield potential. Diseases such as rhizomania, aphanomyces, fusarium wilt, rhizoctonia root rot, and cercospora leaf spot are endemic to the RRV sugarbeet growing region. Although there are many fungicides currently labeled for management of these diseases, they are viewed as supplemental tools to aid the genetic defense package of the individual cultivars. Continued advancements in the field of plant breeding and molecular genomics have made it possible to use “customized” cultivars specifically adapted to growing areas within this region. Without this




Fig. 13–15. Average root yield combined with an increased level of quality equate to an increase in recoverable white sugar per hectare of 183 and 148 $\text{kg ha}^{-1} \text{ yr}^{-1}$, for commercially grown fields and the official variety trials over the last 24 yr. Data are from American Crystal Sugar Company, Minn-Dak Farmers Cooperative, and Southern Minnesota Beet Sugar Cooperative.

combination of genetic resistance in the crop, supplemented by chemical protections, diseases like cercospora leaf spot can devastate the region (Khan and Smith, 2005). Continued research and development of genetic resistance to pathogens, insects, and varied environmental conditions has contributed to—and continues to contribute to—the sugarbeet crop's success and progress in yield and quality.

In addition to improved genetics, one of the most notable developments contributing to increased sugar production is that of improved machinery. Over the past 25 yr, farm machines have gotten larger, more powerful, and very much more efficient. In the late 1980s, many growers utilized 8- to 12-row (4.5–6.7 m wide) equipment behind a 74,570 W (100-hp) row crop tractor (Shoptaugh, 1997). Today, sugarbeet growers in the RRV commonly use 24- to 36-row (13.4–20.1 m wide) equipment (a 300% increase in implement size) pulled with row crop tractors that often exceed 246,081 W (330-hp) and are able to guide the tractor from one end of the field to the other with sub-inch (<2.54 cm) accuracy via GPS guidance systems. As a result, the sugarbeet crop is planted earlier, quicker, and is raised far more efficiently than ever before. Since about 1980 target plant population has nearly doubled, which has played an important role in improved yield and quality.

Advancements in crop nutrition and fertilization (variable rate application is growing) also have contributed to improved crop yields. Scientific research has provided the individual grower and their respective cooperative with a much better understanding of the positive effects of fertilizers on sugarbeet quality and the detrimental effects of overuse. This newfound respect for optimized fertility has led to the widespread implementation and refinement of soil sampling. This annual monitoring has transitioned from crude random soil samples to an organized and methodical grid, zone, and topographical soil analysis taking advantage of the integration of GPS technology. The end result is a sugarbeet crop that advances more quickly physiologically and is healthier and more uniform.

Annual sugarbeet production surveys conducted by North Dakota State University from 1983 to 2007 consistently listed weed control as the top response by growers as their “worst production issue” (Carlson et al., 2007). Multiple species of pigweeds (*Amaranthus* spp.), lambsquarters (*Chenopodium album* L.), and Kochia [*Bassia scoparia* (L.) A.J. Scott] were among the weed species causing the largest problem, and they cost the cooperatives millions of dollars in lost revenue each year. Because of this, row crop cultivation and hand labor for weed control was a common practice throughout the growing season. Row crop cultivation was used by 99% of the respondents for each year from 1996 to 2007, while hand labor has steadily declined since the mid 1990s (62% of respondents used hand labor to thin the crop in 1996, 32% in 2002, and 20% in 2007) (Carlson et al., 2007). The addition of several new herbicide chemistries, and new application methodology (i.e., the Micro-Rate Program) (Rothe et al., 2004) were the main causes for the decline in hand weeding.

In 2008, the sugarbeet industry in the RRV transitioned into the use of glyphosate-resistant sugarbeet. Each cooperative held its grower membership to a maximum of 50% of their total allocated acreage during the first season and unlimited use for the second season (2009) and thereafter. The adoption of this new technology by the growers was a nearly 98% use rate by the second season. When comparing commercial yields, the average yield per hectare of sugarbeet

grown in this region from 1988 to 2007 was 43.48 Mg ha⁻¹. This average increased almost 9 Mg ha⁻¹ after the introduction of glyphosate-resistant sugarbeet (52.46 Mg ha⁻¹ average 2008–2012). The impact on quality is just as remarkable with the average kilograms of recoverable white sugar per hectare increasing 1308 kg (1988–2007: 6562 kg ha⁻¹; 2008–2012: 7870 kg ha⁻¹) per harvested hectare.

The Great Lakes Region (Michigan, Ohio, and Ontario, Canada)

Michigan has a long history of sugarbeet promotion and production. In 1881, the Michigan legislature offered a temporary bounty of two cents per pound for sweeteners derived from corn (*Zea mays* L.), sorghum [*Sorghum bicolor* (L.) Moench], and beet. However, it was not until 1890, when the Michigan Agricultural College (now Michigan State University) in East Lansing, MI imported 800 pounds of seed of four sugarbeet cultivars (Kleinwanzlebener), Vilmorin Imperial, Austrian Wohanka, and White Silesian) for distribution to 400 Michigan farmers, that interest was spurred in beet sugar production in Michigan. The Michigan Sugar Bounty Law of 1897 (Act no. 48) codified support for the nascent industry, and with help from the USDA in securing adequate quantities of seed from Europe, sugarbeet was produced in 64 counties with an average of 16% sugar at 83% purity, and interest in this new crop resulted in the distribution of 150 thousand copies of Bulletin 150 (*Sugar Beets in Michigan in 1897*) (Kedzie, 1901; Smith and Kedzie, 1987). Beet for sugar has been grown continuously in Michigan since that time.

Up until World War I, seed was imported from Germany, the majority of seed from the cultivar Kleinwanzlebener (Cox and Hill, 1924). During and after the war, faculty and students at Michigan Agricultural College (e.g., F.A. Spragg, H.B. Smith, and E.E. Down) began selecting within commercial cultivars and European germplasm. By 1923, no fewer than 29 cultivars were being imported from varied suppliers, and eight of these were produced in Michigan or Ontario for the 1921 season (Down, 1925). In 1923, the USDA Bureau of Plant Investigations (currently USDA-ARS) transferred work on sugarbeet from Blissfield, MI to East Lansing, and charged E.E. Down with providing improved lines of sugarbeet for the Michigan industry through a cooperative agreement with Michigan State College. From 1929, Kohls was the sugarbeet breeder for Michigan Agricultural College and was responsible for 'Michigan Hybrid 18,' the original source of genetic monogerm discovered by V.F. Savitsky (Savitsky, 1952). Seed was sent to V.F. Savitsky in about 1945 and two plants he discovered were monogerm (none were ever found in Michigan) (H.L. Kohls, personal notes, circa 1950).

By 1956, most multigerm breeding had ceased at the USDA-ARS station in East Lansing, MI, and efforts were directed towards converting the black-leg- and leaf spot-resistant multigerm populations to monogerm and O-type maintainers for the production of hybrid cultivars. The original source of monogerm and O-type germplasm for East Lansing is unclear; however, it likely involved materials received by H.L. Kohls from F.V. Owen (USDA, Salt Lake City, UT) in 1945. By 1948, these materials were incorporated into the first test hybrids, showing a sugar yield boost of >10% over commercial materials (Kohls, 1951), but they lacked disease resistance. Most or all of the Great Lakes monogerm sources were bred with assistance from USDA-ARS, Beltsville, MD materials (under the direction of G.E. Coe). Releases, 'EL 31' through 'EL 38,' were derived from introgressing monogerm and O-type into the '53AB1' ancestors with continual evaluation and

reselection. A combining ability test in 1962 compared 10 CMS parents. Ultimately, the combination of ‘SP 6822’ as pollinator (Coe and Hogaboam, 1971a) with selections for increased seed set from USDA Salt Lake City females ‘SLC 129’ and ‘SLC 133,’ which were released as ‘EL 44’ and ‘EL 45,’ respectively, were proved to be the most successful USDA hybrid for the Eastern growing regions, embodied in ‘US H20’ (Coe and Hogaboam, 1971b). The last USDA hybrid released for the Great Lakes growing region was ‘US H23,’ which used similar CMS material as ‘US H20,’ but used ‘EL 40’ as the pollinator for greater cercospora leaf spot resistance (Doney, 1995). Breeding for resistance to cercospora leaf spot, a perennial problem in the humid Great Lakes Region, has been recently summarized by Panella and McGrath (2010).

Since about 1985, private seed company breeders have contributed all commercial cultivars, while public breeders have focused on release of parental materials and germplasm enhancement. During the early years of commercial hybrids, performance was somewhat variable. Processors demanded higher sugar content, at the expense of good emergence and stand establishment, which is particularly important in the Great Lakes Region, and yields plummeted. Seed priming and greater attention to emergence genetics (McGrath et al., 2000) resulted in improving yields in the 21st century (Fig. 13–16). Plant populations, and hence yield, improved substantially. Since 2008, with the advent of glyphosate resistance and its attendant lack of injury to young beets from multi-application herbicide phytotoxicity and soil compaction, commercial yields have improved




Fig. 13–16. Research in seed priming and attention to the genetics of emergence has resulted in an increased percent emergence over the past 20 yr in Michigan. Increased emergence has resulted in better stands and more beets harvested, which have helped maintain a steady improvement in yield. Data are from the Michigan Sugar Company.

dramatically. Although there has been a steady increase in yield during the last 25 yr, in the years since 2007 when glyphosate-resistant cultivars began to appear in official variety trials, the rate of yield increase has accelerated dramatically (Fig. 13–17). Although acreage has declined by about 10% sugar production actually has increased in recent years, due to improved yields. Sucrose content of beets has remained relatively stable (Fig. 13–17).

The Great Lakes Region sugarbeet processors have consolidated over time. By the 1940s, there were more than 50 factories and 25 companies throughout Michigan, Ohio, and Ontario, Canada that were associated with the now defunct Farmers and Manufacturers Association. By 1980, two companies with five sugar factories remained, and sugarbeet was no longer grown in Ontario. Today, Michigan Sugar Company is a grower-owned cooperative with four factories and more than 1000 growers in Michigan and Ontario producing beet on more than 56,700 ha (140,000 ac) (Fig. 13–18).

What Does the Future Hold?

How will Climate Change Impact Sugarbeet Production?

Climate change will impact every aspect of agriculture, including sugarbeet cultivation and yield (Bindi and Olesen, 2011; Hatfield et al., 2011) by increasing levels of CO₂ and other greenhouse gases (GHG), increasing temperatures, and changing the patterns of precipitation (IPPC, 2008). Considerable uncertainty exists among




Fig. 13–17. Average yield over the past 25 yr has shown a steady increase in Michigan. Over that time the percent sucrose has remained stable. Data are from the Michigan Sugar Company.




Fig. 13–18. Area of sugarbeet harvested (left vertical axis) and average yield per harvested area (right vertical axis) are shown for the Great Lakes Region (Michigan and Ohio) from 1924 to 2012 based on statistics from the USDA-NASS.

climate models (based on different GHG emission scenarios), although the direct effects of these three main impacts are well supported by modeling (IPCC, 2008). Also predicted are increased variability in rainfall intensity and an increase in the number of sequential record high temperature days (Walthall et al., 2012; IPCC, 2008). Increased temperatures will mean faster snow melt and water that is available earlier in the season, at the expense of water currently stored throughout the season as snow or glacial ice (Bindi and Olesen, 2011; Hatfield et al., 2011; IPCC, 2008; Jaggard et al., 2010; Peltonen-Sainio et al., 2010; Walthall et al., 2012). More intense precipitation events interacting with different soil types may lead to increased erosion and more rapid runoff, affecting water storage in the profile and increasing nutrient contamination in streams and watersheds (Walthall et al., 2012). Flooding may affect both planting and harvest times (Hatfield et al., 2011). Heat stress will affect different crops differently and will be especially important at seed set in grain crops (IPCC, 2008). Increased temperatures may require relocating sugarbeet seed production areas, currently located in Western Oregon and similar environments where sugarbeet roots (stecklings) can be vernalized in situ with little chance of freezing damage (Kockelmann and Meyer, 2006).

Impacts of modifying complex agrobiological interactions are very difficult to predict (Newton et al., 2011). Very little is known about the impact of increased CO₂, temperature, and altered rainfall patterns on the growth and development of crop pests and diseases. Disease will still be a problem (Hatfield et al., 2011; Newton et al., 2011; Savary et al., 2011; Walthall et al., 2012), but the delicately

balanced management schemes that have been achieved in the past may change. For example, nematode pressure might increase with the increase in their number of reproductive cycles as a result of a longer growing season (Sticht et al., 2009). The changes also will affect the interactions of crop plants with weed species, native and invasive, and little research has been done to predict these impacts (Bradley et al., 2010; Walthall et al., 2012).

Experiments examining the effects of increased level of CO₂ on growth, photosynthesis, and dry matter accumulation in sugarbeet suggest increased productivity, but only with attention to different levels of inputs (e.g., Bunce, 1992; Demmers-Derks et al., 1998; Ford and Thorne, 1967; Ignatova et al., 2005; Romanova et al., 2002; Wolf, 1998; Ziska et al., 1995). Most results examined productivity in above-ambient CO₂ concentrations in controlled environments, where light intensity, in particular, may not correlate as well with field environments, or experiments in which plants have not been continuously exposed to high CO₂ throughout their life cycle. Also there is the general conclusion that the yield benefit gained by CO₂ enrichment is rapidly lost with nutrient stress, particularly with suboptimal nitrogen levels (e.g., Bunce, 1992; Demmers-Derks et al., 1998; Ford and Thorne, 1967; Ignatova et al., 2005; Romanova et al., 2002; Wolf, 1998; Ziska et al., 1995).

Sugarbeet was grown season long, without enclosure, under an enriched CO₂ concentration in a free air CO₂ enrichment (FACE) study in Braunschweig, Germany (Burkart et al., 2009; Manderscheid et al., 2010). White sugar yield increased about 13% under the higher concentration of CO₂, which was doubled (28%) when the sugarbeets were shaded, preventing a decrease in leaf area index, which often is seen under the elevated CO₂ concentrations. They concluded that the sugarbeet root was sink-limited under higher (than ambient) CO₂ concentrations, and therefore, the potential effect of elevated CO₂ on sugarbeet yield was minimized (Burkart et al., 2009; Manderscheid et al., 2010). Based on the research to date, elevated CO₂ concentrations will be beneficial for sugarbeet production and yields, but there still is considerable uncertainty as to the magnitude of the benefits.

An additional predicted consequence of the rise in the concentration of GHGs is the increase of ozone in the atmosphere (Walthall et al., 2012; IPCC, 2008). In a 2-yr study in an open topped chamber, sugarbeet exhibited a significant reduction of root yield of 6% and a slight reduction of sucrose content, resulting in a 9% reduction of sugar yield per hectare (De Temmerman et al., 2007). Another study reported that capacity of sugarbeet to recover from ozone exposure was dependent on the duration of exposure and the time in the plant's life cycle when it was exposed (Köllner and Krause, 2003). Although chlorophyll and sucrose content in the leaves showed some recovery, longer lasting effects were found in taproot glucose levels (Köllner and Krause, 2003). Increased levels of ozone may reduce yield potential of sugarbeet cultivars, although this biennial crop appeared less sensitive than annual crops such as potato (*Solanum tuberosum* L.) or wheat (*Triticum aestivum* L.) (De Temmerman et al., 2007).

Water and heat stress often are considered together when looking at abiotic challenges to sugarbeet production, and reviews are available (Morillo-Velarde and Ober, 2006; Ober and Rajabi, 2010; Stevanato, 2005). Heat and drought are of greatest concern in rainfed areas in the United States (i.e., the Red River Valley and Michigan), where access to irrigation to help alleviate water or heat stress is limited. Both the low and high GHG emission models examined by Walthall et

al. (2012) predicted that the rainfed areas will receive more precipitation, while western growing areas are predicted to have reduced precipitation, although both regions will have increased temperatures. Jaggard and coworkers (2007) attributed $0.025 \text{ t ha}^{-1} \text{ yr}^{-1}$ of a total $0.204 \text{ t ha}^{-1} \text{ yr}^{-1}$ increase in yield (1976–2004) to an earlier sowing date. Yield increases were attributed mostly to variation in the weather, and only about 30% of the yield increase over this time period was attributed to technological advances, including plant breeding (Jaggard et al., 2007). Similarly, Richter and coworkers (2006) concluded that growers could compensate for drought-related losses on sandy soils from climate change by planting earlier and harvesting later. The predicted warmer temperatures across the United States should allow for earlier planting dates, which could increase yields in rainfed and irrigated systems if moisture levels are moderate. In a simulation of sugarbeet production in northern and central Italy, the authors concluded that “sugarbeet production in northern and central Italy may not be greatly affected by future climate change, if global warming will be characterized by increased temperature and increased precipitation regimes” (Donatelli et al., 2002).

Each report and review lists a number of research areas that need to be pursued and cropping systems that need to be adjusted and integrated into current production systems to meet climate change challenges. Sugarbeet research always has been an area where private, state, and federal researchers have collaborated well. Current active research projects are focused on drought and heat tolerance in sugarbeet (Bloch et al., 2006; Ober and Rajabi, 2010; Ober et al., 2004; Pidgeon et al., 2006; Romano et al., 2013; Werker and Jaggard, 1998), but there is much more to be done. Walthall et al. (2012) stated, “Increases in temperature coupled with more variable precipitation will reduce productivity of crops, and these effects will outweigh the benefits of increasing carbon dioxide.” Areas of research often mentioned (Hatfield et al., 2011; IPCC, 2008, Walthall et al., 2012) and of particular interest to sugarbeet production include the following: We need to better understand the sensitivity of sugarbeet to key direct and indirect climate change effects and their interactions. Adaptive management strategies and climate risk tools need to be developed. More genetic improvement programs targeted to specific climate change factors such as tolerance to drought, heat, and ozone must be initiated and supported to increase yield potential. Accomplishing these goals will require multidisciplinary teams of scientists from industry, university, and federal research programs to work together. Sugarbeet has proven to be a very adaptable crop, and there likely is sufficient genetic variation within the *Beta* gene pool to deploy for future adaptations of the crop to changes in climate (Streibig et al., 2013).

Future Production and Uses of Sugarbeet

The 2008 Farm Bill (and previous farm bills) had provisions to stabilize the cost and production of sugar in the United States (information available online through USDA-ERS, 2013). The program is designed to operate to the greatest extent possible at no cost to the U.S. taxpayer. The approach is to stabilize prices using domestic marketing allotments and tariff-rate quotas to supply the U.S. need for sucrose. Sugar in excess of the domestic food consumption requirement is diverted into ethanol production. A current major challenge of this program is that unlimited, duty-free imports of sugar can now enter the United States from Mexico under the North American Free Trade Act.

Marketing allotments are important because they determine how much sugar is produced domestically. Each year these allotments are set based on USDA forecasts for consumer sugar requirements. There are two conditions that are met in the allotments. First, the overall allotment quantity must be at least 85% of estimated human consumption. Second, the domestic sugar prices must remain above the level that would cause forfeiture of government loans to the sugar processors (which are paid to the growers). The allotments are divided between sugarbeet and sugarcane producers, with sugarbeet producers receiving 54.35% of the allotment. This is divided among beet processors based on their production in crop years 1998 to 2000. Average per capita sugar consumption in the United States from 1988 to 2012 is constant, averaging 46.76 g d^{-1} (SD = 1.169) (1.65 oz.), and the population has grown an average of 0.91% per year since 2000 (The World Bank, 2013). As yield gains outpace population gains, the acreage needed to produce sugar will decline.

If sugarbeet growers are to maintain or expand acreage in the United States, it will be through new uses of beet. "Energy beet" as a feedstock for biofuel production already is being used in many European countries, and there are groups in North Dakota and California exploring its use in the United States (Panella, 2010). By dry weight, roughly 75% of the sugarbeet is sucrose (Harland et al., 2006), which easily is fermented into ethanol, but also could be fermented to a more energy dense methane, biobutanol, ETBE (ethyl tert-butyl ether), or biohydrogen (Antoni et al., 2007; Sanderson, 2006; von Felde, 2008; Zhang et al., 2007). The pulp or marc of sugarbeet, what is left after the sugar has been extracted, is comprised of more than 70% enzyme digestible complex saccharides (C-5/C-6) and only a small percentage of lignin (Eggleston et al., 2010; Harland et al., 2006; Kozak and Laufer, 2009). Many of the current cultivars used as energy beet are sugarbeet cultivars, but as the use of energy beets increases, breeding objectives will diverge from sugarbeet. One of the first objectives will be a suite of disease resistances suited to the climate in which the energy beet will be grown. If the entire beet and not just the pulp and molasses are being used as a feedstock, biomass might become a breeding objective (Panella and Kaffka, 2010). Breeding energy beets is just getting underway.

Very little of the sucrose produced or sugarbeet marc produced is used as value-added feedstock (Eggleston, 2008). Both hemicellulose and pectin are abundant, and there is value for all of these compounds as feedstock chemicals (Harland et al., 2006; Turley, 2008; van Beilen, 2008). Recently, the use of sugarbeet marc as an industrial source of carboxyl methyl cellulose and polysaccharides was reported (Fishman et al., 2010; Hotchkiss et al., 2010). There are a large number of uses for beet and its coproducts as the United States moves toward a more sustainable economy (McGrath and Townsend, 2014; Panella and Kaffka, 2010; Panella 2010), and there is the genetic variation within the beet genepool to be adapted through breeding to both changing climate and changing end uses (Streibig et al., 2013).

Acknowledgments

The authors would like to thank the U.S. sugarbeet grower-owned cooperatives, which provided much of the data used in this study. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the United States Department of Agriculture. USDA is an equal opportunity provider and employer.

References

- Antoni, D., V. Zverlov, and W. Schwarz. 2007. Biofuels from microbes. *Appl. Microbiol. Biotechnol.* 77:23–35. doi:10.1007/s00253-007-1163-x
- Asher, M.J.C., D.M. Chwarszczynska, and M. Leaman. 2002. The evaluation of rhizomania resistant sugar beet for the UK. *Ann. Appl. Biol.* 141:101–109. doi:10.1111/j.1744-7348.2002.tb00201.x
- Bangsund, D.A., N.M. Hodur, and F.L. Leistritz. 2012. Economic contribution of the sugarbeet industry in Minnesota and North Dakota. Rep. Agribusiness & Applied Economics Rep. 688. North Dakota State University, Department of Agribusiness and Applied Economics. <http://ideas.repec.org/p/ags/nddaae/121494.html> (accessed 18 Feb. 2014).
- Bennett, C.W. 1971. The curly top disease of sugarbeet and other plants. The American Phytopathological Society, St. Paul, MN.
- Bennett, C.W., and L.D. Leach. 1971. Diseases and their control. In: R.T. Johnson, J.T. Alexander, G.E. Rush, and G.R. Hawkes, editors, *Advances in sugarbeet production: Principles and practices*. The Iowa State Univ. Press, Ames, IA. p. 223–285.
- Bennett, C.W., C. Price, and J.S. McFarlane. 1957. Effects of Virus yellows on sugar beet with a consideration of some of the factors involved in changes produced by the disease. *Proc. Am. Soc. Sugar Beet Technol.* 9:479–494. doi:10.5274/jsbr.9.6.479
- Biancardi, E., L.G. Campbell, G.N. Skaracis, and M. De Biaggi. 2005. Genetics and breeding of sugar beet. *Science Publ.*, Enfield, NH.
- Biancardi, E., R.T. Lewellen, M. de Biaggi, A.W. Erichsen, and P. Stevanato. 2002. The origin of rhizomania resistance in sugar beet. *Euphytica* 127:383–397. doi:10.1023/A:1020310718166
- Biancardi, E., J.M. McGrath, L.W. Panella, R.T. Lewellen, and P. Stevanato. 2010. Sugar beet. In: J.E. Bradshaw, editor, *Root and tuber crops. Handbook of Plant Breeding Vol. 7*. Springer Science+Business Media, New York. p. 173–219.
- Biancardi, E., L.W. Panella, and R.T. Lewellen. 2012. *Beta maritima*, the origin of beets. Springer Science+Business Media, New York.
- Bindi, M., and J.E. Olesen. 2011. The responses of agriculture in Europe to climate change. *Reg. Environ. Change* 11:151–158. doi:10.1007/s10113-010-0173-x
- Bloch, D., C.M. Hoffmann, and B. Marlander. 2006. Impact of water supply on photosynthesis, water use and carbon isotope discrimination of sugar beet genotypes. *Eur. J. Agron.* 24:218–225. doi:10.1016/j.eja.2005.08.004
- Bockstähler, H.W., and O.E. Reece. 1948. Progress report on breeding of sugar beet in Minnesota for resistance to black root. *Proc. Am. Soc. Sugar Beet Technol.* 5:137–141.
- Bolton, M.D., L. Panella, L. Campbell, and M.F.R. Khan. 2010. Temperature, moisture, and fungicide effects in managing *Rhizoctonia* root and crown rot of sugar beet. *Phytopathology* 100:689–697. doi:10.1094/PHYTO-100-7-0689
- Bradley, B.A., D.M. Blumenthal, D.S. Wilcove, and L.H. Ziska. 2010. Predicting plant invasions in an era of global change. *Trends Ecol. Evol.* 25:310–318. doi:10.1016/j.tree.2009.12.003
- Bunce, J.A. 1992. Light, temperature and nutrients as factors in photosynthetic adjustment to an elevated concentration of carbon dioxide. *Physiol. Plant.* 86:173–179. doi:10.1111/j.1399-3054.1992.tb01327.x
- Buhre, C., G. Wagner, S. Kluth, C. Kluth, R. Apfelbeck, and M. Varrelmann. 2007. Resistance of sugarbeet varieties as basis for the integrated control of root and crown rot (*Rhizoctonia solani*). *Zuckerindustrie* 132:50–55.
- Burkart, S., R. Manderscheid, and H.J. Weigel. 2009. Canopy CO₂ exchange of sugar beet under different CO₂ concentrations and nitrogen supply: Results from a free-air CO₂ enrichment study. *Plant Biol.* 11:109–123. doi:10.1111/j.1438-8677.2009.00240.x
- Büttner, G., B. Märlander, and R. Manthey. 1995. Breeding for resistance to rhizomania in sugarbeet (*Beta vulgaris* L.). *Plant Breed.* 114:160–164. doi:10.1111/j.1439-0523.1995.tb00782.x
- Carlson, A.L., J.L. Luecke, M.F.R. Khan, and A.G. Dexter. 2007. Survey of weed control and production practices on sugarbeet in Minnesota and eastern North Dakota. Sugarbeet Research and Education Board Extension Rep. 38. p.40–64.
- Carsner, E. 1924. Studies on curly-top disease of the sugar beet. *J. Agric. Res.* 28:297–320.
- Carsner, E. 1926a. Resistance in sugar beets to curly top. *USDA Circular* 388:7.
- Carsner, E. 1926b. Seasonal and regional variations in curly-top of sugar beets. *Science* 63:213–214. doi:10.1126/science.63.1625.213
- Carsner, E. 1933. Curly-top resistance in sugar beets and test of the resistant variety U.S. No. 1. *USDA Technical Bull.* 360. USDA, U.S. Gov. Print. Office, Washington, DC.

- Coe, G.E., and G.J. Hogaboam. 1971a. Registration of sugarbeet parental line SP 6322-0. *Crop Sci.* 11:947. doi:10.2135/cropsci1971.0011183X001100060080x
- Coe, G.E., and G.J. Hogaboam. 1971b. Registration of US H20 sugarbeet. *Crop Sci.* 11:942. doi:10.2135/cropsci1971.0011183X001100060066x
- Coons, G.H. 1953. Breeding for resistance to disease. *Yearbook of Agriculture 1953*. USDA, U.S. Gov. Print. Office, Washington, DC. p.174–192.
- Coons, G.H., and J.E. Kotila. 1951. Virus yellows of sugar beets and tests for its occurrence in the United States. *Phytopathology* 41:559.
- Coons, G.H., F.V. Owen, and D. Stewart. 1955. Improvement of the sugar beet in the United States. *Adv. Agron.* 7:89–139. doi:10.1016/S0065-2113(08)60336-3
- Coons, G.H., D. Stewart, H.W. Bockstahler, G.W. Deming, J.O. Gaskill, G.J. Hogaboam, and C.L. Schneider. 1954. Evaluation tests in 1953 of U.S. 400 and related black root- and leaf spot-resistant varieties of the U.S. Department of Agriculture. *Proc. Am. Soc. Sugar Beet Technol.* 8:112–117.
- Coons, G.H., D. Stewart, and H.A. Elcock. 1931. Sugar-beet strains resistant to leaf spot and curly top. *Yearbook of Agriculture 1931*. USDA, U.S. Gov. Print. Office, Washington, DC. p. 493–496.
- Cox, J.F., and B.B. Hill. 1924. Sugar beet growing in Michigan. *Spec. Bull.* 106. Michigan Agric. Exp. Stn., East Lansing.
- Culbertson, J.O. 1942. Inheritance of factors influencing sucrose percentage in *Beta vulgaris*. *J. Agric. Res.* 64:153–172.
- Dahlberg, H.W., A.C. Maxon, and H.E. Brewbaker. 1940. Breeding for resistance to leaf spot and other characters. *Proc. Am. Soc. Sugar Beet Technol.* 2:169–180.
- De Temmerman, L., G. Legrand, and K. Vandermeiren. 2007. Effects of ozone on sugar beet grown in open-top chambers. *Eur. J. Agron.* 26:1–9. doi:10.1016/j.eja.2006.08.001
- Demmers-Derks, H., R.A.C. Mitchell, V.J. Mitchell, and D.W. Lawlor. 1998. Response of sugar beet (*Beta vulgaris* L.) yield and biochemical composition to elevated CO₂ and temperature at two nitrogen applications. *Plant Cell Environ.* 21:829–836. doi:10.1046/j.1365-3040.1998.00327.x
- Donatelli, M., F.N. Tubiello, U. Peruch, and C. Rosenzweig. 2002. Impacts of climate change and elevated CO₂ on sugar beet production in northern and central Italy. *Ital. J. Agron.* 6:133–142.
- Doney, D.L. 1995. USDA-ARS sugarbeet releases. *J. Sugar Beet Res.* 32:229–257. doi:10.5274/jsbr.32.4.227
- Doney, D.L., R.E. Wyse, and J.C. Theurer. 1981. The relationship between cell size, yield, and sucrose concentration of the sugarbeet root. *Can. J. Plant Sci.* 61:447–453. doi:10.4141/cjps81-060
- Down, E.E. 1925. Tests with sugar beet. *Circular Bull.* 66. Michigan Agric. Exp. Stn., East Lansing.
- Duffus, J.E. 1978. The impact of yellows control on California sugarbeets. *J. Am. Soc. Sugar Beet Technol.* 20:1–5. doi:10.5274/jsbr.20.1.1
- Duffus, J.E., E.D. Whitney, R.C. Larsen, H.Y. Liu, and R.T. Lewellen. 1984. First report in western hemisphere of rhizomania of sugar beet caused by *Beet necrotic yellow vein virus*. *Plant Dis.* 68:251.
- Eggleston, G. 2008. Sugar and related oligosaccharides. In: B. Fraser-Reid and T.K. Thiem, editors, *Glycoscience*. Springer Verlag, Heidelberg, Germany. p. 1164–1183.
- Eggleston, G., T. Tew, L. Panella, and T. Klasson. 2010. Ethanol from sugar crops. In: B.P. Singh, editor, *Industrial crops and uses*. CAB International, Chippenham, UK. p. 60–83.
- Elliott, M.C., and G.D. Weston. 1993. Biology and physiology of the sugar-beet plant. In: D.A. Cooke and R.K. Scott, editors, *The sugar beet crop: Science into practice*. Chapman and Hall, London. p. 37–65.
- Fischer, H.E. 1989. Origin of the 'Weisse Schlesische Rübe' (white Silesian beet) and resynthesis of sugar beet. *Euphytica* 41:75–80. doi:10.1007/BF00022414
- Fishman, M.L., H.K. Chau, P.H. Cooke, D.R. Coffin, and H. Hotchkiss, Jr. 2010. Sugar beet pulp: A sustainable source of carboxyl methyl cellulose (CMC) and other polysaccharides. American Chemical Society 239th ACS meeting, San Francisco, CA. 22 March 2010.
- Follett, R.F., W.R. Schmehl, and F.G.J. Viets. 1970. Seasonal leaf area, dry weight, and sucrose accumulation by sugarbeets. *J. Am. Soc. Sugar Beet Technol.* 16:235–252. doi:10.5274/jsbr.16.3.235
- Ford, M.A., and G.N. Thorne. 1967. Effect of CO₂ concentration on growth of sugar-beet, barley, kale, and maize. *Ann. Bot. (Lond.)* 31:629–644.

- Ford-Lloyd, B.V., A.L.S. Williams, and J.T. Williams. 1975. A revision of *Beta* section *Vulgares* (*Chenopodiaceae*), with new light on the origin of cultivated beets. *Bot. J. Linn. Soc.* 71:89–102. doi:10.1111/j.1095-8339.1975.tb02448.x
- Francis, S.A. 2006. Development of Sugar Beet. In: A.P. Draycott, editor, *Sugar beet*. Blackwell Publishing Ltd, Oxford, UK. p. 9–29.
- Francis, S.A., M. Redfearn, D.M. Chwarszczynska, M.J.C. Asher, and R.T. Lewellen. 1998. Use of molecular markers in breeding for disease resistance in sugar beet (*Beta vulgaris* L.). *Aspects App. Biol.* 52:279–285.
- Gayon, J., and D.T. Zallen. 1998. The role of the Vilmorin Company in the promotion and diffusion of the experimental science of heredity in France, 1840–1920. *J. Hist. Biol.* 31:241–262. doi:10.1023/A:1004335619901
- Gidner, S., B.-L. Lennefors, N.-O. Nilsson, J. Bensefelt, E. Johansson, U. Gyllenspetz, and T. Kraft. 2005. QTL mapping of BNYVV resistance from the WB41 source in sugar beet. *Genome* 48:279–285. doi:10.1139/g04-108
- Goldman, I.L., and J.P. Navazio. 2008. Table beet. In: J. Prohens and F. Nuez, editors, *Vegetables I. Asteraceae, Brassicaceae, Chenopodiaceae, and Cucurbitaceae*. Handbook of Plant Breeding. Springer, New York. p. 219–238.
- Grant, E.B. 1871. *Beet-root sugar and cultivation of the beet*. Lee, Shepard and Dillingham, New York.
- Grimmer, M., S. Trybush, S. Hanley, S. Francis, A. Karp, and M. Asher. 2007. An anchored linkage map for sugar beet based on AFLP, SNP and RAPD markers and QTL mapping of a new source of resistance to Beet necrotic yellow vein virus. *Theor. Appl. Genet.* 114:1151–1160. doi:10.1007/s00122-007-0507-3
- Grimmer, M.K., T. Kraft, S.A. Francis, and M.J.C. Asher. 2008. QTL mapping of BNYVV resistance from the WB258 source in sugar beet. *Plant Breed.* 127:650–652. doi:10.1111/j.1439-0523.2008.01539.x
- Hamilton, C. 2009. *Footprints in the sugar: A history of the Great Western Sugar Company*. Hamilton Bates, Caldwell, ID.
- Harland, J.I., C.K. Jones, and C. Hufford. 2006. Co-products. In: A.P. Draycott, editor, *Sugar beet*. Blackwell Publ., Oxford, UK. p. 443–463.
- Harris, F.S. 1919. *The sugar-beet in America*. Macmillan, New York.
- Hatfield, J.L., K.J. Boote, B.A. Kimball, L.H. Ziska, R.C. Izaurralde, D. Ort, A.M. Thomson, and D. Wolfe. 2011. Climate impacts on agriculture: Implications for crop production. *Agron. J.* 103:351–370.
- Hecker, R.J. 2004. Sugar Beet. In: W.F. Keim, editor, *A history of the Department of Agronomy/Soil & Crop Sciences*. Citizen Printing Company, Fort Collins, CO. p. 48–54.
- Henderson, R.W., and H.W. Bockstahler. 1946. Reaction of sugar beet strains to *Aphanomyces cochlioides*. *Proc. Am. Soc. Sugar Beet Technol.* 4:237–245.
- Hogaboam, G.J., R.C. Zielke, and C.L. Schneider. 1982. Registration of EL40 sugarbeet parental line. *Crop Sci.* 22:700.
- Hotchkiss, A., M. Fishman, and L. Liu. 2010. The role of sugar beet pulp polysaccharides in the sustainability of the sugar beet industry. In: *Sustainability of the sugar and sugar ethanol industries*. ACS Symposium Series. American Chemical Society, Washington, DC. p. 283–290.
- Ignatova, L.K., N.S. Novichkova, V.A. Mudrik, V.Y. Lyubimov, B.N. Ivanov, and A.K. Romanova. 2005. Growth, photosynthesis, and metabolism of sugar beet at an early stage of exposure to elevated CO₂. *Russ. J. Plant Physiol.* 52:158–164. doi:10.1007/s11183-005-0023-8
- IPCC. 2008. *Climate change 2007: Synthesis report*. Contribution of Working Groups I, II and III to the fourth assessment report of the Intergovernmental Panel on Climate Change. R.K. Pachauri and A. Reisinger, editors. IPCC, Geneva, Switzerland. http://www.ipcc.ch/publications_and_data/ar4/syr/en/contents.html (accessed 13 Feb. 2014).
- Jaggard, K.W., A. Qi, and M.A. Semenov. 2007. The impact of climate change on sugarbeet yield in the UK: 1976–2004. *J. Agric. Sci.* 145:367–375. doi:10.1017/S0021859607006922
- Jaggard, K.W., A. Qi, and E.S. Ober. 2010. Possible changes to arable crop yields by 2050. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 365:2835–2851. doi:10.1098/rstb.2010.0153
- Johnson, R.T., and G.E. Burt. 1990. *West Coast Beet Seed Company: 50th anniversary, 1940–1990*. West Coast Beet Seed Company, Salem, OR.
- Kedzie, R.C. 1901. The sugar beet industry in Michigan. In: *Proceedings of the Twenty-second Annual Meeting of the Society for the Promotion of Agricultural Science*, Denver, CO. p. 99–106.

- Khan, M.F.R., and L.J. Smith. 2005. Evaluating fungicides for controlling *Cercospora* leaf spot on sugar beet. *Crop Prot.* 24:79–86. doi:10.1016/j.cropro.2004.06.010
- Kiewnick, S., B.J. Jacobsen, A. Braun-Kiewnick, J.L.A. Eckhoff, and J.W. Bergman. 2001. Integrated control of *Rhizoctonia* crown and root rot of sugar beet with fungicides and antagonistic bacteria. *Plant Dis.* 85:718–722. doi:10.1094/PDIS.2001.85.7.718
- Kirk, W.W., P.S. Wharton, R.L. Schafer, P. Tumbalam, S. Poindexter, C. Guza, R. Fogg, T. Schlatter, J. Stewart, L. Hubbell, and D. Ruppel. 2008. Optimizing fungicide timing for the control of *Rhizoctonia* crown and root rot of sugar beet using soil temperature and plant growth stages. *Plant Dis.* 92:1091–1098. doi:10.1094/PDIS-92-7-1091
- Kockelmann, A., and U. Meyer. 2006. Seed Production and Quality. In: A.P. Draycott, editor, *Sugar beet*. Blackwell Publ., Oxford. p. 89–113.
- Kohls, H. 1951. New objectives in sugar beet breeding. Proc. 6th Regional Meeting of the ASSBT, Eastern US and Canada. American Society of Sugar Beet Technologists, Fort Collins, CO. p. 59–62.
- Köllner, B., and G.H.M. Krause. 2003. Effects of two different ozone exposure regimes on chlorophyll and sucrose content of leaves and yield parameters of sugar beet (*Beta vulgaris* L.) and rape (*Brassica napus* L.). *Water Air Soil Pollut.* 144:317–332. doi:10.1023/A:1022913116566
- Kozak, R., and C.S. Laufer. 2009. Addition of a thermostabilized pectin methylesterase significantly enhances the rate of saccharification of sugar beet pulp by the commercial pectinase preparation Pectinex® Ultra SPL. *J. Sugar Beet Res.* 46:71–72.
- Lange, W., W.A. Brandenburg, and T.S.M. de Bock. 1999. Taxonomy and cultonomy of beet (*Beta vulgaris* L.). *Bot. J. Linn. Soc.* 130:81–96. doi:10.1111/j.1095-8339.1999.tb00785.x
- Lewellen, R.T. 1992. Use of plant introductions to improve populations and hybrids of sugarbeet In: Use of plant introductions in cultivar development, Part 2. CSSA, Madison, WI. p. 117–135.
- Lewellen, R.T. 1997. Registration of 11 sugarbeet germplasm C79 lines with resistance to rhizomania. *Crop Sci.* 37:1026. doi:10.2135/cropsci1997.0011183X003700030087x
- Lewellen, R.T., H.Y. Liu, A. Gillen, and C. Strausbaugh. 2007. Performance of rhizomania resistant sugarbeet under normal and resistance-breaking strains of beet necrotic yellow vein virus. In: Proceedings from the 34th Biennial Meeting of American Society of Sugarbeet Technologists. 28 Feb.–3 Mar. 2007, Salt Lake City, UT. <http://assbt-proceedings.org/ASSBT2007Proceedings/PassbtVol34Agp171PerformanceofRhizomaniaResistantSugarbeetUnderNormalandResistanceBreakingStrainsofBeetNecroticYellowVeinVirus.pdf> (accessed 18 Feb. 2014).
- Lewellen, R.T., I.O. Skoyen, and A.W. Erichsen. 1987. Breeding sugarbeet for resistance to rhizomania: Evaluation of host-plant reactions and selections for and inheritance of resistance. In: Proc. Proceedings of the 50th Winter Congress of the IIRB. International Institute for Beet Research, Brussels, Belgium. p. 136–156.
- Lilleboe, D. 2012. Another world record sugarbeet crop in California. *The Sugarbeet Grower* 51(3):22.
- Liu, H.-Y., and R.T. Lewellen. 2007. Distribution and molecular characterization of resistance-breaking isolates of *Beet necrotic yellow vein virus* in the United States. *Plant Dis.* 91:847–851. doi:10.1094/PDIS-91-7-0847
- Liu, H.-Y., J.L. Sears, and R.T. Lewellen. 2005. Occurrence of resistance-breaking *Beet necrotic yellow vein virus* of sugar beet. *Plant Dis.* 89:464–468. doi:10.1094/PD-89-0464
- Manderscheid, R., A. Pacholski, and H.J. Weigel. 2010. Effect of free air carbon dioxide enrichment combined with two nitrogen levels on growth, yield and yield quality of sugar beet: Evidence for a sink limitation of beet growth under elevated CO₂. *Eur. J. Agron.* 32:228–239. doi:10.1016/j.eja.2009.12.002
- McGrann, G.R.D., M.K. Grimmer, E. Mutasa-Göttgens, and M. Stevens. 2009. Progress towards the understanding and control of sugar beet rhizomania disease. *Mol. Plant Pathol.* 10:129–141. doi:10.1111/j.1364-3703.2008.00514.x
- McGrath, J.M., C.A. Derrico, M. Morales, L.O. Copeland, and D.R. Christenson. 2000. Germination of sugar beet (*Beta vulgaris* L.) seed submerged in hydrogen peroxide and water as a means to discriminate cultivar and seedlot vigor. *Seed Sci. Technol.* 28:607–620.
- McGrath, J.M., C.A. Derrico, and Y. Yu. 1999. Genetic diversity in selected, historical USDA sugarbeet germplasm releases and *Beta vulgaris* ssp. *maritima*. *Theor. Appl. Genet.* 98:968–976. doi:10.1007/s001220051157
- McGrath, J.M., and K.K. Fugate. 2012. Analysis of sucrose from sugar beet. In: V.R. Preedy, editor, *Dietary sugars: Chemistry, analysis, function and effects*. Food and Nutritional Components in Focus No. 3. Royal Society of Chemistry Publishing, Cambridge, UK. p. 526–545.

- McGrath, J.M., L. Panella, and L. Frese. 2011. Beta. In: C. Kole, editor, *Wild crop relatives: Genomic and breeding resources, industrial crops*. Springer-Verlag, Berlin, Heidelberg. p. 1–28.
- McGrath, J.M., and B.J. Townsend. 2014. Sugar beet, energy beet, and industrial beet. In: V.M.V. Cruz and D.A. Dierig, editors, *Industrial crops breeding for bioenergy and bioproducts. Handbook of Plant Breeding*. Springer, New York. (in press.)
- Morillo-Velarde, R., and E.S. Ober. 2006. Water use and irrigation. In: A.P. Draycott, editor, *Sugar beet*. Blackwell Publishing Ltd, Oxford. p. 221–255.
- Murphy, A.M. 1942. Production of heavy curly-top exposures in sugarbeet breeding fields. *Proc. Am. Soc. Sugar Beet Technol.* 3:459–462.
- Murphy, A.M. 1946. Sugar beet and curly top history in Southern Idaho 1912–1945. *Proc. Am. Soc. Sugar Beet Technol.* 4:408–412.
- Newton, A., S. Johnson, and P. Gregory. 2011. Implications of climate change for diseases, crop yields and food security. *Euphytica* 179:3–18. doi:10.1007/s10681-011-0359-4
- Ober, E., and A. Rajabi. 2010. Abiotic stress in sugar beet. *Sugar Tech* 12:294–298. doi:10.1007/s12355-010-0035-3
- Ober, E.S., C.J.A. Clark, M. Le Bloa, A. Royal, K.W. Jaggard, and J.D. Pidgeon. 2004. Assessing the genetic resources to improve drought tolerance in sugar beet: Agronomic traits of diverse genotypes under droughted and irrigated conditions. *Field Crops Res.* 90:213–234. doi:10.1016/j.fcr.2004.03.004
- Owen, F.V. 1945. Cytoplasmically inherited male-sterility in sugar beets. *J. Agric. Res.* 71:423–440.
- Owen, F.V., F.A. Abegg, A.M. Murphy, B. Tolman, C. Price, F.G. Larmer, and E. Carsner. 1939. Curly-top-resistant sugar-beet varieties in 1938. Circular 513. USDA, Washington, DC.
- Panella, L. 2005. Curly top. In: E. Biancardi, L.G. Campbell, G.N. Skaracis, and M. de Biaggi, editors, *Genetics and breeding of sugar beet*. Science Publishers, Enfield, NH. p. 74–76.
- Panella, L. 2010. Sugar beet as an energy crop. *Sugar Tech.* 12:288–293. doi:10.1007/s12355-010-0041-5
- Panella, L., and S.R. Kaffka. 2010. Sugar beet (*Beta vulgaris* L) as a biofuel feedstock in the United States. In: G.Eggleston, editor, *Sustainability of the sugar and sugar ethanol industries*. ACS Symposium Series. American Chemical Society, Washington, DC. p. 163–175.
- Panella, L., and R.T. Lewellen. 2007. Broadening the genetic base of sugar beet: Introgression from wild relatives. *Euphytica* 154:383–400. doi:10.1007/s10681-006-9209-1
- Panella, L., and J.M. McGrath. 2010. The history of public breeding for resistance to *Cercospora* leaf spot in North America. In: R.T. Lartey et al., editors, *Cercospora leaf spot of sugar beet and related species*. APS Press, St. Paul, MN. p. 141–156.
- Pavli, O.I., P. Stevanato, E. Biancardi, and G.N. Skaracis. 2011. Achievements and prospects in breeding for rhizomania resistance in sugar beet. *Field Crops Res.* 122:165–172. doi:10.1016/j.fcr.2011.03.019
- Peltonen-Sainio, P., L. Jauhainen, M. Trnka, J.E. Olesen, P. Calanca, H. Eckersten, J. Eitzinger, A. Gobin, K.C. Kersebaum, J. Kozyra, S. Kumar, A.D. Marta, F. Micale, B. Schaap, B. Seguin, A.O. Skjelvåg, and S. Orlandini. 2010. Coincidence of variation in yield and climate in Europe. *Agric. Ecosyst. Environ.* 139:483–489. doi:10.1016/j.agee.2010.09.006
- Peterson, C.L., M.C. Hall, and D.J. Traveller. 1984. Sugarbeet storage experiments. ASAE Paper 84-4040. ASAE, Knoxville, TN.
- Pidgeon, J.D., E.S. Ober, A. Qi, C.J.A. Clark, A. Royal, and K.W. Jaggard. 2006. Using multi-environment sugar beet variety trials to screen for drought tolerance. *Field Crops Res.* 95:268–279. doi:10.1016/j.fcr.2005.04.010
- Richter, G.M., A. Qi, M.A. Semenov, and K.W. Jaggard. 2006. Modelling the variability of UK sugar beet yields under climate change and husbandry adaptations. *Soil Use Manage.* 22:39–47. doi:10.1111/j.1475-2743.2006.00018.x
- Romano, A., A. Sorgonà, A. Lupini, F. Araniti, P. Stevanato, G. Cacco, and M. Abenavoli. 2013. Morpho-physiological responses of sugar beet (*Beta vulgaris* L.) genotypes to drought stress. *Acta Physiol. Plant.* 35:853–865. doi:10.1007/s11738-012-1129-1
- Romanova, A.K., V.A. Mudrik, N.S. Novichkova, R.N. Demidova, and V.A. Polyakova. 2002. Physiological and biochemical characteristics of sugar beet plants grown at an increased carbon dioxide concentration and at various nitrate doses. *Russ. J. Plant Physiol.* 49:204–210. doi:10.1023/A:1014897422414
- Rothe, I., A.G. Dexter, and J.L. Luecke. 2004. Impact of injury from late season sequential glufosinate applications on sugarbeet yield. *J. Sugar Beet Res.* 41:119–135. doi:10.5274/jsbr.41.4.119

- Rush, C.M., H.-Y. Liu, R.T. Lewellen, and R. Acosta-Leal. 2006. The continuing saga of rhizomania of sugar beets in the United States. *Plant Dis.* 90:4–15. doi:10.1094/PD-90-0004
- Rush, C.M., and S.R. Winter. 1990. Influence of previous crops on *Rhizoctonia* root and crown rot of sugar beet. *Plant Dis.* 74:421–425. doi:10.1094/PD-74-0421
- Rush, C.M. 2003. Ecology and epidemiology of *Benyviruses* and plasmodiophorid vectors. *Annu. Rev. Phytopathol.* 41:567–592. doi:10.1146/annurev.phyto.41.052002.095705
- Sanderson, K. 2006. US biofuels: A field in ferment. *Nature* 444:673–676. doi:10.1038/444673a
- Savary, S., A. Nelson, A.H. Sparks, L. Willocquet, E. Duveiller, G. Mahuku, G. Forbes, K.A. Garrett, D. Hodson, J. Padgham, S. Pande, M. Sharma, J. Yuen, and A. Djurle. 2011. International agricultural research tackling the effects of global and climate changes on plant diseases in the developing world. *Plant Dis.* 95:1204–1216. doi:10.1094/PDIS-04-11-0316
- Savitsky, V.F. 1940. Genetics of sugar beets. USDA-National Agricultural Library Translation 5825. Translated from Russian, original source unavailable.
- Savitsky, V.F. 1950. Monogerm sugar beets in the United States. *Proc. Am. Soc. Sugar Beet Technol.* 6:156–159.
- Savitsky, V.F. 1952. Monogerm Sugar Beets in the United States. *Proc. Am. Soc. Sugar Beet Technol.* 7:156–159.
- Schneider, K., R. Schäfer-Pregl, D.C. Borchardt, and F. Salamini. 2002. Mapping QTLs for sucrose content, yield and quality in a sugar beet population fingerprinted by EST-related markers. *Theor. Appl. Genet.* 104:1107–1113. doi:10.1007/s00122-002-0890-8
- Scholten, O.E., T.S.M. de Bock, R. Klein-Lankhorst, and W. Lange. 1999. Inheritance of resistance to *Beet necrotic yellow vein virus* in *Beta vulgaris* conferred by a second gene for resistance. *Theor. Appl. Genet.* 99:740–746. doi:10.1007/s001220051292
- Scholten, O.E., R.M. Klein-Lankhorst, D.G. Esselink, T.S.M. DeBock, and W. Lange. 1997. Identification and mapping of random amplified polymorphic DNA (RAPD) markers linked to resistance against *Beet necrotic yellow vein virus* (BNYVV) in *Beta* accessions. *Theor. Appl. Genet.* 94:123–130. doi:10.1007/s001220050390
- Shoptaugh, T.L. 1997. Roots of success: History of the Red River Valley sugarbeet growers. Institute for Regional Studies, North Dakota State Univ., Fargo, ND.
- Skuderna, A.W. 1925. Sugar beet breeding in the Arkansas Valley of Colorado. *J. Am. Soc. Agron.* 17:631–634. doi:10.2134/agronj1925.00021962001700100009x
- Smith, C.D., and R.C. Kedzie. 1987. Sugar beets in Michigan in 1897. Bulletin 150 Agricultural College, Mich.: Michigan State Agricultural College Experiment Station. S75.E232 no.131–169 1896–1899.
- Smith, S., M. Cooper, J. Gogerty, C. Loeffler, D. Borcharding, and K. Wright. 2014. Maize. In: S. Smith, B. Diers, J. Specht, and B. Carver, editors, Yield gains in major U.S. field crops. CSSA Spec. Publ. 33. ASA, CSSA, and SSSA, Madison, WI. p. xxx–xxx. doi: 0.2135/cssaspecpub33.c6
- Souder, E., Jr. 1970. The century of the sugar beet. *The California sugar beet* 1970:34–36.
- Strausbaugh, C.A., and I.A. Eujayl. 2012. Influence of sugarbeet tillage systems on the *Rhizoctonia*-bacterial root rot complex. *J. Sugar Beet Res.* 49:57–76. doi:10.5274/jsbr.49.3.57
- Strausbaugh, C.A., I.A. Eujayl, and P. Foote. 2010. Seed treatments for the control of insects and diseases in sugarbeet. *J. Sugar Beet Res.* 47:105–125. doi:10.5274/jsbr.47.3.105
- Strausbaugh, C.A., I.A. Eujayl, and P. Foote. 2013a. Selection for resistance to the *Rhizoctonia*-bacterial root rot complex in sugar beet. *Plant Dis.* 97:93–100. doi:10.1094/PDIS-05-12-0511-RE
- Strausbaugh, C.A., I.A. Eujayl, L.W. Panella, and L.E. Hanson. 2011. Virulence, distribution and diversity of *Rhizoctonia solani* from sugar beet in Idaho and Oregon. *Can. J. Plant Pathol.* 33:210–226. doi:10.1080/07060661.2011.558523
- Strausbaugh, C.A., I.A. Eujayl, and L. Panella. 2013b. Interaction of sugar beet host resistance and *Rhizoctonia solani* AG-2-2 IIIB Strains. *Plant Dis.* doi:10.1094/PDIS-11-12-1078-RE.
- Strausbaugh, C.A., and A.M. Gillen. 2008. Bacteria and yeast associated with sugar beet root rot at harvest in the Intermountain West. *Plant Dis.* 92:357–363. doi:10.1094/PDIS-92-3-0357
- Strausbaugh, C.A., and A.M. Gillen. 2009. Sugar beet root rot at harvest in the US Intermountain West. *Can. J. Plant Pathol.* 31:232–240. doi:10.1080/07060660909507596
- Strausbaugh, C.A., A.M. Gillen, J.J. Gallian, S. Camp, and J.R. Stander. 2006. Influence of host resistance and insecticide seed treatments on curly top in sugar beets. *Plant Dis.* 90:1539–1544. doi:10.1094/PD-90-1539
- Strausbaugh, C.A., E.J. Wenninger, and I.A. Eujayl. 2012. Management of severe curly top in sugar beet with insecticides. *Plant Dis.* 96:1159–1164. doi:10.1094/PDIS-01-12-0106-RE

- Strausbaugh, C.A., W.M. Wintermantel, A.M. Gillen, and I.A. Eujayl. 2008. Curly top survey in the western United States. *Phytopathology* 98:1212–1217. doi:10.1094/PHYTO-98-11-1212
- Stenger, D.C., and C.L. McMahon. 1997. Genotypic diversity of Beet curly top virus populations in the western United States. *Phytopathology* 87:737–744. doi:10.1094/PHYTO.1997.87.7.737
- Stevanato, P. 2005. Resistance to abiotic stresses. In: E. Biancardi, L.G. Campbell, G.N. Skaracis, and M. de Biaggi, editors, *Genetics and breeding of sugar beet*. Science Publishers, Enfield, NH. p. 116–119.
- Sticht, C., S. Schrader, A. Giesemann, and H.J. Weigel. 2009. Sensitivity of nematode feeding types in arable soil to free air CO₂ enrichment (FACE) is crop specific. *Pedobiologia* 52:337–349. doi:10.1016/j.pedobi.2008.12.001
- Stump, W.L., G.D. Franc, R.M. Harveson, and R.G. Wilson. 2004. Strobilurin fungicide timing for *Rhizoctonia* root and crown rot suppression in sugarbeet. *J. Sugar Beet Res.* 41:17–38. doi:10.5274/jsbr.41.1.17
- Streibig, J.C., C. Ritz, C.B. Pipper, F. Yndgaard, and J.N. Tomsen. 2013. Sugar beet, bioethanol, and climate change. IOP Conf. Series: Earth and Environmental Science. *Climate Change: Global Risks, Challenges and Decisions* 6. doi:10.1088/1755–1307/6/24/242040.
- The World Bank. 2013. The World Bank: Data: Population growth (annual %). <http://data.worldbank.org/indicator/SP.POP.GROW/countries/1W?display=default> 8-21-2013 (accessed 18 Feb. 2014).
- Theis, T. 1971. A food resource. In: R.T. Johnson et al., editors, *Advances in sugarbeet production: Principles and practices*. The Iowa State Univ. Press, Ames, IA. p. 3–18.
- Turley, D.B. 2008. The chemical value of biomass. In: J.H. Clark and F.E.I. Deswarte, editors, *Introduction to chemicals from biomass*. Wiley, Sussex, UK. p. 21–46.
- United States Beet Sugar Association. 1936. *The silver wedge: The sugar beet in the United States*. United States Beet Sugar Association, Washington, DC.
- USDA-ERS. 2013. Sugar & sweeteners: Policy. <http://www.ers.usda.gov/topics/crops/sugar-sweeteners/policy.aspx> (accessed 23 Jan. 2013).
- USDA-NASS. National Agricultural Statistics Service/USDA. http://quickstats.nass.usda.gov/results/2F65366F-16C2-384B-9B53-ABFE2FE826F1?pivot=short_desc (accessed 9 Apr. 2013).
- van Beilen, J.B. 2008. Transgenic plant factories for the production of biopolymers and platform chemicals. *Biofuels Bioprod. Bioref.* 2:215–228. doi:10.1002/bbb.64
- Vilmorin, P.-P.-A.L. 1859. Notice sur l'amélioration de la Carotte sauvage. In: L. Vilmorin, editor, *Notices sur l'amélioration des plantes par le semis et considérations sur l'hérédité dans les végétaux*. 1st ed. Librairie Agricole, Paris.
- von Felde, A. 2008. Trends and developments in energy plant breeding—Special features of sugarbeet. *Zuckerindustrie* 133:342–345.
- Walthall, C.L., J. Hatfield, P. Backlund, L. Lengnick, E. Marshall, M. Walsh, S. Adkins, M. Aillery, E.A. Ainsworth, C. Ammann, C.J. Anderson, I. Bartomeus, L.H. Baumgard, F. Booker, B. Bradley, D.M. Blumenthal, J. Bunce, K. Burkey, S.M. Dabney, J.A. Delgado, J. Dukes, A. Funk, K. Garrett, M. Glenn, D.A. Grantz, D. Goodrich, S. Hu, R.C. Izaurralde, R.A.C. Jones, S.-H. Kim, A.D.B. Leaky, K. Lewers, T.L. Mader, A. McClung, J. Morgan, D.J. Muth, M. Nearing, D.M. Oosterhuis, D. Ort, C. Parmesan, W.T. Pettigrew, W. Polley, R. Rader, C. Rice, M. Rivington, E. Rosskopf, W.A. Salas, L.E. Sollenberger, R. Srygley, C. Stöckle, E.S. Takle, D. Timlin, J.W. White, R. Winfree, L. Wright-Morton, and L.H. Ziska. 2012. Climate change and agriculture in the United States: Effects and adaptation. *USDA Tech. Bull.* 1935. Washington, DC. [http://www.usda.gov/oce/climate_change/effects_2012/CC%20and%20Agriculture%20Report%20\(02-04-2013\)b.pdf](http://www.usda.gov/oce/climate_change/effects_2012/CC%20and%20Agriculture%20Report%20(02-04-2013)b.pdf).
- Ware, L.S. 1880. *The sugar beet: Including a history of the beet sugar industry in Europe*. Henry, Carey, Baird, and Co., Philadelphia, PA.
- Werker, A.R., and K.W. Jaggard. 1998. Dependence of sugar beet yield on light interception and evapotranspiration. *Agric. For. Meteorol.* 89:229–240. doi:10.1016/S0168-1923(97)00081-6
- Whitney, E.D. 1989. Identification, distribution, and testing for resistance to rhizomania in *Beta maritima*. *Plant Dis.* 73:287–290. doi:10.1094/PD-73-0287
- Windels, C.E., and J.R. Brantner. 2005. Early-season application of azoxystrobin to sugarbeet for control of *Rhizoctonia solani* AG 4 and AG 2-2. *J. Sugar Beet Res.* 42:1–16. doi:10.5274/jsbr.42.1.1
- Wisler, G.C., and J.E. Duffus. 2000. A century of plant virus management in the Salinas Valley of California, 'East of Eden'. *Virus Res.* 71:161–169. doi:10.1016/S0168-1702(00)00196-9

- Wolf, J. 1998. Effects of nutrient (NPK) supply on sugar beet response to elevated atmospheric CO₂. *NJAS-Wageningen J. Life Sci.* 48:157–170.
- Zhang, Y.-H.P., B.R. Evans, J.R. Mielenz, R.C. Hopkins, and M.W.W. Adams. 2007. High-yield hydrogen production from starch and water by a synthetic enzymatic pathway. *PLoS ONE* 2:e456. doi:10.1371/journal.pone.0000456
- Zhao, B., I.J. Mackay, P.D.S. Caligari, and R. Mead. 1997. The efficiency of between and within full-sib family selection in a recurrent selection programme in sugar beet (*Beta vulgaris* L). *Euphytica* 95:355–359. doi:10.1023/A:1003026800450
- Zimmermann, B., and J. Zeddies. 2000. Evolution of sugar beet breeding and its contribution to value added. Univ. of Hohenheim, Dep. Agricultural Economics, Hohenheim, Germany.
- Ziska, L.H., R.C. Sicher, and D.F. Kremer. 1995. Reversibility of photosynthetic acclimation of swiss chard and sugarbeet grown at elevated concentrations of CO₂. *Physiol. Plant.* 95:355–364. doi:10.1111/j.1399-3054.1995.tb00849.x

