This does not constitute a formal recommendation. When using herbicides always read the label, and when in doubt consult your farm advisor or county agent.

This is an excerpt from the book Weed Control in Natural Areas in the Western United States and is available wholesale through the UC Weed Research & Information Center (wric.ucdavis.edu) or retail through the Western Society of Weed Science (wsweedscience.org) or the California Invasive Species Council (cal-ipc.org).

Plantago lanceolata

Buckhorn plantain

Family: Plantaginaceae

NON-CHEMICAL CONTROL

Grazing	P	can reduce growth only slightly
Prescribed burning	P	
Mowing and cutting	P	may give some reduction of growth, but low growing habit will reduce control
Tillage	G	
Grubbing, digging or hand pulling	G	difficult but much more effective than cutting

CHEMICAL CONTROL

The following specific use information is based on published papers and reports by researchers and land managers. Other trade names may be available, and other compounds also are labeled for this weed. Directions for use may vary between brands; see label before use.

2,4-D		
Aminocyclopyrachlor + chlorsulfuron		
Aminopyralid		
Chlorsulfuron	NIA	
Clopyralid	P	
Dicamba	E	
Glyphosate	G	
Hexazinone	G	

Imazapic	G
Imazapyr	NIA
Metsulfuron	E
Paraquat	P
Picloram	E
Rimsulfuron	G
Sulfometuron	E
Sulfosulfuron	NIA
Triclopyr	G

= Excellent control, generally better than 95%

G = Good control, 80-95%

= Fair control, 50-80%

= Poor control, below 50%

Control includes effects within the season of treatment.

= Likely based on results of observations of related species

FLW = flowering

NIA = No information available

= Fall = Spring = Summer

Control is followed by best timing, if known, when efficacy is \mathbf{E} or \mathbf{G} .

RECOMMENDED CITATION: DiTomaso, J.M., G.B. Kyser et al. 2013. Weed Control in Natural Areas in the Western United States. Weed Research and Information Center, University of California. 544 pp.

> I of I 2013