

09/02/22 through 09/08/22

Crops (Leafout Date)	#148 Merced				#39 Parlier				#258 Lemon Cove			
	9/2 - 9/8 Water Use	Accum'd Seasonal Water Use	9/9 - 9/15 Estimated ETc		9/2 - 9/8 Water Use	Accum'd Seasonal Water Use	9/9 - 9/15 Estimated ETc		9/2 - 9/8 Water Use	Accum'd Seasonal Water Use	9/9 - 9/15 Estimated ETc	
Almonds (3/1) *	1.73	41.47	1.47		1.79	42.57	1.35		1.72	39.60	1.26	
Pistachio (4/8) * **	1.73	36.47	1.38		1.77	37.63	1.26		1.69	34.75	1.17	
Citrus (2/1)	1.16	31.50	1.09		1.21	32.37	0.99		1.16	30.32	0.90	
Raisin Grapes (3/15) (11 ft. row spacing)	1.15	24.08	1.01		1.17	24.86	0.91		1.12	22.92	0.83	
Winegrapes (3/15) (10 ft. spacing on California Sprawl Trellis) ***	1.31	26.48	1.13		1.35	27.28	1.03		1.29	25.16	0.95	
Walnuts (4/8)	1.54	37.12	1.33		1.59	38.29	1.21		1.52	35.48	1.14	
Stone Fruit (3/10)	1.77	34.78	1.54		1.82	35.81	1.42		1.73	33.00	1.33	
Past 7 days precipitation (inches)	0.00				0.00				0.00			
Accumulated precipitation (inches) (1/1/2022)	1.98				1.94				1.82			

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹

Crops	#148 Merced				#39 Parlier				#258 Lemon Cove			
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	2.7	2.3	2.0	1.8	2.8	2.4	2.1	1.9	2.6	2.3	2.0	1.8
Pistachio (4/8)	2.7	2.3	2.0	1.8	2.7	2.4	2.1	1.9	2.6	2.3	2.0	1.8
Citrus (2/1)	1.8	1.5	1.4	1.2	1.9	1.6	1.4	1.3	1.8	1.5	1.4	1.2
Raisin Grapes (3/15) (11 ft. row spacing)***	Assume all grape			1.2	Assume all grape			1.2	Assume all grape			1.2
Winegrapes (3/15) (10 ft. spacing on California Sprawl Trellis) ***	irrigation type is drip			1.4	irrigation type is drip			1.4	irrigation type is drip			1.4
Walnuts (4/8)	2.4	2.1	1.8	1.6	2.4	2.1	1.9	1.7	2.3	2.0	1.8	1.6
Stone Fruit (3/10)	2.7	2.4	2.1	1.9	2.8	2.4	2.1	1.9	2.7	2.3	2.0	1.8

¹ The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

Crops	#148 Merced				#39 Parlier				#258 Lemon Cove			
Almonds 115 Trees/A	638	543	472	425	661	567	496	449	614	543	472	425
Pistachio 106 Trees/A	673	573	498	448	673	598	523	473	648	573	498	448
Citrus 110 Trees/A	444	370	346	296	469	395	346	321	444	370	346	296
Raisin Grapes 566 Vines/A	Assume all grape			58	Assume all grape			58	Assume all grape			58
Winegrapes 622 Vines/A	irrigation type is drip			61	irrigation type is drip			61	irrigation type is drip			61
Walnuts 76 Trees/A	857	750	643	572	857	750	679	607	822	715	643	572
Stonefruit 172 Trees/A	426	379	332	300	442	379	332	300	426	363	316	284

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

Crops	#124 Panoche				#2 Five Points				#15 Stratford			
System Efficiency >>>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	2.9	2.5	2.2	2.0	2.8	2.5	2.2	1.9	3.2	2.8	2.5	2.2
Pistachio (4/8)	2.9	2.5	2.2	2.0	2.8	2.5	2.2	1.9	3.2	2.8	2.5	2.2
Citrus (2/1)	2.0	1.8	1.6	1.4	2.0	1.7	1.5	1.4	2.3	2.0	1.7	1.5
Raisin Grapes (3/15) (11 ft. row spacing)	Assume all grape			1.3	Assume all grape			1.3	Assume all grape			1.5
Winegrapes (3/15) (10 ft. spacing on California Sprawl Trellis)	irrigation type is drip			1.5	irrigation type is drip			1.5	irrigation type is drip			1.7
Walnuts (4/8)	2.6	2.3	2.0	1.8	2.6	2.2	2.0	1.7	2.9	2.5	2.2	2.0
Stone Fruit (3/10)	3.0	2.6	2.3	2.1	2.9	2.5	2.2	2.0	3.3	2.9	2.5	2.3

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.