

TABLE 5. Percentage of mold development on prunes harvested by hand from Tulare and incubated at 4 C.

Treatment	Percentage of mold ^a after incubation days:				
	8	11	15	20	30
Unwashed	8.62 Ab	24.03 Ab	40.10 Ab	48.27 ABC	64.35 AC
Washed	6.71 A	25.70 A	38.30 A	49.31 A	61.04 A
Chlorine	1.15 B	10.21 B	17.30 B	35.72 B	47.54 AB
K-sorbate	0.00 B	8.19 B	4.11 C	57.12 A	67.21 A
CGA	0.00 B	1.36 C	2.07 C	9.17 C	32.49 B

^aMean of 3 replications.

^bSignificant differences at 1% level.

^cSignificant differences at 5% level.

TABLE 6. Percentage of mold development on prunes harvested by hand from Tulare and incubated at 20 and 28 C.

Treatment	Percentage of mold ^a at:			
	20 C		28 C	
	4 days	5 days	4 days	5 days
Unwashed	38.86 Ab	67.91 Ab	35.20 A	70.83 A
Washed	36.19 A	71.68 A	29.88 A	74.78 A
K-sorbate	22.62 AB	68.30 A	23.31 A	55.44 A
Chlorine	13.29 B	35.33 B	12.51 A	49.55 A
CGA	4.97 B	39.10 B	12.56 A	59.28 A

^aMean of 3 replicates.

^bMeans differ significantly at 1% level.