

Inadequate drainage was the cause of a soil salinity problem on 25,000 acres of Tulelake land during a five-year study of water tables and soil salinity.

The predominant crop in the Tulelake area is barley, a crop that can stand higher soil salt concentration than most other crops. However, when the salinity level exceeds four millimhos—unit of measurement of electrical conductivity of saline—a reduction in barley yield begins. At eight millimhos the yield is reduced about 20% and at 16 millimhos there is a 50% reduction in yield.

A field examination of the salinity problem in Tulelake was initiated in 1955 and during the growing season water table heights were measured by continuous recorders and observation wells. Soil samples were taken adjacent to the recorders three times during the growing season—at the start, the middle, and the end—and analyzed for total salts. Because of the limited number of sampling sites included in the study, the salinity readings are merely indicative of the general situation. Undoubtedly there are many salty areas not sampled, and some of the salty land may include areas of low salt.

Soil samples from several locations at the end of the growing season were examined to determine the vertical distribution of salt in the soil profile. One representative analysis showed a high concentration of salt in the surface soil layers. The top inch of soil contained about twice as much salt as the second half foot.

Soil samples taken at the end of the growing season clearly indicated that the salinity problems in the land studied were the result of the high water table.

JAMES N. LUTHIN AND KENNETH BAGHOTT

# Soil Salinity

## on Tulelake lands

Water moves upward from the water table level by capillary action to the soil surface and evaporates, leaving the salts in the soil. However, the water table must be close to the soil surface for long periods of time for the upward movement and evaporation of the water to take place. Poor soil drainage and inadequate

rains depends on the total seasonal rainfall, the amount of each storm, the level of the water table during the rainy season, and on the adequacy of field drains for rapid removal of water from the soil.

When the water table is close to the soil surface, the winter rains are not effective for leaching. However, when the water table is kept 4' or 5' below the soil surface by a system of drains, winter rains will wash out an appreciable amount of salt.

To keep the water table sufficiently below the soil surface, some sort of drainage—tile lines or open ditches—must be provided at a depth of at least 4' and spaced close enough together to lower the water table rapidly.

Water must move through and out of the soil to get rid of the salts, so ponding water on the soil without adequate drainage accomplishes little in the way of leaching.

Flooding land lacking drainage prior to the growing season, as has been the practice for many years in the Southwest Sump of the Tulelake lands, has resulted in a continued increase in the soil salinity problem.

*James N. Luthin is Associate Professor of Irrigation, University of California, Davis.*

*Kenneth Baghott is Farm Advisor, Modoc-Siskiyou Counties, University of California.*

Salt Content in Surface 6"

Depth	Salt Millimhos
0" - 1/2"	14.80
1/2" - 1 1/2"	14.80
1 1/2" - 3"	9.12
3" - 6"	8.48

Salt Content at Same Location with 1/2 foot Sampling

0" - 1/2"	10.22
1/2" - 1"	6.90
1" - 1 1/2"	6.23
1 1/2" - 2"	6.01
2" - 2 1/2"	5.32
2 1/2" - 3"	4.64

control of irrigation water contribute to the salinity problem.

The following spring more soil samples were taken to determine the effect of the winter rains in soil leaching. There was considerable leaching of the salt and the average salinity was below the levels found at the start of the field examination. The amount of leaching by winter

## TENSIOMETERS

Continued from preceding page

chard would reduce the number of trees in an irrigation run and give more flexibility in the irrigation practices of the two areas. A three-week irrigation schedule with a reduced amount of water would prevent excessive saturated soil conditions and benefit the orchard.

Another orchard on Ramona sandy loam appears to have a very active root system at both the 12" and 24" soil depths. A slowly permeable subsoil restricts the downward movement of water and forces it to move laterally until the whole soil area around the tree is saturated. A three-week irrigation schedule

with not more than 24 hours of water applied at one time would be a better irrigation program than a monthly program for a three-day period.

A third orchard—on a Hanford sandy loam—has a deep root system and root activity causes the soil suctions to increase more rapidly at the 24" and 36" soil depths than at the 12" depth. Apparently some restriction in the soil between the 12" and the 24" depths prevents the soil suction values at the deeper depths from reaching saturation. The recovery from saturated conditions at the 12" depth makes a favorable condition for feeder roots. Tensiometer readings indicate a good irrigation program in the orchard.

Citrus trees are conditioned by cultural practices and a shallow-rooted orchard might not stand a sudden change to longer intervals between irrigations. Citrus trees under sprinkler irrigation tend to develop root systems that are shallower than those of furrow-irrigated trees. Also, sprinkler irrigated citrus trees wilt at much lower suction values, for comparable depths, than trees under furrow irrigation.

*L. H. Stolzy is Assistant Irrigation Engineer, University of California, Riverside.*

*A. W. Marsh is Extension Irrigation and Soil Specialist, University of California, Riverside.*

*R. E. Puffer is Farm Advisor, San Bernardino County, University of California.*

*D. C. Baier is Farm Advisor, Orange County, University of California.*