

Annual Forage Plants

aids plant use of available soil moisture

Cyrus M. McKell, Jack Major, and Eugene R. Perrier

Seed germination and early plant growth of most annual forage species on approximately 18 million acres of California's rangeland take place soon after the first fall rains, but the greatest growth is in late winter and spring when temperatures and soil moisture supplies are favorable.

Fertilization of annual rangelands is an important range improvement technique. Fertilization increases plant growth and therefore should stimulate plant use of soil moisture. The amount of available moisture and its distribution throughout the growing season are critical factors in plant growth on annual rangelands and may be significantly altered by fertilizer application.

The pattern of soil moisture depletion under moderately high rates of nitrogen and phosphorus fertilization was studied in a completely randomized block experiment in Solano County that included four fertilizer treatments replicated four times. Treatments applied before the first fall rain consisted of 150 pounds of nitrogen per acre, 200 pounds of phosphorus per

acre, a combination of 150 pounds of nitrogen and 200 pounds of phosphorus per acre, and an untreated check plot. Ammonium nitrate and treble superphosphate were the fertilizer materials applied.

The test area is on the west slope of the Sacramento Valley near the Coast Range and has an average annual rainfall of 18" to 20" from September

the yield on non-fertilized plots or on plots receiving only phosphorus. The combination of nitrogen and phosphorus produced a greater forage yield than did any other treatment. The nitrogen treatment produced more than twice the yield of plots given no treatment, and on the nitrogen-phosphorus-treated plots the yield was about four times as much as on the non-fertilized check plots.

Yield and Moisture Use Efficiency of Annual Range Following Various Fertilizer Treatments

Item	Treatment			
	No fertilizer	Phosphorus 200 lbs./a	Nitrogen 150 lbs./a	Nitrogen 150 lbs./a + phosphorus 200 lbs./a
Inches of rainfall, (October-June)	17.01	17.01	17.01	17.01
Forage yield in lbs./a	1,440	1,704	3,169	5,929
Differences			1,465*	2,760*
Lbs. forage/inch of rainfall	84.7	100.2	186.3	348.6
Differences		15.5	86.1*	162.3*

* Indicates significance at the 1% level.

through May. The topography is slightly undulating and soils are derived from alluvium and old valley terraces. Soil of the study site is a mosaic of Olcott fine sandy loam and Clear Lake clay adobe. Vegetation consisted of resident annual grasses and forbs.

The experimental area was fenced, and plots 10' x 11' were centrally equipped with Bouyoucos resin-impregnated gypsum blocks so that soil moisture depletion at the 6", 12", 20", and 36" depths could be measured. In April, 20"-deep columns 6" x 12" were excavated from the variously treated plots, and plant roots were removed by washing.

Forage yield was measured at the peak of growth by clipping, at the 1" height, one square foot of area from each of the plots.

Within two weeks after the first rain—2.5"—a plant growth response was noted, and the growth continued during the cool winter months at a greater rate than that on non-fertilized or phosphorus-fertilized plots.


Forage yield on plots fertilized with nitrogen was significantly greater than

A comparison of forage yield in relation to soil moisture used for each treatment indicates that fertilization enables more efficient plant use of rainfall. In range areas, where rainfall is usually the only source of water for plant growth, fertilization produces more forage with a given amount of moisture.

Root excavations on April 15 indicated that a majority of the plant roots in the non-fertilized plots were located within the upper 12" of soil. Excavation for a soil profile pit to the 6' depth at the end of the summer in a non-fertilized area revealed that roots were present in abundance at the surface and gradually diminished toward the bottom of the pit. In the nitrogen-phosphorus plots roots were uniformly abundant to the 20" depth. A large number of roots in this plot were severed when the soil columns 20" deep were removed. No deeper excavations were made in the fertilized plots. The over-all reactions of the vegetation to fertilization were illustrated by the course of soil moisture depletion during the growing season.

Continued on next page

Calibration curve for Bouyoucos resin-impregnated gypsum blocks.



FORAGE PLANTS

Continued from preceding page

All plots were at field capacity on March 12 even though the nitrogen and nitrogen-phosphorus plots were supporting luxuriant stands of annual grasses and forbs. Therefore, the increased plant growth in winter did not appreciably drain the moisture reserve in the soil, and moisture was equally available on all plots for the rapid growth that occurred as soon as spring temperatures were favorable.

No summer-growing annual weeds, such as turkey mullein, spikeweed, and tarweed, were noted on the nitrogen-phosphorus plots. Observations indicated that a greater number of plants matured earlier on the nitrogen and nitrogen-phosphorus plots than on the unfertilized plots.

The graphs below give the moisture-extraction pattern for each of the treatments and the rainfall, in inches, as a function of time. The extraction patterns are expressed in terms of suction, in bars,

and as the suction force increases the availability of water to the plants is reduced. Plants on the nitrogen and nitrogen-phosphorus plots were subjected to higher suction forces in general than were those on the phosphorus or the non-fertilized plots. These suction forces ranged from 15 bars to 25 bars and were associated with extent and time of maturity.


Plants that received an application of nitrogen reduced the soil moisture below the field capacity—0.3 bar—earlier in the season, near the first of April, than did the phosphorus or non-fertilized plots. Also, the rate of soil-moisture use was higher in these plots, particularly at the lower depths. It appears that, in the nitrogen-phosphorus plots, root development was greater to the 36" depth than in plots with the other treatments as indicated by the rate and time of moisture extraction. Removal of available soil moisture from the 36" depth began in the nitrogen-phosphorus plots on March 28; in the nitrogen plots, on April 8; in the phosphorus plots, on April 12; and

in the non-fertilized check plots, on April 26.

A rainfall of 1.90" and an accompanying low transpiration period during the last half of April were major causes of a reduced rate of moisture depletion. Toward the end of April, moisture use at the shallow depth in all plots increased because of warmer weather, but on the nitrogen-phosphorus and the nitrogen plots moisture use began increasing at all depths.

At the 6", 12", and 20" depths the nitrogen-phosphorus and nitrogen plots reached the wilting point—15 bar suction—by May 10. However, on nitrogen plots the plants continued to use moisture at all depths but, in general, were later in maturity than those on nitrogen-phosphorus plots. On phosphorus plots the pattern of water use was similar to that on nitrogen plots, but water-use rates were slightly lower, particularly at the 36" depth. The rate of water use at the 6" depth, on the non-fertilized plot, was similar to that on fertilized plots, but rate of water use at the 12", 20", and 36"

Rainfall and moisture extraction patterns for fertilizer treatments as a function of time. Left—Non-fertilized and phosphorus-fertilized plots. Right—Nitrogen and nitrogen plus phosphorus plots.



depths was less and the wilting point was not reached by the end of the experiment.

Rainfall of 1.85" in May resulted in: 1, reduced suction at all depths on the nitrogen-phosphorus plot; 2, reduced suction at the 6" depth only, on the nitrogen plot; 3, reduced suction at the 6" and 12" depths on the phosphorus plot, but only a slowing of the rate of moisture removal at the 20" and 36" depths; and 4, reduced suction at the 6" depth on the non-fertilized plot but only a slight retardation of the rate of moisture extraction at the other depths.

A late rain, on May 20, was not used

by the plants on the nitrogen-phosphorus plots because they were completely matured and dry. A slight increase in suction at the 6" depth near the first of June probably resulted from evaporation at the soil surface. When this particular soil dries, fissures are opened and evaporation can occur within the soil profile. On the nitrogen plots, the June rain was used by the plants in the final stages of maturity, and only a slight amount of depletion could be related to evaporation. The plants on the phosphorus and non-fertilized plots were not so mature as those on the nitrogen plots and continued

to deplete the soil moisture until matured. The continuing rate of depletion of soil moisture on these plots appeared to be caused by the undesirable, late-maturing medusahead and summer-growing annual weeds.

Fertilization of annual range plants with nitrogen resulted in a greater forage production and earlier maturity than occurred on the plots with phosphorus only and on the check plots. Combination of nitrogen and phosphorus resulted in a significant interaction that gave increased yield of forage.

Abundant plant growth and more uniform early plant maturity, due to the application of nitrogen, removed more soil moisture than was removed from the soil in which plants did not receive nitrogen. Also, growth of summer-growing annual weeds was retarded considerably by depletion of soil moisture at all depths of the soil in plots where nitrogen fertilizer was applied.

Cyrus M. McKell is Plant Physiologist, Crops Research Division, A.R.S., U.S.D.A., and Associate in Agronomy, University of California, Davis.

Jack Major is Assistant Professor, Department of Botany, University of California, Davis.

Eugene R. Perrier is Laboratory Technician, Department of Irrigation, University of California, Davis.

The above progress report is based on Research Project No. 4635.

Growth of forage in response to various fertilizers. Left—nitrogen, 150 pounds per acre; center—nitrogen, 150 pounds, plus phosphorus, 200 pounds per acre; right—no fertilizer.



California Mastitis Test

for dairy herd improvement

O. W. SCHALM

About 30,000 cows in California are being screened every month by the California Mastitis Test—CMT—applied to milk samples collected for butterfat determination. Owners of dairy herds which show a high level of CMT-positive cows are receiving advice on management. Where indicated, medical treatment is given during the 2-3 months when the milk producing glands are in the resting or dry state.

The CMT is a chemical test for the immediate detection of high leucocyte—white cell—count, and it can be applied to milk samples on the farm. The discovery was reported in 1957 by the School

of Veterinary Medicine at Davis. Because the leucocyte content of milk is increased in all conditions which lead to irritation of tissues within the mammary gland, the CMT is a quick and inexpensive way to check all the cows in a herd for detection of udder abnormalities.

Part of the program of the Dairy Herd Improvement Association is a monthly analysis of udder health by CMT. By demonstrating the extreme importance of proper functioning and use of the milking machine in mastitis prevention, the program is leading to improved design of milking machine auxiliaries by manufacturers. Also, the local representatives

of milking machine companies have improved the services rendered to dairy-men, to maintain milking machines at peak efficiency.

In some herds, milk production has increased as much as 20% from one year to the next. The need for antibiotic treatment of clinical mastitis has been markedly reduced in herds which follow the recommended program.

Continuing research on bovine mastitis at Davis is a dual program involving large-scale herd trials as well as investigations on fundamental aspects of the problem. The nature of the chemical reaction which leads to a visual change in a high leucocyte milk sample is an interesting aspect. Other investigations are under way on the factors involved in development of bacterial infection within the mammary gland, so as to learn more about preventing such infection.

O. W. Schalm is Professor of Veterinary Science, University of California, Davis.

The above progress report is based on Research Project No. H 1064.