

Brown-bagging Granny Smith apples on trees stops codling moth damage

Walter J. Bentley □ Mario Viveros

In a 2-year study, thinning Granny Smith apples to one per cluster and covering them with brown paper bags when they were the size of a golf-ball resulted in significantly fewer fruits damaged by codling moth compared with fruits hand-thinned and left untreated. Sunburned fruit were also reduced both years by bagging, and fruit firmness and sweetness were improved in 1 of the 2 years. A similar experiment is now underway with Fuji apples.

During 1989 and 1990, field trials were conducted in a commercial apple orchard near Arvin, California, to evaluate the feasibility of growing Granny Smith apples in home gardens without using insecticides to control codling moth. Instead, small, brown paper bags were placed around individual apples as a barrier to oviposition by moths. The bags also protected the apples from sunburn. The Granny Smith cultivar was chosen because of its long period between bloom and harvest when the probability of infestation increases.

Since it was first detected in 1747 in an area that would soon become the United States, codling moth, *Cydia pomonella* (Linnaeus), has become the major pest of apples. In addition to apples, codling moth infests English and Asian pears, English walnut, quince, crab apple, hawthorne, and various stone fruits. However, apple and pear are preferred; when left unsprayed, a complete crop loss can result. In the southern San Joaquin Valley, codling moth is particularly troublesome because it can complete four generations in most years.

Home gardeners in particular are troubled by codling moth. Many do not

understand the importance of appropriately timing insecticide sprays. Often, they are applied when they do the least good, leading gardeners to step up the dosage or the frequency of application. That, in turn, can lead to unacceptable residues on fruit. Some gardeners are reluctant to apply synthetic materials and are unable to harvest fruit without worms. Many give up trying to grow apples.

Codling moth behavior

Female codling moths lay eggs either on leaves or on developing fruit. After the eggs hatch, the larvae enter the fruit through the calyx end, where two fruit touch, or along the sides of the fruit. Once in the fruit, they quickly move to the core where they feed on seeds and then exit to pupate or to overwinter as mature larvae in silken cocoons. Overwintering larvae can be found under tree bark, under plant debris, or on the soil. By March, wintering larvae are pupating and adult moths begin emergence.

Overwintering moth emergence extends from March through late May in the San Joaquin Valley, and in most years two distinct flight peaks occur. First summer generation flight occurs from early June through early July. By mid-July second summer generation adults are active; the third summer generation occurs in late August into September. Activity in the fourth and final flight (third summer generation) is not as prolonged as the previous flights. Typical seasonal flight activity is seen in figure 1.

Control of codling moth has been traditionally based on timely insecticide applications. If minimal sprays are to be used, they must be timed at egg hatch so that the larvae will pick up a lethal dose upon contact or while feeding. The only alternative to timing insecticides has been a calendar-based spray schedule. In areas with four generations, this could result in as many as 12 sprays with organophosphate or carbamate insecticides to keep apples worm free. If biological insecticides — such as various formulations of *Bacillus thuringiensis* (Bt) — are used, as many as 24 sprays could be required, with weekly applications from mid-March through August. This procedure is sometimes used by those attempting to market apples under the organic label.

The field trials

Twenty-four trees selected for the field trials, among the 518 grown per acre, were on a two-wire trellis with a 14- by 6-foot planting density. The trees, ranging in height from 12 to 15 feet and irrigated with low-volume sprinklers, received a standard dormant spray with an organophosphate and a supreme oil for control of San Jose scale and rosey apple aphid. No other pesticides were applied during the study.

The field trial consisted of three treatments applied to single trees and replicated eight times. The treatments in-




Fig. 1. Seasonal flight of codling moth, Arvin, California, 1990.

cluded: (1) bagging individual fruit after hand-thinning to one per cluster, (2) hand-thinning fruit to one per cluster but not bagging them, and (3) not thinning or bagging.

The bags used were #2 Townsend paper bags measuring 18.5 cm by 10.5 cm. In

categories: category 1, no evidence of sunburn; category 2, slight yellowing; category 3, a uniform yellowing of an exposed side; and category 4, some browning. Because the number of fruit varied from tree to tree, results of both codling moth infestation and sunburn are ex-

difference among treatments due to fruit loss from codling moth. However, more apples were harvested from the bagged and thinned treatments than from the others.

Where infestation levels are high, the percentage of infested fruit can be mis-

TABLE 4. Proportion of sunburned fruit classed into four categories*
Percentage of yield (lb) in each category†

Treatment	1		2		3		4	
	1989	1990	1989	1990	1989	1990	1989	1990
Bagged + thinned	86.5 A	96.1 A	8.0 A	3.1 A	3.8 A	0.6 A	1.8 A	0.3 A
Thinned	48.2 B	72.9 B	23.7 B	18.0 B	18.8 B	5.6 B	9.4 B	3.5 B
Non-thinned	52.5 B	69.6 B	20.4 B	19.1 B	16.7 B	6.2 B	10.4 B	5.2 B

*Categories: (1) no sunburn, (2) slight yellowing, (3) uniform yellowing, (4) browning.

†Means followed by the same letter in columns are not significantly different at the 1% level, by Duncan's multiple range test. (Harvested September 7, 1989 and August 29, 1990.)

TABLE 5. Quality effects of bagging and thinning Granny Smith apples

Treatment	PSI*		Soluble solids*		Diameter (mm)*	
	1989	1990	1989	1990	1989	1990
Bagged + thinned	18.8 A	20.2 A	14.2 A	13.0 A	73.5 A	73.6 A
Thinned	17.9 B	20.0 A	13.7 B	12.9 A	74.1 A	74.9 A
Nonthinned	17.5 B	20.2 A	13.5 B	13.0 A	71.9 B	71.9 B

*Means followed by the same letter in columns are not significantly different ($P < 0.01$), by Duncan's multiple range test. (Harvested September 7, 1989 [20 fruit per plot] and August 29, 1990 [24 fruit per plot].)

TABLE 6. Cost/revenue comparison of bagging and thinning with chemical pest control and thinning for Granny Smith apples*

Operation	Treatment	
	Chemical thinning and pest control	Hand thinning and bagging
Codling moth control and thinning costs/ac*		
Materials @ 4 treatments/ac†	\$140.00	\$2,274.00
Equipment	\$75.00	NA
Labor	\$25.00	\$7,770.00
Total cost/ac	\$240.00	\$10,044.00
Additional cost/ac (bagging costs minus chemical cost)	NA	\$9,804.00
Marketable yield/ac†	27,226 lb	35,804 lb
Estimated return/lb	\$0.15	\$0.15
Revenue/ac	\$4,083.90	\$5,370.60
Increased return/ac with bagging	NA	\$1,286.70
Net loss/ac (total revenue minus cost)	NA	\$8,517.30

*Other costs of production assumed to be equal, yield of 230 fruit/tree and labor based on 3 man-hours/tree (bagging and thinning) at \$5/hr. Four treatments assumed for codling moth control to equal the bagged treatment.

†Based on marketable yields of 72 lb/tree with 518 trees per acre and sunburn (4% sunburn in bagged and 73% in non-bagged).


sunburn by protecting fruit throughout the growing season with paper bags. As much as 30 to 40% additional yield in Grade 1 fruit can apparently be achieved with bagging in the southern San Joaquin Valley compared with nontreatment. This is mainly thanks to sunburn protection. Sunburn protection may not be as necessary in other parts of California where summers are not as hot. Where high winds occur and make paper bags not feasible, cloth bags with string ties could be used to provide comparable protection.

For individuals with few trees, bagging can result in quality fruit without significant loss due to codling moth or sunburn. This technique should be applicable to other apple cultivars as well, and at present a similar experiment with Fuji apples is underway. The experiment with Fuji apples may be commercially feasible inasmuch as consumers are apparently willing to pay top price for them. The technique should also be useful for Asian pears, which have to be thinned heavily and are therefore more expensive than other pears on the market.

Bagging can be done on the entire tree or on as many fruit as desired. This technique takes more time than spraying, but it demonstrates that apples can be grown without insecticides under heavy codling moth pressure during a long growing season. It also eliminates the problem of pesticide disposal often encountered by gardeners.

W. J. Bentley is Extension Entomologist and Mario Viveros is Extension Pomologist with UC Cooperative Extension, Kern County.

The authors wish to thank Martha Gonzalez, John Fritz, Lynette Beurman, and Nancy Medina for treatment and harvest assistance, as well as John Wood and Jim Carlisle of Green Valley Fruit for providing a site and managing the trial.



After thinning the immature fruit, Farm Advisor Mario Viveros encloses individual Granny Smith apples on the tree in brown paper bags. With its open end stapled shut, each bag will form a physical barrier to codling moth attack, and also prevent sunburn damage. A similar experiment is now being conducted with Fuji apples. *Inset:* Fruit diameters averaged 2.4 cm when bagged on May 1, 1990.

CALIFORNIA AGRICULTURE ASSOCIATE EDITORS

Animal, Avian, Aquaculture and Veterinary Sciences

Richard H. McCapes
(2nd assoc. editor to be announced)

Economics and Public Policy

Harold O. Carter
(2nd assoc. editor to be announced)

Food and Nutrition

Eunice Williamson

Human and Community Development

Linda M. Manton
Karen P. Varcoe

Land, Air & Water Sciences

Garrison Sposito
Henry J. Vaux, Jr.

Natural Resources

Daniel W. Anderson
John Helms
Richard B. Standiford

Pest Management

Thomas C. Baker
(2nd assoc. editor to be announced)

Plant Sciences

Calvin O. Qualset
G. Steven Sibbett

