



In addition to controlling noxious annual weeds, prescribed burns on rangeland can remove litter, recycle nutrients and stimulate the emergence of desirable perennial grasses.

References

- DiTomaso JM, Kyser GB, Hastings MS. 1999. Prescribed burning for control of yellow starthistle (*Centaurea solstitialis*) and enhanced native plant diversity. *Weed Sci* 47(2):233-42.
- Donald WW. 1991. Seed survival, germination ability, and emergence of jointed goatgrass (*Aegilops cylindrica*). *Weed Sci* 39(2):210-6.
- Hopkinson P, Fehmi JS, Bartolome JW, et al. 1999. Adaptive management and fire control of barb goatgrass. *Res Manage Notes* 17(3):168-9.
- Jacobsen WC. 1929. Goatgrass — A weed pest of the range. *Monthly Bull, Calif Dept Agr* 18(1):37-41.
- Kennedy PB. 1928. Goatgrass or wild wheat (*Aegilops triuncialis*). *J Am Soc Agr* 20(12):1292-6.
- Ludwig JA, Reynolds JF. 1988. *Statistical Ecology*. New York: J Wiley. 337 p.
- Menke JW. 1992. Grazing and fire management for native perennial grass restoration in California grasslands. *Fremontia* 20(12):22-5.
- Meyer MD, Schiffman PM. 1999. Fire season and mulch reduction in a California grassland: A comparison of restoration strategies. *Madroño* 46(1):25-7.
- Murphy AH, Lusk WC. 1961. Timing medusahead burns to destroy more seed and save good grasses. *Cal Ag* 15(11):6-7.
- Peters A, Johnson DE, George MR. 1996. Barb goatgrass: A threat to California rangelands. *Rangelands* 18(1):8-10.
- Talbot MW, Smith LS. 1930. The goatgrass situation in California. *Monthly Bull, Calif Dept Agr* 19(1):40-6.
- Willis BD, Evans JO, Dewey SA. 1988. Effects of temperature and flaming on germinability of jointed goatgrass (*Aegilops cylindrica* Host) seed. *Proc West Soc Weed Sci* 41:49-55.
- Young FL, Ogg AG Jr, Dotray PA. 1990. Effect of postharvest field burning on jointed goatgrass (*Aegilops cylindrica*) germination. *Weed Tech* 4(1):123-7.


Animals and fungi can affect goatgrass establishment

Valerie T. Eviner
F. Stuart Chapin III

The establishment of barb goatgrass is greatly affected by its interactions with other species in California grassland. For example, goatgrass establishment is enhanced by a fungus, *Ulocladium atrum*. This fungus is visible as black lesions on the surface of the seedhead; it helps break down the tough, woody seedhead surrounding goatgrass seeds. This speeds up germination and leads to a 65% increase in goatgrass biomass at the end of the growing season.

In contrast, small mammals help to slow goatgrass invasion. In two separate experiments, more than 3,000 individually marked goatgrass seedheads were laid out in plots at the UC Hopland Research and Extension Center (HREC). Within days, new rodent holes had formed in the areas where the seeds were planted, and every single seedhead was missing. It was not determined if this predation was due to mice or voles.

Gophers have a dramatic effect on goatgrass establishment. We established 10.76-square-foot plots of eight different plant species commonly found in California annual grassland at HREC. There were 18 replicate plots of each species randomly distributed throughout the study area. Gophers selectively built mounds in goatgrass plots, completely burying the goatgrass and killing most of the seedlings. Gophers buried every goatgrass plot, while only a



Controlling gophers on rangeland can have an unintended side effect: dramatically increased establishment of barb goatgrass. In experiments at Hopland, gophers selectively disturbed goatgrass plots, burying almost all the seedlings, left, while goatgrass plots protected from gophers have dense seedling establishment.

handful of other plots had significant levels of gopher disturbance. Goatgrass biomass was 3.5 times higher in gopher exclosures than in the presence of gophers. A preliminary survey of the pastures at HREC indicated that where gophers were present, goatgrass coexists with other plant species, while in areas without gopher activity, goatgrass formed a dense monoculture stand.

While much more research is needed to determine the relative importance of the fungus *Ulocladium*, mice and voles, gophers and competition with plants on goatgrass establishment, it is clear that these interactions can play a significant role in the establishment of this invasive species.

V.T. Eviner is Ph.D. Candidate, Department of Integrative Biology, UC Berkeley; and F.S. Chapin III is Professor, Institute of Arctic Biology, University of Alaska, Fairbanks.