

# RANGE SCIENCE REPORT

Agricultural Experiment Station

Cooperative Extension

NO. 35

April 1998

## Water Quality Effects of Rangeland Beef Cattle Excrement

Glenn Nader<sup>1</sup>, Kenneth W. Tate<sup>2</sup>, Robert Atwill<sup>3</sup>, and Dan Drake<sup>4</sup>

Causes of rangeland nonpoint source pollution are difficult to identify and quantify due to the complexity of watersheds. Concerns regarding rangeland cattle excrement on water quality has focused on nutrient and pathogen loading to water bodies. Pollutants from rangeland beef cattle excrement can effect water quality if the pollutants are directly deposited in the water body or if they are transported to the water body during storm events.

Nutrient and pathogen dynamics on rangeland watersheds are complex and vary through time and space. Often background levels of nutrients and pathogens are unknown. The spatial distribution of rangeland beef cattle excrement varies within and across watersheds. Cattle distribution is a function of topography, soils, vegetation, air temperature, water and supplement location, animal class, and fencing.

Some studies have attempted to compare cattle excrement deposited on the range to human waste deposited in a septic system (Lahonton 1985, DWR 1979). An important concept involved in this comparison is that humans import their food sources into a watershed while cattle predominantly consume forage produced in the watershed. Cattle export nutrients out of the watershed in the form of body mass. Beef calves that gain 2 to 2.5 pounds per day, of which 2.4% is nitrogen, 0.8 % is phosphorus (Azevedo and Stout 1974), illustrate the amount of nutrients that can be removed from grazed watersheds.

Nitrogen exported in tissues of domestic ungulates was estimated to be 17% of the N in ingested forage (Dean et al. 1975). Thus, assuming 70% moisture and 10% crude protein forage, for every ton of forage consumed about 1.6 pounds of nitrogen is removed from the system.

---

<sup>1</sup> U.C. Cooperative Extension Farm Advisor, Sutter/Yuba/Butte Counties.

<sup>2</sup> Rangeland Watershed Specialist, U.C. Cooperative Extension, Agronomy and Range Dept., U.C. Davis.

<sup>3</sup> U.C. Dept. of Population Medicine and Reproduction and Veterinarian Medicine Extension, Vet. Med. Teaching and Research Center, Tulare, CA.


<sup>4</sup> U.C. Cooperative Extension Farm Advisor, Siskiyou County.

To understand and quantify the fate of nitrogen on a watershed several things must be considered:

- 1.) Quality and quantity of the forage.
- 2.) Retention by the animal.
- 3.) Losses through volatilization and leaching of  $\text{NH}_3$ .

- 4.) Soil incorporation.
- 5.) Plant uptake.
- 6.) Spatial distribution of the feces and urine.
- 7.) Hydrology of the watershed.

Figure 1. illustrates the complex processes of the nitrogen cycle on a watershed, and the role of herbivores in the nitrogen cycle.



**Figure 1. Nitrogen Cycle**

Evaluation of nutrient discharges must also consider the natural input by rain. Olness et al. (1975) found that the rangeland watershed received more total inorganic nitrogen in rainfall than was lost in with surface runoff. Ritter (1986) found that both nitrogen and phosphorus contributed by rainfall was greater than the rates occurring in stream flow. Menzel et al. (1978) during a 4-year study found rainfall added four times the nitrogen compared to nitrogen discharged in runoff from rotational grazed pastures and about equaled the amount discharged from continuously grazed pastures.

Evaluation of rangeland cattle excrement impacts on water quality require consideration of natural variability in the hydrologic cycle, the nutrient cycles, and the pathogen cycle as well as how grazing modifies each of these processes. These factors make it difficult to quantify the amount of nutrient and pathogen loading that is attributable to rangeland beef cattle on a watershed. The purpose of this paper is to review the literature to evaluate our current knowledge of the potential for rangeland beef cattle excrement to impact water quality.

### Components of Range Cattle Excrement

The feces consist of slowly digestible plant material, enzymes, mucous cells of the digestive tract, pathogens, surplus salts, wastes from cellular metabolism, and water (Azevedo and Stout 1974). Urine contains waste products of metabolism. The amount of excreted feces will depend on forage intake (driven by body size and physiological function) and digestibility.

The actual amount of urine produced daily varies according to production (growth, lactation, or conception), air temperature, and water consumption. (NRC 1984). A review of range forage intakes by Cordova et al. (1978) showed that they were highly variable with a ranging from 40 to 90 g Dry Matter /Weight(kg)<sup>75</sup>. Several studies in the western United States estimates intake ranging from 1 to 2.8% of body weight. Many water quality studies (Lahonton 1985, DWR 1979) are based on confined beef cattle excretion data, which should be used as only crude first estimates of rangeland cattle excretion. The following are values used by agricultural engineers based on a wide range of confined beef cattle diets and conditions. Beef cattle produce 30 to 49 pounds of urine per day. Beef cattle produce between 29 and 72 lb. of feces per day. For every ton of live animal mass, beef cattle excrete 0.748 lbs. of Kjeldahl nitrogen, 0.189 lbs. of ammonia nitrogen, 0.20 lbs. of phosphorus per day (ASAE 1992). The ASAE data did not give a dry matter output, but Azevedo and Stout (1974) give the range of percent of fresh weight to dry matter of 15 - 27%. The range in confined beef cattle output was 29 to 60 lbs. of raw and 4.6 to 10.2 lbs. on a dry matter basis.

Actual rangeland fecal output studies using collection bags illustrate the amount of variation found under rangeland conditions (Table 1). Authors report their findings on a daily fecal output on a dry matter basis. Connor et al. (1963) was the only one to publish the dry matter digested percents which were 40.4 for Southern Nevada and 53.3 for Northern Nevada.

Table 1. - Fecal Output Studies

| Animal Weight (Lbs.) | Daily Fecal Output (Lbs.) | Location       | Forage             | Month           | Author                    |
|----------------------|---------------------------|----------------|--------------------|-----------------|---------------------------|
| 460                  | 3.78                      | So. Nevada     | desert Shrub       | July to October | Connor et al. 1963        |
| 460                  | 2.68                      | No. Nevada     | sagebrush/grass    | June to Sept.   | Connor et al. 1963        |
| 605                  | 5.1 - 7                   | Eastern Oregon | crested wheatgrass | April-May       | Handle et al. 1972        |
| 605                  | 5.1 - 7                   | Eastern Oregon | crested wheatgrass | June            | Handle et al. 1972        |
| 726                  | 5.9                       | Nebraska       | tall wheatgrass    | September       | Adams et al. 1991         |
| 880                  | 5.5                       | Nebraska       | meadow             | July            | Hollingsworth et al. 1995 |
| 880                  | 8.4                       | Nebraska       | meadow             | September       | Hollingsworth et al. 1995 |
| 880                  | 7.9                       | Nebraska       | meadow             | October         | Hollingsworth et al. 1995 |

## **Nutrients**

Nitrogen and phosphorus are the nutrients which cause most water quality concerns.

### **Nitrogen**

Cattle intake nitrogen mainly in the form of plant protein. Nitrogen is lost through eructation, belching, and excretion. The amount of nitrogen consumed by beef cattle that is utilized depends on the demands for growth, maintenance, reproduction, and lactation. This leads to a wide variation in reported utilization. Young calves utilize about 42% of consumed nitrogen (Salter and Schollenbergen 1939). Woodmansee et al. (1981) stated that cattle commonly retain 15 to 20% nitrogen of ingested forage. While Afzal and Adams (1992) indicated that typically 75% of the ingested N is returned in dung and urine. Azevedo and Stout (1974) reported nitrogen was excreted in urine (47.6%) and feces (52.4%) by weight. However, many researchers suggest urea accounts for about 75 percent of the excreted nitrogen (Afzal and Adams 1992). Excreted nitrogen, mainly in the form of urea, is rapidly hydrolyzed by ubiquitous urea-decomposing enzymes yielding ammonia. More than 80% of the nitrogen in urine may be lost by volatilization.

Under simulated feedlot conditions, Stewart (1970) found that 85 to 90% of nitrogen in urine was lost as ammonia. Under ambient conditions losses are probably about 50%, which is the often used value (Woodmansee et al. 1981). Nitrogen in feces that is not volatilized is slowly released from complex organic compounds present in manure as a result of microbial activity. The microorganisms which decompose manure demand a carbon-to-nitrogen ratio of less than 15 or 20 before ammonia can be split off and released from nitrogenous organic compounds in sufficient quantities for good plant growth. As decomposition proceeds, the various organic constituents of the substrate are attacked at different rates. Stewart (1970) reported that 37.3% of the nitrogen present in fresh feces was volatilized within one week. Wilkinson and Lowery (1973) reported soil N is affected in an

area .97 square feet around each defecation 3.0 square feet around each urination. Grass growth was affected within an area of 10.76 square feet around each urination spot. Woodmansee et al. (1981) estimated that in low productivity systems, the amount of nitrogen added in one urine patch may be 10 times greater than the uptake capacity of the plants, and in highly productive systems, the amount of nitrogen added maybe three times the uptake capacity. Nitrogen not taken up by the plants may be immobilized by soil microorganisms or eventually transferred to soil organic matter. Ammonium may be absorbed onto soil colloids or fixed and lost from the rapid cycling pools, but would slowly become available. Afzal and Adams (1992) found that total mineral nitrogen under feces was always shallow (0-.78 inches). The depth of total mineral nitrogen from urine changed with time. The change in form of total mineral nitrogen to nitrate and depth was observed at 56 days after simulated urine application with an increase of nitrate from 61% in the 0 to .78 inches depth to 98% in the 1.57 to 2.36 inches depth. Dormaar et al. (1990) found that grazing did not change the total nitrogen in the Ah horizon, but the forms were different with higher ammonia and nitrate present. Nitrate is susceptible to loss by leaching if precipitation is heavy, but in most grasslands such losses are probably small (Woodmansee et al. 1981). Most elements in feces of large animals are bound in relatively resistant organic fractions (Float and Torrance 1970). The bulk of bound elements remains for many years at the surface in feces (Angel and Wicklow 1975). Fecal nitrogen is very efficient for plant growth because of the slow release (Dormaar et al. 1990). Cattle grazing removes herbage from large areas in a pasture, but deposits feces in a small area. Buckhouse and Gifford (1976) found .02% of a semi arid range covered with bovine feces under a stocking rate of 4.9 acres/AUM. Uneven distribution of excreta may affect the nitrogen cycling in the soil-plant-animal system.

### **Phosphorus**

Phosphorus is an essential nutrient for growth, maintenance, lactation, and reproduction of cattle.

Phosphorus and calcium are important in the formation of bones. Dietary phosphorus retained by the animal varies from 78% for growing calves to 58% for lactating cows (NRC 1984). Every 2.2 pounds of calf gain contains nine grams of phosphorus, while every 2.2 pounds of cow gain contains six grams. Most excreted phosphorus (97.3%) is in the feces. Of the nutrients present in manure, phosphorus is the second most resistant to leaching. In general, phosphorus from applied manure is not leached from soils (Azevedo and Stout 1974). Phosphorus is rapidly hydrolyzed and chemically precipitated or absorbed by other soil minerals. Most soils are able to rapidly tie up large amounts of this element in forms not readily available to plants. Phosphorus is absorbed by plants as  $H_2 PO_4$ ,  $HPO_4$  or  $PO_4$ . Most soil phosphorus is tied up chemically in compounds of limited solubility. In neutral to alkaline soils, calcium phosphate is formed, while in acid soil, iron and aluminum phosphates are produced.

## **Pathogens**

### **Bacteria**

The primary pathogenic bacteria found in beef cattle excrement includes *Escherichia coli*, *Leptospira interrogans*, *Salmonella spp.*, *Campylobacter jejuni* and *Yersinia enterocolitica* (Gary et al. 1983, Altekruze et al. 1994, Whipp et al. 1994).

### **Protozoal Parasites**

The primary water-borne protozoa potentially transmitted by cattle excrement includes *Cryptosporidia parvum* and *Giardia duodenalis* (also known as *Giardia lamblia*) (Fayer and Ungar 1986, Craun 1990, Atwill 1996). *C. parvum* is a tiny protozoal parasite that can cause gastrointestinal illness in a wide variety of mammals, including humans, cattle, sheep, goats, pigs, and horses. It also occurs in various wildlife species such as deer raccoons, opossums, rabbits, rats, mice, and squirrels (Fayer and Ungar 1986). In cattle, shedding of the parasite is usually limited to calves, but there are a few reports of subclinical shedding in adult cattle (Lorenzo et al. 1993). Dairy calves are commonly infected with *C. parvum* and *G. duodenalis* (Ongerth and Stibbs

1989, Xiao 1994), but little is known of their distribution in beef cattle herds, particularly in those herds located on open range.

## **Water Quality Impacts Related to Rangeland Beef Cattle Excrement**

### **Nutrients**

Hathaway and Todd (1993), studied the contribution of different cultural activities in the Wood River sub basin in eastern Oregon. They found that continuously grazed irrigated meadows did not influence the nitrogen loading of streams. The daily load of phosphorus was lower on the downstream grazed area than the upstream pristine land use area. Gary et al. (1983) studied a moderately grazed pastures bisected by a small perennial stream in central Colorado. Only minor effects on water quality were detected during a two-year study. Cow excretion was monitored for an eleven hour period both years and 6.7 to 10.5% of the defecations and 6.3 to 9% of the urinations were deposited directly in the stream. Nitrate nitrogen did not increase, and ammonia nitrogen increased significantly only once during this study. Tanner and Terry (1991) found no significant differences in N, P, chlorophyll, dissolved oxygen, and pH of surface water collected from light to moderately grazed and ungrazed wetlands in south Florida. Dahlgren and Singer (1991) found that grazing of northern California oak woodlands had no effect on major nutrients. Coltharp and Darling (1975) found no difference in water chemical levels between grazed and ungrazed areas along three mountain streams. Robbins (1979) stated that all the available data indicate that pollutant yields from rangeland are not directly related to the number of animals or amount of waste involved, but are related to hydrological and management factors involving erosion/sedimentation.

### **Pathogens**

Detailed studies that attempt to link rangeland cattle grazing with the presence of water-borne pathogenic bacteria have for the most part not been done (Atwill 1996). Instead, indicator bacteria have been used. These studies need to be

interpreted with some caution since indicator bacteria have been shown to be poorly correlated with some pathogenic bacteria such as *Campylobacter jejuni* (Carter et al. 1987, Bohn and Buckhouse 1985). An increase in indicator bacteria in waterways, due to cattle grazing has been documented in many studies (Gary et al. 1983, Robbins 1979, Dixon et al. 1979, Stephenson and Street 1978). However grazing has also been found to have little or no effect on fecal indicator counts (Frear 1983, Buckhouse and Gifford 1976). Fecal indicators may not always signify the presence of pathogens in the water column (Bohn and Buckhouse 1985). When contamination does occur, it may be temporary and short-lived (Gary et al. 1983, Robbins 1979), or may persist for several months. (Stephenson and Street 1978). Furthermore, concentrations tend to decrease downstream (Robbins 1979).

A special concern for bacterial pollutants is their ability to survive in the environment, only to become a factor in pollution at a later time. Bacteria such as *Salmonella newport* and *E. coli* have been shown to survive several months in freshwater sediments (Burton et al. 1987). Fecal coliforms may survive up to two months in soil, but in the protective medium of feces, can persist up to a year (Bohn and Buckhouse 1985). Bottom sediments have been found to harbor concentrations of indicator organisms up to 760 times greater than the overlying water (Stephenson and Rychert 1982).

Studies that carefully evaluate the association between rangeland cattle and the presence of these water-borne protozoa have not been done. The majority of the existing literature deals with dairy cattle, or was conducted in laboratory settings. These studies do not explicitly state how the cattle were managed nor define the cattle's proximity to contaminated water bodies. Madore et al. (1987) measured 5,800 *Cryptosporidium* oocysts/L in irrigation canal water running through agricultural acreage with cattle pastures compared to 127 oocysts/L in river water subject to human recreation and 0.8 oocysts/L for stream water exposed to ranch land runoff. Unfortunately, the authors do not specify if the cattle were beef or

dairy cattle or if the species of *Cryptosporidia* was the of human health concern, *parvum*. Presently, there are no data that indicates rangeland cattle are a significant source of *C. parvum* (Atwill 1996).

### **Spatial Distribution of Cattle Excrement**

Nonpoint pollution caused by cattle excrement may be aggravated or ameliorated by the proximity of deposition to water bodies. Deposition outside of riparian areas may pose no pathogen or nutrient problems (Blackburn et al. 1982). Larsen et al. (1993) utilized a rainfall simulator in a laboratory environment to assess the effectiveness of vegetative filter strips to attenuate fecal coliforms, a questionable indicator of pathogens. Results during a 30 minute simulation indicate that distance of the fecal material up slope from the collection point significantly influences loading. In a situation where a stream was the only source of water and cattle spent 65% of the day within 328 feet of the stream channel 6.7 to 10.5% of defecations and 6.3 to 9.0% of urinations were deposited directly into the stream (Gary et al. 1983). Larsen et al. (1988) found that free ranging cattle in Oregon deposited an average of 3.4% of their feces in the stream in August and 1.7% in November. Powell (1984) found that direct overland flow movement of dung into stream channels was minimal because of standing vegetation and ground litter.

On a grazing allotment in the Blue Mountains of Oregon, riparian areas covered only 2% of the acreage, yet accounted for 81% of the herbaceous vegetation consumed by cattle (Kauffman and Krueger 1984). The preference of cattle for riparian areas has been attributed to the quality and quantity of forage, slope, microclimate, and availability of water (Bryant 1982, Marlow and Pogacinik 1986, Kauffman et al. 1983). The season of grazing can also greatly affect the use of upland pastures. Cattle do find other watershed areas preferable to riparian areas at various times of the year. Bryant (1982) found that from mid-July to mid-September, cattle in northeastern Oregon tended to graze under the forest canopy both for the shade provided and the quality of

forage during this period. He also found that from mid July to mid August, cows tended to stay closer to water than yearlings. From mid August to mid September, both tended to remain near the watering sites equally. From mid September to mid October, water was ineffective for distributing both cows and yearlings. It is generally found that cows not nursing calves and yearlings tend to distribute more on the rangeland. In tallgrass prairie, the preference for bed grounds and shaded areas played a larger role in dung distribution than did quality of forage (Powell 1984). Uplands are favored over forested and riparian zones late in the season and following thunderstorms (Fear 1983, Bryant 1982). Dung fouling could act as a natural concentration constraint for high stocking rates, as cattle will avoid grazing an area 6 times the size of the feces (Sweeten and Reddell 1978).

### **Strategy for influencing livestock distribution**

Livestock's distribution within a watershed can be manipulated using sound range management practices such as salting, water location, fencing, and selecting against cattle that graze riparian areas. Salt, mineral or protein supplements placed next to the streams can result in direct pollution of the water as well as increase cattle dung, urine and trampling next to the stream. Salt should be placed in areas away from stream courses to help distribute cattle. It is best to familiarize animals with the location of salt by driving them there, especially in an area not frequently grazed. Alternative water sources, such as windmill or solar powered wells, reservoirs, and guzzlers, can be developed in upland areas to draw cattle away from streams. Miner et al. (1992) found that a water trough 328 feet from a stream during the winter reduced the amount of time cattle spent in a stream by 90%. In the spring time, Clawson (1993) found that water trough placement reduced the range of stream use from (3.9 - 8.3) to (.9 - 4.7) minutes/cow/day. He also found that a water gap completely eliminated fecal deposition into the stream. Livestock distribution away from riparian areas may be improved through training and selection (Gillen et al. 1984, Howery 1993, Roath and Krueger 1982, Walker 1995).

Subdividing large pastures to exert more control over the frequency and timing of grazing can be used to improve grazing distribution. Rotational grazing management can be used. Continuously grazed rangelands contributed at least four times more nitrogen and phosphorous to the watershed compared to rotationally grazed rangelands. (Khaleel et al. 1979).

## **Conclusions**

### **Nutrients**

Water quality data should be examined carefully before assigning a cause and effect relationship between cattle grazing and non point pollution. Natural background levels of nutrient and pathogen loading can be quite high during storm events. Non point pollution from pastured and rangeland livestock depends on the stocking rate, length of grazing period, the season of use, manure deposition sites and concentration. Normally, pastures and rangelands have not presented water quality problems caused by cattle excrement, except under special circumstances. Several studies have concluded that cattle excrement contributes negligible nutrient pollution to waterways (Hathaway and Todd 1993, Frear 1983, Robbins 1979, Dixon 1979, Johnson et al. 1978, Tanner 1991, Coaltharp and Darling 1975, Milne 1976). Unfortunately, none of the studies defined the treatments well enough to describe the intensity and timing of grazing. The main water quality concerns are from cattle feces and urine deposited directly into the water. Potential problems occur in cases where animals congregate for feeding, watering, resting, in proximity to waterways, (Khaleel et al. 1979).

There is little scientific evidence that excrement from beef cattle on rangelands significantly impacts water quality. When significant nutrient contaminations do occur, especially phosphorus, they are more likely explained by erosion and sediment processes in the watershed (Khaleel et al. 1979, Robbins 1979). Cattle can effect the erosion and sediment process through vegetation removal.

### **Pathogens**

The scientific evidence implicating beef cattle as a significant source of *C. parvum* or *G. duodenalis* for surface water is incomplete and contradictory. Given the lack of scientific investigation, it would be premature to claim that rangeland cattle production is the leading source of *C. parvum* or *G. duodenalis* for surface water contamination (Atwill 1996). Rangeland beef cattle excrement may increase pathogen contamination in water ways beyond background levels, but studies have shown that background levels are not zero. Wildlife species, including muskrats, coyotes, mule deer, waterfowl, elk, etc. shed pathogenic bacteria such as *Campylobacter jejuni* (Altekruse et al. 1994). *Giardia* has been repeatedly isolated from wildlife (Thompson and Reynoldson 1993). Furthermore, high counts of indicator bacteria are often found upstream from grazed areas and are attributed to wildlife (Gary et al. 1983). Concentrations of *Cryptosporidium* oocysts from pristine surface waters have been 0.005-18 oocysts/L, indicating that this organism occurs naturally in pristine watersheds (Sterling and Arrowood 1993).

### **Management Implication**

Rangeland water quality can be managed by implementing spatial distribution of cattle through salting, upland water developments, fences for pasture rotation, and even by training or selection of the cattle grazed. These methods address the deposition of excrement near waterways, but also other, hydrologic, ecologic, and economic issues.

### **Future Direction**

Future research needs to be focused directly on monitoring grazing impacts on nutrient and pathogen dynamics at the watershed scale. Clearly defining the site conditions, grazing management, and excrement depositional patterns on the watershed are critical for interpreting and applying this information.

### **Literature Cited**

- Adams, D.C., R.E. Short, M.M. Borman, and M.D. Macneil. 1991. Estimates of fecal output with an intra-ruminal continuous release marker device. *J. Range Man.* 44(3)204-207.
- Afzal, M and W. A. Adams 1992. Heterogeneity of soil nitrogen in pastures grazed by cattle. *Soil Sci. Soc. of Am. J.* 56: 1160-1166.
- Altekruse S.F., J.M. Hunt, L.K. Tollefson, and J.M. Madden. 1994. Food and animal sources of human *Campylobacter jejuni* infection. *J. Amer. Vet. Medicine Association.* 204:57-61.
- Angel, K. and D. T. Wicklow 1975. Relationships between coprophilous fungi and fecal substrates in Colorado grasslands. *Mycologia* 67: 63-74
- ASAE. 1992. Manure production and characteristics. American Society of Agriculture Engineers Standards (D384) 485-487.
- Atwill, E. R. 1996. Assessing the Link between Rangeland Cattle and Waterborne *Cryptosporidium parvum* Infections in Humans. *Rangelands* 18 (2), April 1996.
- Azevedo, J., and P.R. Stout. 1974. Farm animal manures: An overview of their role in the agricultural environment. California Agric. Exp. Station Extension Service. Manual 44. Ag. Publication. Univ. of Calif. Berkeley, Calif. 94720
- Blackburn, W.H., R.W. Knight, M.K. Wood. 1982. Impact of grazing on watersheds: a state of knowledge. Texas A&M Univ., 32p.
- Bohn, C.C. and J.C. Buckhouse. 1985. Coliforms as an indicator of water quality in wildland streams. *J. Soil and Water Conservation.* 40:95-97.



- Bohn, C.C. and J.C. Buckhouse. 1981. Interpreting coliform counts in northeast Oregon rangeland streams. 1981 Research in Rangeland Management. Oregon State University Agric. Exp. Station. Special Report 620. June 1981. pp. 15-17.
- Bryant, L.D. 1982. Response of livestock to riparian zone exclusion. *J. of Range Manage.* 35:780-785.
- Buckhouse, J.C. and G.F. Gifford. 1976. Water Quality Implications of cattle grazing on a semiarid watershed in southeastern Utah. *J. Range Manage.* 29:109-113.
- Burton G.A., Gunnison D and Lanza G.R. 1987. Survival of pathogenic bacteria in various freshwater sediments. *Applied Envir. Microb.* 53:633-638.
- Carter A.M., Pacha R.E., Clark G.W. and Williams E.A. 1987. Seasonal occurrence of *Campylobacter* spp. in surface waters and their correlation with indicator bacteria. *Appl. Environ. Microbiology* 53:523-526.
- Clawson, J.E. 1993. The use of off-stream water developments and various water gap configurations to modify the watering behavior of grazing cattle. M.S. Thesis, Oregon State Univ., Corvallis, Or. Pp. 80.
- Coltharp, G.B. and L.A. Darling. 1975. Livestock grazing a nonpoint source of water pollution in rural areas. *Water Pollution control in low Density Areas.* University Press of New England. Hanover, New Hampshire. p. 341-352.
- Connor, J. M., V.R. Behman, A.L. Lesperance, and F.E. Kinsinger. 1963. Nutritive evaluation of summer range forage with cattle. *J. Anim. Sci.* 22:961-969.
- Cordova F. J., J.D. Wallace, and R.D. Pieper. 1978. Forage intake by grazing livestock: A Review. *J. of Range Manage.* 31 (6) 430-436.
- Council for Agriculture Science and Tech Agriculture and Groundwater Quality Report No. 103, May 1985 pp. 19-37, ISSN 0194-4088. Ames, Iowa
- Craun G.F. 1990. Waterborne giardiasis. In: *Giardiasis*, E.A. Meyer (ed.), Elsevier Science Publishers, Amsterdam.
- Dept. Of Water Resources (DWR) State of California. 1979. Status Report on Lake Davis.
- Dixon, J.E., G.R. Stephenson, A.J. Lingg, D.V. Naylor, and D.D. Hinman. 1979. Stocking rate influence upon pollutant losses from land wintering rangeland cattle: a preliminary study. *Amer. Soc. of Ag. Engineers. Livestock Waste: A Renewable Resource:* 261-264. St. Joseph, Michigan.
- Dormaar, J. F., S. Smoliak, and W.D. Willms 1990. Distribution of nitrogen fractions in grazed and ungrazed fescue Ah horizons. *J. Range Manage.* 43 (1) 6-9.
- Dean, R. E. , J. E. Ellis, and R. W. Rice 1975. Nutrient removal by cattle from a shortgrass prairie. *J. Appl. Ecol.* 12 (1) 25-29.
- Dahlgren, R. and M. J. Singer 1991. In: *Proceedings of the Symposium on Oak Woodlands and Hardwood Rangeland Management*, Davis, Ca. USDA Forest Service General Technical Report PSW-126. 337-341.
- Fayer, R., and B. Ungar. 1986. *Cryptosporidium* spp. and cryptosporidiosis. *Microbiol. Rev.* 50:458-483.
- Floate, M.J. and C.J. Torrance. 1970. Decomposition of the organic materials from hill soils and pastures. *J. Sci. Food Agric.* 21:116-120.
- Frear, S.T. 1983. High country streams, cattle are compatible. *Beef* May edition 68-69. Gary, H.L., and J.C. Adams. 1984. Indicator bacteria in water and stream sediments near the snowy range in southern Wyoming. *Water, Air, and Soil Pollution.* 25:133-144.
- Gary, H.L., S.R. Johnson, and S.L. Ponce. 1983. Cattle grazing impact on surface water quality in a Colorado front range stream. *J. Soil and Water Conserv.* 38:124-128.
- Gillen, R.L., W.C. Krueger, and R.F. Miller. 1984. Cattle distribution on mountain rangeland in northeastern Oregon. *J. Range Manage.* 49:386-400.
- Handl. W.P. and L.R. Rittenhouse. 1972. Herbage yield and intake of grazing steers. *Proc. West. Sec. Amer. Soc. Sci.* 23:197:200

- Hathaway, R.L., and R. Todd. 1993. Wood River sub-basin water quality as related to land use associated with forest and livestock grazing. Final Technical Completion Report, Oregon Water Resources Research Institute, Oregon State Univ., Corvallis, Oregon. 97331
- Hollingsworth, K.J., D.C. Adams, T.J. Klopfenstein, J.B. Lamb, and G. Villalobos. 1995. Supplement and forage effects on fecal output estimates from an intra-ruminal marker device. *J. Range Man.* 48(2)137-140
- Howery, L.D. 1993. Social factors influence intraspecific differences in distribution patterns among individuals in cattle herd. Ph.D. Diss. Utah State Univ. Logan, UT.
- Kauffman, J.B. and W.C. Krueger. 1984. Livestock impacts on riparian ecosystems and streamside management implications... a review. *J. Range Manage.* 37:430-436.
- Kauffman, J.B., W.C. Krueger, and M. Vavra. 1983. Impacts of cattle on streambanks in Northeastern Oregon. *J. Range Manage.* 36:683-685.
- Khaleel, R., D.R. Keddy, and M.R. Overcash. 1979. Transport of potential pollutants in runoff water from land areas receiving animal wastes: a review. *Water Resources Res.* 14:421-436.
- Lahonton Regional Water Quality Control Board, Staff Report for Basin Plan Amendments for Eagle Lake. 1985. South Tahoe, Calif.
- Larsen, R.E., J.C. Buckhouse, J. A. Moore, and J. R. Miner 1988. Rangeland Cattle and manure placement: A link to water quality. *Proceedings of Oregon Academy of Science* 24:7-15 Oregon Academy of Science, Portland Or.
- Larsen, R.E., J. R. Miner, J.C. Buckhouse, and J. A. Moore. 1993 Water Quality benefits of having cattle manure deposited away from the streams. *Bioresources Technology*, 48 p 113-118.
- Lorenzo M.J., E. Ares-Mazás and V. Martínez de Maturana I. 1993. Detection of oocysts and IgG antibodies to *Cryptosporidium parvum* in asymptotic adult cattle. *Veterinary Parasitology*. 47:9-15.
- Madore M.S., J.B. Rose, C.P. Gerba, M.J. Arrowood, and C.R. Sterling. 1987. Occurrence of *Cryptosporidium* oocysts in sewage effluents and selected surface waters. *J. Parasitology*. 73:702-5.
- Marlow, C.B. and T. M. Pagoacknik 1986. Cattle feeding and resting patterns in a foothills riparian zone. *J. of Range Management* 39. P 212-216.
- Menzel, R.G.; Rhoades, E.D.; Olness, A.E. and Smith, S.J. 1978. Variability of annual nutrient and sediment discharges in runoff from Oklahoma cropland and rangeland Nutrient and Bacterial Pollution. *J. Envir. Qual.*, Vol. 7, No. 3, p. 401-406.
- Milne, C. M. 1976. Effect of a livestock wintering operation on western mountain stream. *Transactions of the ASAE* 19(4):749-752.
- Miner, J.R., J.C. Buckhouse, and J.A. Moore. 1992. Will a water trough reduce the amount of time spend in the stream. *Rangelands* 14(1) p.35-38.
- NRC. 1984. Nutrient requirements of beef cattle. Sixth edition.
- Olness, A.; Smith, S.J.; Rhoades, E.D.; and Menzel, R.G. 1975. Nutrient and sediment discharge from agricultural watersheds in Oklahoma. *J. Environmental Quality*. 43:331-336.
- Ongerth J.E. and Stibbs H.H. 1989. Prevalence of *Cryptosporidium* infection in dairy calves of western Washington. *American J. Veterinary Research*. 50:1069-1070.
- Powell, J. 1984. Dung distribution on a tallgrass prairie watershed. University of Wyoming, Range Management Department. Laramine, Wy. 82071.
- Ritter, W.F. 1986. Water quality of agricultural coastal plain watersheds. *Agricultural Wastes* 16:201-216
- Roath, L.R. and W.C. Krueger. 1982. Cattle grazing and behavior on forested range. *J. Range Mange.* 35:332-338
- Robbins, W.D. 1979. Impact of Unconfined Livestock Activities on Water Quality *Transactions of the Amer. Soc. Ag. Engineering*. p.1317-1323.
- Salter, R.M., and C.J. Schollenbergen. 1939. Farm Manure. *Ohio Agr. Exp. Sta. Bul.* 605, p.65.

- Stephenson, G.R. and R.C. Rychert. 1982. Bottom sediment: a reservoir of *Escherichia coli* in rangeland streams. *J. Range Manage.* 35:119-123.
- Stephenson, G.R., and L.V. Street. 1978. Bacterial variations from a southwest Idaho rangeland watershed. *J. Environ. Qual.* 7:150-157.
- Sterling R.S. and Arrowood M.J. 1993. Cryptosporidia. In: *Parasitic Protozoa*. J.P. Kreier (ed.), Academic Press, Inc. San Diego, California.
- Stewart, B.A. 1970. Volatilization and nitrification of nitrogen from urine. *Environmental Science Tech.* 4:579-82.
- Sweeten, J.M., and D.L. Reddell. 1978. Nonpoint Sources: State-of-art overview. *Transactions of the Am. Soc. Ag. Engineers.* 21(3):474-478.
- Tanner, G.W. and W.S. Terry 1991. *Soil and Crop Science Soc. of Florida. Proceedings, Vol. 50, p. 80- 84.*
- Thompson R.C.A., and J.A. Reynoldson. 1993. *Giardia and Giardiasis. Advances in Parasitology 32:72-99.*
- Walker, J.W. 1995. Viewpoint: grazing management and research now and in the next millennium. *J. Range Manage.* 48:350-357
- Whipp S.C., M.A. Rasmussen, and W.C. Cray, Jr. 1994. Animals as a source of *Escherichia coli* pathogenic for human beings. *J. Amer. Vet. Med. Assoc.* 204:1168-1175.
- Wilkinson, S. R. and R. W. Lorry 1973. Cycling of mineral nutrients in pasture ecosystems. In G. W. Butler and R. W. Bailey. *Chemistry and Biochemistry of herbage. Vol. Academic Press, New York. p. 247-315.*
- Woodmansee R. G., I. Allis, and J. J. Mott. 1981. Grassland Nitrogen. *Ecol. Bull* 33: 443-462.
- Xiao L. 1994. *Giardia* infection in farm animals. *Parasitology Today* 10:436-438.