

---

## **Using a 3-D Computer Simulation Model of Peach Tree Architectural Growth and Dry Matter Partitioning to Evaluate the Production Potential of Peach Trees on Size-Controlling Rootstocks**

---

**Project Leader: Dr. Theodore DeJong**

**Cooperators: Kevin Day, Romeo Favreau, Scott Johnson**

### **ABSTRACT**

The overall objective of this project is to develop a comprehensive functional-structural tree model of peach tree architectural development and growth, and carbon partitioning/source sink interactions within the plant. This model simulates tree and fruit growth and physiological responses to light distribution within the canopy, hourly temperature changes, user-imposed pruning and fruit thinning practices, and the use of different rootstocks.

We have developed sub-models of the architectural characteristics and bud fates on four categories of shoots growing on pruned trees growing on Nemaguard and HBOK 32 rootstocks growing at the Kearney Agricultural Research and Extension Center. These sub-models have been inserted into the overall P-PEACH model to provide architectural information concerning the differences between trees growing on standard vs. size-controlling rootstocks.

We have also completed the second aspect of the project that incorporated a sub-model to simulate water flow from the soil through the tree and the calculation of stem water potential at specific locations throughout the tree. Since we know that the size-controlling behavior of our new peach rootstocks is mediated by tree water relations this will enable mechanistic modeling of that size-controlling behavior. However, developing the water sub-model required us to reprogram the overall model to use hourly, instead of daily, time-steps. This required other reprogramming that allowed further mechanistic modeling of leaf, stem and fruit growth and functioning. This aspect of the research is ground-breaking and the majority of it is being funded by other sources.

As a corollary to this work we completed new studies of the xylem anatomy of size-controlling and standard rootstocks and estimated the root hydraulic conductance of all of the rootstock genotypes that are of primary interest in the rootstock project. These analyses showed that there are clear differences in the xylem anatomy of the size-controlling rootstocks that directly correspond with the amount of vigor expressed by the rootstock. In conjunction with the modeling work this anatomical research provides a clear mechanistic understanding of how dwarfing peach rootstocks work as well as how tree water stress influences tree growth.

## DISCUSSION/CONCLUSION

Crop growth and yield is dependent on a complex set of interactions involving the tree scion and rootstock genotype, the physiological and developmental processes that occur within the tree, the interaction of these processes with the environment that the plant grows in and responses to horticultural manipulation of the tree by the crop manager. Understanding crop growth and yield responses of trees are more complex than most crops because the effects of all these factors are carried out over multiple years. Most experimental research concerning factors that influence these complex processes and the interactions between them has been limited to dealing with one, two or at most three environmental and/or management factors at a time and then monitoring a limited set of plant responses at the tissue, organ, or whole plant level. While these experimental approaches have yielded substantial information about tree crop responses to specific factors, many times experiments have led to conflicting results and it has been very difficult to develop integrated understanding of crop growth and yield responses over multiple years in complex environments. Because of this lack of integrated understanding, research tends to be repeated in various forms over the years and true progress in some areas tends to stagnate until new experimental approaches are developed. Furthermore research tends to get concentrated on specific topics that are measurable with newly available equipment (like photosynthesis, stomatal conductance, water potential, etc.) while information on other important topics (like canopy development processes, canopy architecture, bud fates, carbohydrate storage, etc.) tends to be neglected.

At the same time, molecular level plant biologists and geneticists are eager to apply their new-found tools of genomics, proteomics and metabolomics to solve crop production problems but they have even less understanding of the complex factors and processes controlling or influencing crop growth and yield than the field biologists/pomologists. If these modern techniques of plant biology are ever to be successfully applied to solving complex crop production problems, a more complete understanding of the factors influencing plant growth processes, the complex interactions between them, and the environment, will be necessary. It will also be important to be able to predict outcomes of specific metabolic or developmental changes over several years.

Recent advances in computer technology have made it possible to develop functional-structural plant models that simultaneously simulate whole plant photosynthesis, tree architectural growth and carbon partitioning within the structure of the tree and simultaneously display tree structural development in three dimensions on a computer screen (Allen et al. 2005, 2007). The most advanced of these types of models is being developed to simulate peach tree growth and development and recent advances have successfully simulated responses to pruning and fruit thinning as well as environmental factors such as light and temperature (Lopez et al.2008).

The overall objective of this proposal is to develop a peach tree model that would adapt all of the features of the L-PEACH model to simulating peach tree growth and crop productivity on size-controlling vs. standard rootstocks. This project can be thought of as an attempt to build a working peach tree *in silico* by assembling all the pertinent physiological and developmental concepts, information and data required to make a peach tree functional into a unified, integrated model. It can be likened to trying to build a working car by studying a car and how it functions and then trying to build a working car by having a third of its parts, no manual and creating the missing parts by understanding the general behavior of how the car is supposed to work; and then assembling the car. This exercise forces one to

pay attention to all parts (not just the ones that appear most important or interesting at first glance, or those that are easy to measure) and develop integrated understanding of tree function. This process points out the most important things that we don't understand about trees but also provides a means for the evaluation of new information or data within the context of whole plant functioning as it becomes available. Previous work on this model led us to the discovery that peach fruit grow according to a relative growth rate function and the importance of early spring temperatures on predicting harvest date and fruit sizing potential. This information is now at the center of recommendations for fruit thinning. This modeling work has also led to greatly increased understanding of tree and fruit growth responses to pruning. This type of understanding is what will be necessary to develop new approaches to manage tree growth, with or without size-controlling rootstocks, and develop more labor efficient orchard management practices.


During the past year we continued to improve the general model by developing a more detailed version of the model that simulates water uptake and transport so that the water potential of every part of a simulated tree is calculated hourly and fluctuates based on time of day, light, temperature, and leaf transpiration. Subsequently we began improving the methods used to calculate carbohydrate transport within the model so that carbohydrate transport would be linked with water transport processes. This has been a very complex problem but we have successfully changed the model so that the modeled carbohydrate transport processes are more realistic. At the same time more realistic sub-models of leaf and stem growth and development were developed and incorporated in the model so that vegetative growth can be updated hourly and is linked to the daily patterns of temperature and water potential. We are currently in the process of refining and validating all of these changes to the model. In 2010 we will begin detailed field studies for leaf and stem growth and development in order to check or model and develop accurate mathematical functions that describe their growth.

In 2009 we also conducted anatomical and physiological studies of the xylem characteristics of size-controlling rootstocks compared to the standard Nemaguard rootstock. These analyses showed that there are clear differences in the xylem anatomy of the size-controlling rootstocks (Figures 1 and 2). The differences in xylem anatomy directly relate to reduced theoretical hydraulic conductance among the various rootstocks (Tombesi, et al. 2010) and thus, coupled with previous physiological studies (Basile et al. 2003 a and b; Solari et al., 2006 a, b, and c), provide anatomical and physiological basis for understanding the size-controller behavior of specific genotypes. The theoretical xylem hydraulic conductance of the specific rootstocks that we have been studying in the rootstock project will be used as inputs to the L-PEACH model to attempt to simulate the dwarfing effect of specific rootstocks of tree growth and development.


This is a very ambitious project that builds on nearly 20 years of modeling experience with peach trees. It will both test our current concepts of how environmental factors such as light and temperature as well as management factors such as pruning, fruit thinning, scion cultivar, rootstock, irrigation and nitrogen fertilization interact to influence tree growth and fruit yield and quality. In doing so, it will provide information about how to optimize management of orchards to meet grower needs.

## REFERENCES

- Allen, M. T., Prusinkiewicz, P., DeJong, T. M., *Using L-Systems for Modelling Source-Sink Interactions, Architecture and Physiology of Growing Trees: The L-PEACH Model*, *New Phytologist* **166**, 869-880, 2005
- Allen, M. T., Prusinkiewicz, P., Favreau, R.R., DeJong, T.M., *L-Peach, an L-System-Based Model for Simulating Architecture, Carbohydrate Source-Sink Interactions and Physiological Responses Of Growing Trees*, In *'Functional-Structural Plant Modeling in Crop Production'* (Eds. J Vos, LFM. de Visser, P.C. Struick, J. B., Evers) pp. 139-150. (Springer: The Netherlands) 2007
- Basile, B., Marsal, J. and DeJong, T. M., *Daily Shoot Extension Growth of Peach Trees Growing on Rootstocks that Reduce Scion Growth is Related to Daily Dynamics of Stem Water Potential*, *Tree Physiology* **23**:695-704, 2003
- Basile, B., Marsal, J., Solari, L. I., Tyree, M. T., Bryla D .R. and DeJong, T. M., *Hydraulic Conductance of Peach Trees Grafted on Rootstocks with Differing Size-Controlling Potential*, *Journal of Horticultural Science & Biotechnology* **78**(5):768-774, 2003
- Lopez, G. Favreau, R. R., Smith, C., Costes, E., Prusinkiewicz, P. and DeJong, T.M., *Integrating Simulation of Architectural Development and Source-Sink Behaviour of Peach Trees by Incorporating Markov Chain and Physiological Organ Function Sub-Models into L-PEACH*, *Functional Plant Biology* **35**:761-771, 2008
- Solari, L. I. and DeJong, T. M., *The Effect of Root Pressurization on Water Relations, Shoot Growth, and Leaf Gas Exchanges of Peach (Prunus Persica) Trees on Rootstocks with Differing Growth Potential and Hydraulic Conductance*, *J. Experimental Botany* **57**: 1981-1989, 2006
- Solari, L. I., Johnson S. and DeJong T. M., *Relationship of Water Status to Vegetative Growth and Leaf Gas Exchange of Peach (Prunus Persica) Trees on Different Rootstocks*, *Tree Physiology* **26**:1333-1341, 2006
- Solari, L.I., Johnson S. and DeJong, T. M., *T Hydraulic Conductance Characteristics of Peach (Prunus Persica) Trees on Different Rootstocks are Related to Biomass Production and Distribution*, *tree Physiology* **26**: 1343-1350, 2006
- Tombesi, S., Johnson R. S., Day K. R. and DeJong, T. M., *Relationships between Xylem Vessel Characteristics, Calculated Axial Hydraulic Conductance and Size-Controlling Capacity of Peach Rootstocks*, *Annals of Botany* in press, 2010



**Figure 1.** Weighted mean xylem vessel diameter and calculated theoretical axial xylem conductance of root, trunk and shoot xylem tissue from Nemaguard, P30-139 (Controller 9) and K146-43 (Controller 5) rootstocks.



**Figure 2.** Weighted mean xylem vessel diameter and calculated theoretical axial xylem conductance of root, trunk and shoot xylem tissue from HBOK50, HBOK 10, HBOK 32 and HBOK27 rootstocks.

This page left intentionally blank.