

BALSAM WOOLLY ADELGID IN CALIFORNIA

MICHAEL JONES PhD, UCCE FOREST ADVISOR – MENDOCINO, LAKE, & SONOMA

CHRIS LEE, CAL FIRE - FOREST HEALTH SPECIALIST

YANA VALACHOVIC, UCCE FOREST ADVISOR – HUMBOLDT & DEL NORTE


NORTH COAST SOD MEETING, MAY 2019



BALSAM WOOLLY ADELGID (BWA)


ADELGES PICEAE RATZEBURG (HEMIPTERA: ADELGIDAE)

- Invasive aphid-like insect
 - likely intro. from Europe on nursery stock
- First detected in eastern NA in 1900
 - balsam fir, Fraser fir
- Detected in western NA in 1928
 - grand fir, noble fir, European silver fir, Pacific silver fir, subalpine fir
- Hosts: All true firs (*Abies* spp.) in NA



Chris Lee, Cal Fire

NORTH AMERICAN DISTRIBUTION




INFESTATION RISK MAP





BWA CHARACTERISTICS

- Tiny wingless insect (0.04 in long)
- Piercing/sucking mouthparts
 - Inject toxic saliva that damages vascular tissues
- Mostly immobile
- Excretes waxy wool-like substance
- Infests stems and twigs (100-200 adelgids/in²)
- Disperse by wind and birds



BWA LIFE CYCLE IN NA

- Asexual repro. & no males (parthenogenetic)
- Single host (anholecyclic)
- Multiple generations



SYMPTOMS OF INFESTATION

- Swelling at bud and branch nodes (gouting)
- Stunted terminal growth
- Top curl
- Woolly masses on stem
- Irregular growth rings in stem (“rotholz”)
- Branch dieback
- Crown thinning



James McGraw, NC State

STAND AND LANDSCAPE IMPACTS

- Changes in forest structure
- Infestations are persistent
 - Can reduce reproductive potential of hosts
- Surviving infested trees susceptible to other pests/diseases
- Host tolerance
 - Subalpine, grand, and pacific silver fir very susceptible
 - noble, white, and European firs less susceptible; unless on site
 - More susceptible at low elevations and moist environments
- Site quality impacts susceptibility of hosts



UGA3227014

MANAGEMENT

- Challenge b/c of persistent infestations
 - Outbreaks can occur with favorable environ conditions
- Silvicultural practices
- Biological control
 - 1950-60's – 25 predators introduced (8 established)
 - Native and introduced predators appear ineffective
- Chemical control
 - Systemic injections of insecticides





Figure 4-2: Balsam woolly adelgid management: (A) In infested stands dominated by true fir, harvest damaged and high-risk species and convert to nonhost species that are adapted to the site. (B) In mixed-species stands, remove damaged and high-risk hosts.



CA INFESTATION

- 1928 - First detected in SF




CA INFESTATION

- 1928 - First detected in SF
- 2012 – extensive grand fir mortality observed near Fort Bragg




CA INFESTATION

- 1928 - First detected in SF
- ~2012 - grand fir mortality observed near Fort Bragg
- ~2017 – BWA detected near Fortuna



CA INFESTATION RESEARCH

- What is BWA doing in CA?
- Extent of the infestation
- Is the BWA infestation in coastal grand fir different
- BWA life cycle
- Coastal grand fir susceptibility
- Range expansion
- Management guidelines



THANK YOU

If you have questions:

mjones@ucanr.edu

<http://ucanr.edu/sites/forestry/>



UGA1515037