

RESEARCH ARTICLE

Open Access

Tomato root transcriptome response to a nitrogen-enriched soil patch

Daniel R Ruzicka¹, Felipe H Barrios-Masias², Natasha T Hausmann², Louise E Jackson² and Daniel P Schachtman^{*1}

Abstract

Background: Nitrogen (N), the primary limiting factor for plant growth and yield in agriculture, has a patchy distribution in soils due to fertilizer application or decomposing organic matter. Studies in solution culture oversimplify the complex soil environment where microbial competition and spatial and temporal heterogeneity challenge roots' ability to acquire adequate amounts of nutrients required for plant growth. In this study, various ammonium treatments (as ¹⁵N) were applied to a discrete volume of soil containing tomato (*Solanum lycopersicum*) roots to simulate encounters with a localized enriched patch of soil. Transcriptome analysis was used to identify genes differentially expressed in roots 53 hrs after treatment.

Results: The ammonium treatments resulted in significantly higher concentrations of both ammonium and nitrate in the patch soil. The plant roots and shoots exhibited increased levels of ¹⁵N over time, indicating a sustained response to the enriched environment. Root transcriptome analysis identified 585 genes differentially regulated 53 hrs after the treatments. Nitrogen metabolism and cell growth genes were induced by the high ammonium (65 μg NH₄⁺-N g⁻¹ soil), while stress response genes were repressed. The complex regulation of specific transporters following the ammonium pulse reflects a simultaneous and synergistic response to rapidly changing concentrations of both forms of inorganic N in the soil patch. Transcriptional analysis of the phosphate transporters demonstrates cross-talk between N and phosphate uptake pathways and suggests that roots increase phosphate uptake via the arbuscular mycorrhizal symbiosis in response to N.

Conclusion: This work enhances our understanding of root function by providing a snapshot of the response of the tomato root transcriptome to a pulse of ammonium in a complex soil environment. This response includes an important role for the mycorrhizal symbiosis in the utilization of an N patch.

Background

Nitrogen (N) is often a primary limiting factor for plant growth and yield in agriculture. Applications of N in conventional agriculture include fertilizer banding to the side of the plants, broadcasting on the surface of soil, and anhydrous ammonia injections. These N application methods as well as localized microbial turnover of organic N can result in spatial and temporal heterogeneity (patchiness) of soil N resulting in non-uniform availability to plant roots. Furthermore, the rapid immobilization and nitrification of N additions by soil

microbes can quickly alter N availability to the root [1-3]. Roots respond to localized nutrient patches by up-regulating ion transporters and by the proliferation of new roots into the patch to capture the additional N [4-6]. Mycorrhizal fungi provide plants with an additional mechanism to explore the soil and capture nutrients from enriched regions, increasing nutrient uptake potential [7,8].

Plant roots predominantly acquire N from the rhizosphere as inorganic ammonium (NH₄⁺) or nitrate (NO₃⁻), and subsequently assimilate intracellular NH₄⁺ into amino acids [9,10]. Roots sense and respond to changes in internal and external N status, which includes the regulation of gene expression, metabolism, and further N uptake and assimilation [11,12]. High and low affinity N

* Correspondence: dschachtman@danforthcenter.org

¹ Donald Danforth Plant Science Center, 975 N Warson Rd., St. Louis MO 63132 USA

Full list of author information is available at the end of the article

transport systems in roots allow plants to maximize uptake depending on soil N availability. High affinity transport systems are induced or activated under conditions where soil N availability is reduced (1 μM to 0.5 mM), while low affinity transport systems (active above ~ 0.5 mM N) may be constitutively expressed and transport N into the plant when soil N concentrations are high [13]. Members of the NH_4^+ transporter (AMT) gene family [14] transport NH_4^+ across the plasma membrane of root epidermal cells where it may be locally assimilated [15]. Intracellular NH_4^+ is assimilated into glutamine and glutamate via the N-regulated glutamine synthetase (GS) and glutamate synthase (GOGAT) enzymes [16]. The NO_3^- transporters (NRT) are also encoded by a large gene family [17]. The NO_3^- taken up by roots can be translocated to the shoot or reduced in roots to nitrite (NO_2^-) and then NH_4^+ via N-regulated NO_3^- reductase and NO_2^- reductase [13,16]. Both AMTs and NRTs exhibit complex gene regulation patterns in response to various forms and concentrations of N. These transporters are regulated by internal and external N and provide roots with a mechanism to mount a coordinated response that may serve to increase N acquisition and metabolism [11,18,19].

Recent studies have moved beyond examining expression changes of single genes or gene families to studying global changes in plant gene regulation by nutrients [19-22]. Microarray analyses of Arabidopsis and tomato roots subjected to increased NO_3^- identified hundreds of differentially regulated genes whose functions included N metabolism, cell growth, and transcription [19,21]. However, most genomics studies on plant nutrient metabolism have utilized hydroponic-grown non-mycorrhizal plants, potentially limiting their translatability to our understanding of roots response in soil where nutrients are not distributed uniformly and inorganic N is being transformed. In this report we used molecular tools to study root response to the application of known concentrations of NH_4^+ in a well defined region of the soil. Our aim was to characterize how roots respond to a nutrient patch in natural soils where complex ecological processes are occurring including mycorrhizal colonization and the microbial transformation of NH_4^+ to NO_3^- . This approach is in contrast to previous studies [19-22] that have used hydroponics to study responses to NH_4^+ or NO_3^- singly rather than a dynamic situation which is more relevant to agriculture or natural ecosystems where NH_4^+ is rapidly transformed to NO_3^- . We report on ^{15}N uptake and translocation and the coordinated changes in gene expression patterns in mycorrhizal roots following a localized pulse of NH_4^+ as NO_3^- gradually became more available.

Results

Soil N, plant status, and plant N uptake

Previous work showed that the field soil used for this study contained low concentrations of inorganic N, high soil N mineralization potential, moderate mycorrhizal colonization of tomato roots, and very few changes in the soil food web after nutrient addition [8]. In order to create a nutrient patch and recover roots that were directly exposed to the treatment, pots were prepared with a soil root in-growth core (ring) buried 5 cm below the soil surface and subsequently referred to as the patch (Figure 1a). A pulse of $^{15}\text{NH}_4^+$ was injected into the soil patch to simulate the short-term effects of soil inorganic N spatial heterogeneity. The experimental design consisted of the addition of a high NH_4^+ treatment (65 $\mu\text{g } ^{15}\text{NH}_4^+\text{-N g}^{-1}$ soil) 100-fold higher than ambient NH_4^+ levels, a low NH_4^+ treatment (6.5 $\mu\text{g } ^{15}\text{NH}_4^+\text{-N g}^{-1}$ soil) 10-fold higher than ambient NH_4^+ levels, and a water treatment to control for any potential mobilization of nutrients that occurs when soil moisture is increased.

The NH_4^+ treatments increased the soil inorganic N concentrations in the patch soil, simulating heterogeneous soil patches. In the high NH_4^+ treatment group, the highest soil NH_4^+ concentration (39.7 $\mu\text{g NH}_4^+\text{-N g}^{-1}$ dry soil) was measured at the time of the first sampling which was 5 hrs after injection. At 53 hrs after treatment it remained significantly higher than controls (25.3 $\mu\text{g NH}_4^+\text{-N g}^{-1}$ dry soil) (Fig 1b), and by 96 hrs, decreased to 3.5 $\mu\text{g NH}_4^+\text{-N g}^{-1}$ dry soil due to microbial transformations and/or plant uptake of the added NH_4^+ . In the high NH_4^+ treatment rings, soil NO_3^- concentrations were above ambient levels within 29 hours, indicating nitrification of NH_4^+ (Figure 1c). Patch soil NO_3^- concentrations increased over the first 53 hrs after injection of the high NH_4^+ treatment. In the low NH_4^+ treatment, soil NH_4^+ and NO_3^- concentrations were similar to the water controls.

Over the course of the experiment, plant shoot growth was unaffected by treatment ($P = 0.78$). Percent total N (mean \pm SE) in the shoots was $2.29\% \pm 0.08$ at the time of treatment and unaffected by the N treatments ($P = 0.85$). Percent phosphate (P) (mean \pm SE) in the shoots was $0.18\% \pm 0.002$ and was unaffected by the N treatments. Roots were significantly colonized by arbuscular mycorrhizal fungi as measured by microscopic counting ($33.5\% \pm 8.4$, mean \pm SE) and fungal transcript analysis (data not shown).

To test whether the NH_4^+ treatments resulted in measurable ^{15}N uptake and translocation, atom percent ^{15}N was assayed in roots and leaves (Figure 1d and 1e).




Figure 1 Ammonium treatments altered soil nitrogen dynamics and plant N uptake. (a) Diagram of pot-grown tomato plants where water, low NH_4^+ ($6.5 \mu\text{g NH}_4^+\text{-N g}^{-1}$ soil), and high NH_4^+ ($65 \mu\text{g NH}_4^+\text{-N g}^{-1}$ soil) treatments were injected into buried soil rings. (b) and (c) $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$ concentrations per gram of patch soil in the three treatment groups at 5, 29, 53, and 96 hrs after injection. (d) and (e) atom percent ^{15}N in roots and shoots 5, 29, 53, and 96 hrs after injection of the three treatment groups with ^{15}N labeled NH_4^+ fertilizer. Data represent the mean \pm SEM of 3 biological replicates. Within each graph, means followed by different letters are significantly different from one another at $P < 0.05$ (two-way ANOVA with Tukey-Kramer HSD test).

Within 29 hrs, patch roots and leaves from high NH_4^+ plants contained increased amounts of ^{15}N compared to naturally occurring ^{15}N levels in the water control. The amount of ^{15}N in these tissues continued to increase over time. In the patch roots from the low NH_4^+ treatment, atom % ^{15}N was not higher compared to the water controls until 53 hrs after injection, and there was no significant enrichment detected in leaves of the low NH_4^+ treatments at any time point. There were significantly higher concentrations of ^{15}N in the high NH_4^+ treatment roots and leaves compared to the low NH_4^+ treatment samples at multiple time points, further confirming a physiological difference between these treatments.

Microarray analysis of patch root transcription

Affymetrix Tomato GeneChips were used to analyze the root transcriptome at 53 hrs post-treatment; when ^{15}N enrichment levels were detected in roots from both high and low NH_4^+ treatment groups. Array analysis detected expression of 5822 of the 9524 transcripts contained on the tomato genechip. Statistical analyses identified 585 genes that were significantly altered in expression among the three treatment groups (Additional file 1). The high

NH_4^+ treatment resulted in a much larger transcriptome response than the low NH_4^+ treatment, with 535 genes differentially expressed between the high NH_4^+ treatment and water control, compared to 89 genes with different expression levels between low NH_4^+ treatment and water control (Figure 2a). While there were many differences between the regulated genes under high or low NH_4^+ vs. water control treatments, 39 genes were identified as differentially regulated in both comparison groups, and all 39 were similarly regulated by the high and low NH_4^+ treatments compared to water (Figure 2a and Additional file 2).

We annotated and categorized nearly 80% of the 585 differentially regulated genes into putative functional classes. The NH_4^+ treatments resulted in the differential regulation of genes in a wide range of functional categories (Figure 2b). In every functional category, the high NH_4^+ vs. water comparison contained a significantly higher number of genes compared to the low NH_4^+ vs. water comparison, further indicating that the high NH_4^+ treatment resulted in a larger scale transcriptional response. Fisher's exact test was used to determine





Figure 2 Classification of nitrogen treatment-regulated genes into defined functional categories. (a) Venn diagram displaying the number of genes identified in either or both the high NH₄⁺ vs. water and low NH₄⁺ vs. water comparisons. (b) Affymetrix probeset sequences were matched to publicly available Genbank accession identifications and categorized according to tomato or Arabidopsis orthologue gene annotations. Differential expression was analyzed at 53h post treatment. Black bars represent the number of genes more highly expressed in the NH₄⁺ treatment compared to the water treatment, and white bars represent the number more highly expressed in the water treatment compared to the NH₄⁺ treatment. Asterisks indicate a significant difference in the proportion of genes in those functional categories between the high NH₄⁺ vs. water and low NH₄⁺ vs. water comparisons.

whether the gene lists from the [high NH₄⁺ vs. water] or [low NH₄⁺ vs. water] comparisons were enriched for different functional categories compared to one another, indicating a unique type of response to the two NH₄⁺ treatments. Significantly more cell growth and division genes were identified in the high NH₄⁺ vs. water comparison (29 out of 535, 5.4%) than the low NH₄⁺ vs. water comparison (0 out of 89, 0%) (P = 0.0245). Conversely, significantly more signal transduction genes were identified in the low NH₄⁺ vs. water comparison (15 out of 89,

16.9%) than the high NH₄⁺ vs. water comparison (45 out of 535, 8.4%) (P = 0.0187).

A significant proportion (39 out of 89, 43.8%) of the genes in the low NH₄⁺ vs. water comparison were similarly regulated in the high NH₄⁺ vs. water comparison (Additional file 2), and included N assimilation and metabolism genes such as glutamine synthetase and tryptophan synthase. Eight of the remaining 50 (16%) genes uniquely identified only in the low NH₄⁺ vs. water comparison function in sugar metabolism including glucosyl

hydrolase, trehalose 6-phosphate synthase, low affinity sucrose transporter *SUT4*, isocitrate lyase, and glucose 6-phosphate translocator.

Identification of genes regulated among the N treatments

Multiple genes in the N transporters, N metabolism, and amino acid metabolism subcategories were highly responsive to the NH_4^+ treatments. The high NH_4^+ treatment resulted in the increased expression level of N transporter, assimilation, and metabolism genes in roots including NH_4^+ transporter *AMT2*, NO_3^- transporter *NRT2.1*, nitrite reductase *Nir2*, glutamine synthetases *GS* (chloroplastic) and *GTS1* (cytosolic), and NADH-dependant glutamate synthase *GLT1* (*GOGAT*) (Table 1). Additional genes including peptide transporter 1 (*LeOPT1*), NH_4^+ transporter 1 (*AMT1*), and nitrilase (*NIT4*) exhibited lower expression levels in the high NH_4^+ treatment roots compared to water control samples. The nitrate transporters *NRT2.3* and *NRT3.2* were found on the array but were not differentially regulated among the treatment groups. Multiple amino acid metabolism genes were regulated by the N treatments including higher expression levels of two aspartate aminotransferases, an alanine aminotransferase, and a tryptophan synthase (Table 1). Lower expression levels of chorismate synthase 2 and alpha-aminoacidic semialdehyde synthase were detected (Table 1). A tomato MADS-Box transcription factor similar to the Arabidopsis N-starvation response transcription factor ANR1 was expressed 3.26-fold higher in the water control treatment compared to high NH_4^+ samples (Table 1).

We used qRT-PCR to quantify the expression of key N metabolism genes to confirm the accuracy of the array results. The qRT-PCR results largely agreed with both the direction and magnitude of expression levels across the three treatments, including three genes that were not significantly different by array or qRT-PCR (glutamate dehydrogenase *GDH1*, ferredoxin-dependant glutamate synthase *GLS1.2* (*GOGAT*), and asparagine synthetase *ASN1*) (Figure 3).

Microarrays identified genes that were differentially regulated among the NH_4^+ treatments from other functional categories including cell growth/division, cell wall biosynthesis, and stress/defense response, and sulfur metabolism (Figure 2b, Tables 2, 3, and 4, and Additional file 1). In the cell growth and division category, there were significantly more genes induced by the high NH_4^+ treatment vs. water control than repressed by the high NH_4^+ treatment (27 out of 29 cell growth and division genes induced by high NH_4^+ , $P < 0.001$) (Table 2). This category of genes included multiple cyclins, histones, and other growth factors. A similar pattern was observed for cell wall biosynthesis genes encoding multiple pectinest-

erases, expansins, endo-xyloglucan transferases, and cellulose synthase (18 out of 20 induced by high NH_4^+ , $P < 0.001$) (Table 3).

The NH_4^+ treatments caused various transcriptional responses in multiple stress and defense-response genes, and genes in this functional category were some of the most differentially regulated genes of the experiment (Table 4). Genes in certain stress and defense response subcategories exhibited similar transcriptional responses including glutathione metabolism genes (6 out of 7 higher expression in water samples), pathogen response factors (5 out of 6 higher expression in water samples), and heat shock proteins (6 out of 7 higher expression in high NH_4^+ samples).

Phosphate transporters

The patch roots in this study were colonized by arbuscular mycorrhizal (AM) fungi therefore we examined the transcriptome data to determine whether the NH_4^+ treatments might affect known symbiosis processes including the transcriptional regulation of the phosphate transporters (PTs). The tomato PTs group into mycorrhiza-specific, mycorrhiza-induced, and nonspecific root expression patterns, and are indicators of Pi status and the mycorrhizal symbiosis [23-26]. The Affymetrix tomato genechip only includes nonspecific phosphate transporter 1 (*PT1*), and it was 1.6-fold higher in the water samples compared to the low N treatment ($P = 0.004$). To test whether the NH_4^+ treatments resulted in the differential regulation of the other PT family genes, we assayed the expression of the tomato phosphate transporters *PT1*, 2, 3, 4, and 5 with qRT-PCR (Figure 4). Expression of nonspecific *PT1* and *PT2* were 1.9- and 3.1-fold higher in the water samples compared to the low N treatment samples ($P = 0.003$ and 0.046 , respectively). Mycorrhiza-induced *PT3* and mycorrhiza-specific *PT4* expression levels were 5.0- and 5.7- fold higher, respectively, in the low NH_4^+ treatment group compared to the water control samples ($P = 0.014$ and 0.045 , respectively). Mycorrhiza-induced *PT5* expression was not different among the treatments. The expression level of the phosphate starvation-induced tomato gene *TPS11* was 7.5-fold higher in the water samples compared to the low NH_4^+ treatment ($P = 0.019$). Similar trends were found for the high NH_4^+ treatments compared to water controls. We analyzed shoot Pi levels to test whether the alterations in *PT* gene expression correlated with or resulted in changes in shoot total Pi levels, but they not significantly different across the treatments.

Discussion

Previous studies have reported the transcriptional regulation of genes in diverse functional groups including

Table 1: Differentially regulated nitrogen assimilation and metabolism genes.

Probe Set ID	Putative Annotation	Fold Change (High N vs. water)	P-value	Fold Change (Low N vs. water)	P-value
Les.224.1.S1_at	glutamine synthetase	16.69	0.002	6.31	0.024
Les.2360.1.S1_at	nitrite reductase	4.25	0.044	1.39	0.364
Les.3640.1.S1_at	ammonium transporter	4.07	0.045	1.24	0.638
Les.2884.1.S1_at	glutamine synthetase	3.23	0.044	1.39	0.365
Les.28.2.S1_a_at	nitrate transporter 2.1	2.61	0.082	1.42	0.402
Les.987.1.A1_at	aspartate aminotransferase	2.29	0.050	-1.02	0.943
Les.987.3.S1_at	aspartate aminotransferase	2.18	0.044	-1.34	0.216
Les.5163.1.S1_at	dicarboxylate transport	1.85	0.075	1.99	0.105
Les.899.1.S1_at	NADH-dependent glutamate synthase	1.76	0.064	-1.05	0.834
Les.3626.1.S1_at	alanine aminotransferase	1.71	0.092	1.36	0.246
Les.2756.1.A1_at	tryptophan synthase-related	1.53	0.095	2.01	0.078
Les.231.1.S1_at	O-acetyl(thiol)serine lyase	1.32	0.136	1.63	0.088
Les.3660.1.S1_at	chorismate synthase 2	-1.73	0.061	-1.36	0.180
Les.299.1.S1_at	peptide transporter 1	-1.75	0.093	-1.37	0.254
Les.797.1.S1_at	ammonium transporter 1	-1.80	0.065	-1.09	0.700
Les.3289.1.S1_at	g-aminobutyrate transaminase subunit precursor	-1.84	0.053	-1.76	0.105

Table 1: Differentially regulated nitrogen assimilation and metabolism genes. (Continued)

Les.7.1.S1_at	homogentisate 1,2-dioxygenase HGO	-1.98	0.092	-1.54	0.214
Les.1493.1.S1_at	nitrilase	-2.02	0.098	-1.65	0.192
LesAffx.3336.1.S1_at	cystathionine beta-synthase domain protein	-2.15	0.108	-3.60	0.078
Les.3071.1.S1_at	alpha- aminoadipic semialdehyde synthase	-2.27	0.067	-2.35	0.105
Les.5024.1.S1_at	ANR1-like MADS- box transcription factor	-3.26	0.067	-2.28	0.156

Probe Set ID; Affymetrix identifier for each microarray probeset. Putative Annotation; functional annotation based on tomato protein function or function of Arabidopsis orthologues identified with BLAST searches. Fold Change; linear fold changes (bold values significant at False Discovery Rate (FDR) adjusted P-value < 0.10). High N = added 65 μg $^{15}\text{NH}_4\text{-N}$ per gram soil, low N = added 6.5 μg $^{15}\text{NH}_4\text{-N}$ per gram soil. Probe Set IDs Les.987.1.A1_at and Les.987.3.S1_at represent the same genes.

metabolism, energy, cell growth, and transcription/translocation in response to N nutrition as NO_3^- or NH_4^+ [19-22]. However, these have utilized hydroponics systems that do not necessarily reflect the rhizosphere environment encountered by plant roots in agricultural and natural ecosystems. Roots grown in solution culture systems do not compete with soil microbes for nutrients, and the concentrations of nutrients in solution are more uniform both spatially and temporally. Fertilizer application and soil processes in conventional and organic farming result in the formation of heterogeneous soil nutrient patches [4], and plant utilization of N patches depends on roots rapidly sensing and response to the local enrichment of nutrients where they are in competition with soil microbial N assimilation and nitrification, leaching, and denitrification [2,3,27,28].

To better understand plant root response to a localized and dynamic inorganic N soil patch, we utilized an experimental design that simulated a more realistic patch environment. The buried ring created the spatial attributes of an N patch by ensuring that harvested roots were localized to the NH_4^+ treatment patch. Measurements of soil NH_4^+ and NO_3^- levels confirmed dynamic soil transformations of available N by 53 hrs when we sampled the roots for microarray analysis. After 96 hrs, we observed a trend of decreasing NH_4^+ and NO_3^- soil levels, indicating plant and microbial uptake of both NH_4^+ and NO_3^- [1,9]. Furthermore, estimates of the % recovery of applied ^{15}N

in shoots from high and low NH_4^+ treatment groups after 96 hrs (22% and 21%, respectively) support the assertion that roots faced significant competition for N in the soil environment.

The rapid N uptake observed in this study demonstrates the ability of tomato roots to quickly capture fertilizer in soil patches [29,30]. The roots that encountered the high NH_4^+ treatment took up and translocated significantly more ^{15}N from the patch than the low NH_4^+ treatment roots. This observation is in agreement with the larger transcriptional response to the high NH_4^+ treatment including multiple nitrogen transport, assimilation, and metabolism genes. However, the relatively low % recovery suggests that actual uptake and assimilation are only a small fraction of what was initially available despite the rapid root responses.

The genes coordinately regulated in both high and low treatments may represent a conserved physiological response to different ranges of N patch conditions. The co-regulated list of genes did not include the NH_4^+ or NO_3^- transporters *AMT1*, *AMT2*, or *NRT2.1* identified in the high NH_4^+ vs. water control comparison. We speculate that roots in the high NH_4^+ patch needed additional transporters to effectively capture the higher soil N levels, while constitutively expressed transporters were sufficient in the low NH_4^+ patch. Of equal interest, however, is the set of 50 genes regulated by the low NH_4^+ treatment




Figure 3 qRT-PCR analysis of key nitrogen metabolism genes. Expression levels of NH₄⁺ transporters AMT1 and AMT2, glutamine synthetases GTS1 and GS, NO₃⁻ transporter NRT2.1, nitrite reductase Nii, NADH-dependant glutamate synthase GLT1 (GOGAT), glutamate dehydrogenase GDH1, ferredoxin-dependant glutamate synthase GLS1.2 (GOGAT), and asparagine synthetase ASN1 in patch roots harvested 53 hrs after high (65 μg NH₄⁺-N g⁻¹ soil), low (6.5 μg NH₄⁺-N g⁻¹ soil), or water control treatments. Relative quantity was calculated using the ΔΔCT method with actin (LeACT) as the reference control, and the water control group normalized to 1. For a given gene, means followed by different letters are significantly different from one another at P < 0.05 (one way ANOVA).

that were not similarly regulated by the high NH₄⁺ treatment. This group of genes included a significant number of sugar metabolism genes for which we have no clear explanation as to the functional significance for the change in regulation. One possibility is that sugar metabolism may have been required for the production of root exudates, which can increase plant growth promoting rhizobacteria, or stimulate N cycling by microbial populations in the rhizosphere [31]. While the total number of genes regulated by the two treatments suggests that the high NH₄⁺ treatment caused a larger response, the responses were in part unique, and may reflect different strategies to effectively utilize the N patch. Increased N availability stimulates highly regulated root development and growth in order to efficiently scavenge and assimilate the additional soil N [32,33]. The induction of multiple histone gene family members such as histones H4, H3 and H2AX in the high NH₄⁺ treatment suggests an increase in DNA replication [34,35], while the increased expression of a mitotic spindle checkpoint gene, replicon protein A, multiple cyclin genes, and a putative cdc20 suggest an increase in cell division processes [36,37]. Furthermore, the induction of cell wall genes including

expansins, pectinesterases, and cellulose synthase suggests an increase in cell wall biosynthesis that would be required during root growth [38]. In Arabidopsis, multiple expansins and other cell wall modification enzymes were up-regulated by NO₃⁻ 3 hrs post-treatment [20], and gene expression profiling of maize roots in early response to a NO₃⁻ treatment identified multiple genes involved in cell growth and lateral root elongation including alpha-expansin, kinesin, and cellulose synthase [39]. These experiments imply that similar N developmental response mechanisms are conserved across maize, Arabidopsis, and tomato roots, and that the root response to an N nutrient patch includes coordinated root growth.

In this study, diverse stress response genes encoding heat-shock proteins, glutathione transferases, thioredoxin, pathogenesis-related proteins, and dehydration/desiccation responsive proteins were found to be differentially expressed among the NH₄⁺ treatment groups. Limiting nutrient conditions cause various stress-related responses including the up-regulation of reactive oxygen species metabolism [40]. Chronic N stress induces a range of plant stress responses which include the transcriptional regulation of numerous stress responsive

Table 2: Differentially regulated cell growth and division genes.

Probe Set ID	Putative Annotation	Fold Change (High N vs. water)	P-value
Les.5082.1.S1_at	cdc20 cell cycle regulator	3.46	0.044
Les.2170.1.A1_at	cyclin	2.89	0.062
Les.5789.1.S1_at	histone 3	2.74	0.045
Les.103.1.S1_at	cyclin A1	2.65	0.044
Les.3009.3.A1_at	histone HTA12	2.52	0.053
Les.3713.1.S1_at	B2-type cyclin dependent kinase	2.48	0.044
Les.677.1.S1_at	histone H2AX	2.33	0.053
Les.677.2.A1_at	histone H2AX	2.31	0.061
Les.5740.1.S1_at	replicon protein A	2.28	0.044
LesAffx.66157.1.S1_at	mitotic arrest deficient-like	2.21	0.068
LesAffx.19390.1.S1_at	cyclin	2.16	0.044
Les.3209.1.S1_at	histone H4 replacement isoform	2.11	0.044
Les.3090.1.S1_at	histone H3	2.08	0.044
Les.3555.1.S1_at	histone H2B-2	2.00	0.044
Les.4603.1.S1_at	histone H3	1.96	0.044
Les.5283.1.S1_at	minichromosome maintenance protein	1.83	0.084
Les.4439.1.S1_at	histone H3	1.80	0.053
Les.4978.1.S1_at	DNA-dependant ATPase	1.77	0.091
Les.4442.1.S1_s_at	histone H2B-1	1.73	0.044
LesAffx.57438.1.S1_at	nucleosome chromatin assembly factor	1.72	0.081
Les.3209.2.A1_at	histone H4 replacement isoform	1.72	0.062

Table 2: Differentially regulated cell growth and division genes. (Continued)

Les.4831.1.S1_at	nucleosome chromatin assembly factor	1.64	0.053
Les.4539.1.S1_a_at	histone H4	1.57	0.091
LesAffx.2226.2.A1_at	ribonucleotide reductase-like	1.57	0.091
Les.4564.1.S1_at	microtubule associated protein	1.54	0.088
Les.2989.1.S1_at	histone HTA7	1.51	0.044
Les.4539.2.S1_at	histone H4	1.50	0.091
Les.3009.2.S1_at	histone HTA12	1.48	0.066
Les.4940.1.S1_at	cyclin	-1.56	0.088
Les.3563.1.S1_at	ER auxin binding protein 1	-1.74	0.082

Probe Set ID; Affymetrix identifier for each microarray probeset. Putative Annotation; functional annotation based on tomato protein function or function of Arabidopsis orthologues identified with BLAST searches. Fold Change; linear fold changes (bold values significant at False Discovery Rate (FDR) adjusted P-value < 0.10). Probe Set IDs representing the same genes include (Les.3009.3.A1_at and Les.3009.2.S1_at), (Les.677.1.S1_at and Les.677.2.A1_at), (Les.3209.1.S1_at and Les.3209.2.A1_at) and (Les.4539.1.S1_a_at and Les.4539.2.S1_at).

genes. In Arabidopsis ~35% of the genes upregulated by a severe chronic N limitation stress were classified as response to abiotic stimulus, general stress, or oxidative stress [41]. Studies of N effects on the expression of stress response genes in rice also indicate that N limiting conditions cause the differential regulation of biotic and abiotic stress genes [42]. From their studies, Lian *et al.* postulate that this could be due to the perception of N limitation as a biotic or abiotic stress that requires a conserved set of regulated genes that play protective roles [42]. Our results suggest that the conditions in the water control samples may have initiated a stress response in roots due to an N limitation, and that the high and low NH₄⁺ treatments alleviated this coordinated stress response.

The complex regulation of specific NH₄⁺ and NO₃⁻ transporters following the NH₄⁺ pulse may reflect a simultaneous and synergistic response to both NH₄⁺ and NO₃⁻ in the soil patch. Both Arabidopsis *AtAMT1.1* and tomato *LeAMT1* exhibit increased transcript levels during N deprivation and are repressed by NO₃⁻ and NH₄⁺ [43-45]. The higher expression of *LeAMT1* under control conditions and its repression by high NH₄⁺ in the present study further support the idea that *LeAMT1* is a high affinity ammonium transporter whose expression is regu-

lated by N-limiting conditions to increase NH₄⁺ uptake. In two hydroponics studies, *LeAMT2* was induced by increased concentrations of NH₄⁺ over the course of 24 hrs but repressed by increased concentrations of NO₃⁻ after 24 hrs [19,44]. The higher expression of *LeAMT2* in response to the soil N patch 53 hrs after treatment in this current study suggests that the positive effects of NH₄⁺ may be stronger than the long-term repressive effects of NO₃⁻ exposure. Arabidopsis, barley and tomato NO₃⁻ transporters *AtNRT2.1*, *HvNRT2*, and *LeNRT2.1* were induced by NO₃⁻ in hydroponic culture, and NH₄⁺ repressed *HvNRT2* expression [13,19]. We report that *LeNRT2.1* was induced in the high NH₄⁺ treatment where we measured increased NO₃⁻ concentrations. The increased NO₃⁻ in the patch may have induced *LeNRT2.1*, although the effects of NH₄⁺ alone in the absence of NO₃⁻ on *LeNRT2.1* remain to be tested. The complex regulation of the NH₄⁺ and NO₃⁻ transporters in this study indicate that tomato roots are able to quickly sense and respond to changing concentrations of NH₄⁺ and NO₃⁻ simultaneously in a localized N patch, enhancing N uptake and utilization. Moreover, growth is known to increase with co-provision of NH₄⁺ and NO₃⁻ [22]. Recent

Table 3: Differentially regulated cell wall metabolism genes.

Probe Set ID	Putative Annotation	Fold Change (High N vs. water)	P-value	Fold Change (Low N vs. water)	P-value
Les.3273.1.S1_at	cell wall-plasma membrane linker	5.98	0.053	1.63	0.424
LesAffx.846.2.S1_at	pectinacylesterase	3.22	0.080	-1.44	0.458
Les.3733.1.S1_at	expansin	3.20	0.067	1.45	0.414
Les.3590.1.S1_at	endo-xyloglucan transferase	3.12	0.081	-1.33	0.565
LesAffx.4617.1.A1_at	pectinesterase	3.10	0.063	-1.41	0.417
Les.2316.1.S1_at	cellulose synthase isomer	3.01	0.067	1.88	0.193
Les.2189.1.S1_at	pectinesterase	2.51	0.070	-1.07	0.868
Les.218.3.S1_at	pectinesterase	2.07	0.094	-1.14	0.710
Les.1604.1.A1_at	cobra-like4 phytocheletin synthase	1.91	0.096	1.51	0.219
Les.369.1.S1_at	expansin	1.87	0.074	1.48	0.186
LesAffx.69659.1.S1_at	chitinase class IV	1.82	0.063	1.65	0.127
Les.5233.1.S1_at	pectinesterase	1.76	0.089	1.06	0.829
Les.218.1.S1_at	pectinesterase	1.73	0.096	1.27	0.362
Les.3523.1.S1_at	polygalacturonase	1.72	0.096	1.17	0.551
Les.4739.1.S1_at	UDP-glucose:protein transglucosylase	1.70	0.048	1.28	0.196
Les.109.1.S1_at	beta-galactosidase	1.61	0.065	-1.02	0.944
Les.4707.1.S1_at	pectate lyase	1.61	0.061	1.67	0.093
Les.2590.2.A1_at	endo-xyloglucan transferase A2-like	1.40	0.085	-1.05	0.743

Table 3: Differentially regulated cell wall metabolism genes. (Continued)

Les.4523.1.S1_at	xyloglucan endotransglucosylas e-hydrolase	-2.53	0.079	-1.58	0.265
Les.4652.1.S1_at	esterase/lipase/ thioesterase	-3.76	0.063	-1.78	0.271

Probe Set ID; Affymetrix identifier for each microarray probeset. Putative Annotation; functional annotation based on tomato protein function or function of Arabidopsis orthologues identified with BLAST searches. Fold Change; linear fold changes (bold values significant at False Discovery Rate (FDR) adjusted P-value < 0.10). Probe Set IDs Les.218.1.S1_at and Les.218.3.S1_at represent the same genes.

studies have also reported root responses to soil glutamate that may have been available to patch roots [46]. This study highlights the ability of plant roots to simultaneously regulate multiple transporters for uptake of both forms of inorganic N as part of a plastic response strategy to quickly exploit the N pulse.

Numerous transcription factors were identified in the microarray study that may function as key regulators of a secondary response to the N enrichment. The tomato MADS-box transcription factor BT013126 shares 67% amino acid sequence similarity with Arabidopsis ANR1 and is expressed 3.26-fold higher in the water control compared to the high NH_4^+ treatment (Table 1). Arabidopsis ANR1 is a key regulator of the developmental response to N in roots and is induced by N starvation and repressed by NO_3^- re-supply [47,48]. Prior to the N additions, the plants were most likely N-limited as the shoot N concentration was below sufficiency levels (mean = 1.94% for water control) [49]. The expression pattern of this tomato *ANR1-like* gene in N patch roots corresponds to what was found in Arabidopsis, suggesting that its functional role to regulate root development in response to N is conserved across species and in diverse root environments.

Root responses to macronutrients N, P, potassium (K), and sulfur (S) are interconnected and may be due to the increased availability of one causing an imbalance in another. Previous studies have shown N addition to increase the expression level of S metabolism genes [21,50], which could account for the changes in S metabolism genes reported here (Additional file 1). Alternatively, these genes may have been affected by sulfate in the NH_4^+ treatment, although soil S concentrations were likely sufficient for the plant. Cross talk between K and N has also been shown where K deficiency alters the transcriptional and post-transcriptional activity of various N uptake, assimilation, and metabolism genes including three nitrate transporters [51,52]. Nitrogen and phosphate metabolism have been shown to be closely linked where N uptake results in coordinated P uptake [53,54]. However, in response to a 3 hr nitrate pulse, phosphate

transporter expression levels in hydroponics-grown Arabidopsis did not change [20]. In soils, mineral availability and acquisition is additionally affected by the mycorrhizal symbiosis, and previous work has linked the up-regulation of the fungal phosphate transporter *GiPT* to the presence of N [55]. In this current study, multiple phosphate transporters were regulated by the NH_4^+ treatments, in contrast to the Arabidopsis findings [20]. We observed that mycorrhizal-induced *PT3* and mycorrhizal-specific *PT4* were more highly expressed when more NH_4^+ was present in the soil. The *PT3* and *PT4* expression patterns suggest that arbuscular mycorrhizal Pi uptake may be enhanced by NH_4^+ soil enrichment. Phosphate transporters *PT1* and *PT2* are found in both mycorrhizal and nonmycorrhizal root tissues, but are repressed in mycorrhizal roots [56-58]. The repression of *PT1* and *PT2* in the NH_4^+ treatments in the present study further supports the conclusion that the NH_4^+ treatments promoted the symbiotic Pi uptake pathway. In fact, Wang *et al.* reported that tomato *PT2* was induced by NO_3^- in hydroponic-grown non-mycorrhizal roots [19], and thus it appears that *PT2* regulation in the current study was in response to up-regulation of the mycorrhizal Pi uptake pathway rather than soil NO_3^- directly. We can speculate that the NH_4^+ soil enrichment induced root growth in the nutrient patch, resulting in a localized P deficiency that promoted the mycorrhizal Pi uptake pathway. Importantly, the lower expression level of phosphate-starvation induced *TPSII* in the low and high NH_4^+ treatment plants suggests that these roots were receiving more Pi than the water control samples [59], although this was not measured directly. This shift towards the mycorrhizal Pi uptake pathway may have resulted in increased Pi uptake, possibly as a mechanism to support N-induced growth. Our results detail a novel and complex interaction between inorganic N, the arbuscular mycorrhizal symbiosis, and the tomato phosphate transporter gene family, and suggest an important role for the symbiosis in the utilization of an N patch to increase P uptake and maintain N-induced growth.

Table 4: Differentially regulated stress and defense response genes.

Probe Set ID	Putative Annotation	Fold Change (High N vs. water)	P-value	Fold Change (Low N vs. water)	P-value
Les.2287.3.A1_at	TAS14 peptide dehydrin	-24.08	0.053	-40.75	0.078
Les.5957.1.S1_at	lactoylglutathione lyase	-6.87	0.111	-17.15	0.078
Les.293.1.S1_at	hydroxyacylglutathione hydrolase	-4.66	0.513	-4.44	0.078
Les.23.1.S1_at	glutathione S-transferase	-4.54	0.569	-3.07	0.078
Les.124.1.S1_at	glutathione transferase	-4.38	0.044	-2.54	0.107
Les.5100.1.S1_at	type I small heat shock protein	-3.85	0.044	-2.20	0.107
Les.4789.1.S1_at	pathogenesis-related protein	-3.47	0.062	-2.12	0.115
Les.5341.1.S1_at	pathogen responsive alpha-dioxygenase 2	-2.76	0.098	-1.85	0.131
Les.1645.1.A1_at	pathogenesis-related chitin-binding protein	-2.44	0.065	-1.74	0.133
Les.4004.1.S1_a_at	pathogenesis related PR5-like protein	-2.26	0.070	-1.59	0.143
Les.5098.1.S1_at	early responsive to dehydration 7-like	-2.20	0.064	-1.42	0.152
LesAffx.47187.1.S1_at	responsive to dehydration 22-like	-2.12	0.058	-1.34	0.165
Les.5103.1.S1_at	pathogenesis-related protein 1 like	-2.04	0.062	-1.33	0.168
Les.3151.1.S1_at	universal stress protein	-1.99	0.084	-1.33	0.168
Les.253.1.S1_at	pathogenesis related protein 1-like	-1.95	0.044	-1.29	0.229
Les.4910.1.S1_at	stress enhanced protein 2-like	-1.93	0.085	-1.26	0.232

Table 4: Differentially regulated stress and defense response genes. (Continued)

Les.1498.1.S1_at	dehydration responsive	-1.87	0.096	-1.25	0.289
Les.208.1.S1_at	glutathione S-transferase	-1.72	0.062	-1.22	0.340
Les.2657.1.S1_at	rare cold inducible protein-like	-1.70	0.095	-1.20	0.377
Les.3194.1.S1_at	universal stress protein	-1.62	0.088	-1.16	0.386
Les.5128.1.S1_at	responsive to dehydration 22-like	-1.61	0.044	-1.14	0.454
Les.3276.3.S1_at	monocysteinic thioredoxin	-1.56	0.093	-1.11	0.509
Les.252.1.S1_at	wound-responsive protein-related	-1.50	0.077	-1.11	0.524
Les.54.1.S1_at	sulfiredoxin	1.10	0.044	-1.10	0.537
Les.248.2.A1_at	glutathione S-transferase	1.14	0.095	-1.08	0.796
Les.384.1.A1_at	thaumatin-like pathogenesis-related PR-5 like protein	1.42	0.070	-1.07	0.850
Les.4307.1.S1_at	early responsive to dehydration 3-like	1.47	0.053	-1.05	0.870
Les.1394.1.A1_at	heat shock factor binding protein 1	1.82	0.063	1.22	0.185
Les.3593.1.S1_at	heat shock protein	2.13	0.044	1.45	0.204
Les.2409.1.S1_at	dnaJ related molecular chaperone	2.30	0.050	1.57	0.537
LesAffx.43379.1.S1_at	dnaJ homologue 3	2.71	0.052	1.64	0.623
LesAffx.71535.1.S1_at	heat shock protein	2.81	0.077	1.72	0.639
Les.641.1.S1_at	dnaJ heat shock protein	3.01	0.067	1.95	0.746
LesAffx.66226.2.S1_at	cold-regulated plasma membrane 1 protein	3.28	0.044	1.97	0.180

Table 4: Differentially regulated stress and defense response genes. (Continued)

Les.5158.1.S1_at	dehydration response element B1A	3.36	0.092	1.99	0.218
LesAffx.59336.1.S1_at	response to desiccation 26-like transcription factor	3.59	0.082	2.25	0.231

Probe Set ID; Affymetrix identifier for each microarray probeset. Putative Annotation; functional annotation based on tomato protein function or function of Arabidopsis orthologues identified with BLAST searches. Fold Change; linear fold changes (bold values significant at False Discovery Rate (FDR) adjusted P-value < 0.10).

Conclusions

Spatially discrete NH_4^+ is quickly transformed in the soil and taken up by plants, and the tomato root transcriptome reflects levels of N availability and transformations of N that occur in the soil. The dynamic regulation of both NH_4^+ and NO_3^- transporters in N-patch roots demonstrates that roots are able to simultaneously sense and respond to both forms of inorganic N, in ways that are likely to increase root competition with microbial immobilization, nitrification, and denitrification, and conserve N within cropping systems. The arbuscular mycorrhizal symbiosis may further increase the effective recovery of other nutrients such as P in an N patch. The strong and diverse transcriptional response to the soil N patch illustrates the utility of applying transcriptomic studies to plants growing in realistic soil environments and the key genes co-regulated under high and low N conditions in this study may serve as molecular tools for monitoring plant N status in agricultural sites for finer tuning of fertilizer application, soil microbial N processes, and ultimately, to develop more efficient agriculture methods.




Figure 4 qRT-PCR analysis of phosphate physiology genes. Expression levels of the tomato phosphate transporters and phosphate starvation response gene TPSII in patch roots harvested 53 hrs after high ($65 \mu\text{g NH}_4^+\text{-N g}^{-1}$ soil), low ($6.5 \mu\text{g NH}_4^+\text{-N g}^{-1}$ soil), or water control treatments. Relative quantity was calculated using the $\Delta\Delta\text{CT}$ method with actin (LeACT) as the reference control, and the water control group normalized to 1. For a given gene, means followed by different letters are significantly different from one another at $P < 0.05$ (one way ANOVA).

Methods

Soil and plant material

Seeds of *Solanum lycopersicum* L. Cv. 76R [60] were surface sterilized, germinated with mist irrigation and then watered with one-tenth strength Long Ashton's solution containing N as $(\text{NH}_4)_2\text{SO}_4$ (4 mM) and NaNO_3 (8 mM). Plants were maintained under day/night length of 16/8 hr in UC Davis glasshouses. Seven week old seedlings were transplanted into 12-L pots containing buried rings with field-collected soil (Zamora loam, a fine silty, mixed thermic, Mollic Haploxeralfs) collected on an organically managed farm (Jim and Deborah Durst Farming in Esparto, Yolo County, California) [8]. The buried soil root in-growth cores (rings) were 7.3 cm in diameter and 4.2 cm tall (total volume 176 cm^3) and were filled with 210 g of field soil to a final bulk density equal to that of the surrounding soil (1.2 g cm^{-3}). The broad ends of the ring were covered with 1 mm plastic mesh to easily allow roots to grow up and down into the ring. The soil was passed through a 1 cm sieve before packing into pots at a bulk density of 1.2 g cm^{-3} . Extractable inorganic N (mean \pm standard error) was $0.20 \pm 0.02 \mu\text{g NH}_4^+\text{-N g}^{-1}$ dry soil and $9.7 \pm 1.09 \mu\text{g NO}_3^-\text{-N g}^{-1}$ dry soil at the time the rings were prepared. An application of one tenth strength Long Ashton's solution was applied two weeks after transplanting. Soil moisture was maintained gravimetrically at 19% before and after treatment injection by weighing the pots daily, and watering to compensate for evapotranspiration water loss.

Experimental design

We applied the nutrient treatments five weeks after transplantation by injecting 9 aliquots of 2 mL solution inside the buried ring using a template placed on the soil surface to assure an even distribution of nutrients and minimal loss of the solution. We added $6.5 \mu\text{g}$ and $65 \mu\text{g }^{15}\text{NH}_4^+\text{-N}$ (99 atom percent) g^{-1} dry soil in the ring for the low and high NH_4^+ treatments, respectively ($1.35 \text{ mg }^{15}\text{NH}_4^+\text{-N}$ per ring and $13.5 \text{ mg }^{15}\text{NH}_4^+\text{-N}$ per ring, respectively). Water was used as a control. Each treatment consisted of three biological replicates, and plants were destructively harvested at 5, 29, 53, and 96 hrs after $^{15}\text{NH}_4^+\text{-N}$ addition,

for a total number of 36 plants in an unreplicated block design. The transcriptome analyses were performed on roots harvested at 53 hrs post-treatment.

Harvest and sample analysis

At harvest, shoots were severed at the soil surface, dried at 60°C, weighed, and ground to a fine powder for isotope analyses. Immediately following the harvest of the above-ground biomass, the root in-growth rings were carefully exposed, and the roots growing into and out of the ring were severed. Patch roots for transcriptome analysis were rinsed and immediately frozen in liquid nitrogen. A homogenous sub-sample of the patch soil and the surrounding pot soil was immediately removed for gravimetric water content and soil inorganic N concentrations. Soil NH_4^+ and NO_3^- were analyzed after KCl extractions and colorimetric determination using modifications of Miranda *et al.*, [61] and Foster *et al.*, [62] respectively. A small subsample of ring roots was scored for arbuscular mycorrhizal fungi at 200 \times [63]. The remaining patch roots were washed by wet sieving, dried at 60°C, weighed, and ground to a fine powder for isotope analyses. All dried plant material was analyzed for $\delta^{15}\text{N}$ on a PDZ Europa ANCA-GSL elemental analyzer and a PDZ Europa 20-20 isotope ratio mass spectrometer (Sercon Ltd., Cheshire, UK) at the UC Davis Stable Isotope Facility, USA. Background ^{15}N was calculated as the average atom percent ^{15}N in the water samples (mean atom percent $^{15}\text{N} \pm \text{SD}$ in shoots: 0.369% \pm 0.0003). Leaf total P was analyzed by microwave digestion with nitric acid/hydrogen peroxide [64] followed by atomic absorption spectrometry and inductively coupled plasma atomic emission spectrometry at the UC Davis Division of Agriculture and Natural Resource Laboratory.

Soil analysis

Soil nutrient and plant isotope data were analyzed with a two-way analysis of variance (ANOVA) with harvest time, nutrient addition, and block as fixed main effects. All two-way interactions were tested. The three way interaction was not tested because of insufficient degrees of freedom. Data was checked to assure that the ANOVA assumptions were met and was transformed as necessary. Tukey-Kramer Honestly Significant Difference test was used to determine differences between means at $P < 0.05$. All data were analyzed using R (R Core Development Team 2007).

RNA isolation

Root RNA samples were extracted using the RNeasy Plant Mini Kit (Qiagen Sciences, Germantown, MD, USA) following the manufactures guidelines plus a third wash step before elution. RNA concentrations and quality were assessed using the Agilent Nanodrop and the RNA

6000 Nano Assay (Bioanalyzer 2100, Agilent, Santa Clara CA). RNA samples had RNA integrity numbers (RIN) of at least 7.0. DNase digestion was performed on 20 ug total RNA using RQ1 RNase-free DNase (Promega, Madison WI). These RNA were used for both microarray analysis and cDNA synthesis for qRT-PCR analysis.

Microarray analysis

Transcriptome profiling of each 53h-post injection RNA sample was performed using the Tomato Genome Array Chip (Affymetrix, Santa Clara, CA, USA). RNA samples were prepared for microarray analysis using the MessageAmp Premier RNA Amplification Kit (Ambion, Foster City CA) with 200 ng total RNA as input. Fragmented cRNA samples were then sent to the University of Missouri's DNA Core Facility for Array hybridization and scanning. Each array's CEL file was summarized in Affymetrix Expression Console software using the MAS5 algorithm. The signal intensities were log transformed, and quality control analysis performed. This array data has been made available on the Gene Expression Omnibus (GEO; <http://www.ncbi.nlm.nih.gov/projects/geo/>) accession #GSE21020. The data were filtered to remove probesets whose log signal intensity was below 4.605 in all 9 arrays. For each probeset, which represents the combined expression data from all relevant probe pairs on the chip, the generalized linear model $Y_{ij} = \mu + T_i + \epsilon_{ij}$ was fit. In each ANOVA, Y_{ij} is the log normalized transcript level for the i^{th} treatment and the j^{th} replicate, μ is the overall mean expression for the probeset and T_i represents the i^{th} treatment (water, low nutrient, and high nutrient). The null hypothesis $t_1 = t_0$ (i.e., mean expression not different between a pair of treatments) was tested using an F-test. We examined the model for conformation to the assumption of normality of the residuals testing the null hypothesis that the residuals for each gene were normally distributed using the Shapiro-Wilk Test. All analyses were performed in JMP Genomics 3.0 (SAS Institute, Cary NC). An FDR level of 10% was used for declaring findings significant, and a stringent rate of 5% was also examined [65,66]. ANOVA analysis of all differentially expressed genes can be found in Additional file 1. The list of statistically significant transcripts was initially annotated using the Affymetrix NetAffyx annotation file to match a representative Genbank public ID and Unigene to each probeset. The functional annotations of the Unigenes were grouped into functional categories as described in Bevan *et al.* [67]. In cases where the tomato gene did not have a matching Unigene or had not been functionally annotated, the tomato sequence was used to identify Arabidopsis orthologues by WU-BLAST searches at TAIR [68]. To test for enrichment of specific functional categories between the high vs water and low vs. water pairwise comparisons, Fisher's exact test was performed using

GraphPad Prism 5.0, San Diego CA). Within specific categories, binomial distribution probability tests were performed to test for enrichment in the up/down regulation patterns of functionally related genes (expectation = 0.5).

Quantitative real-time RT-PCR

cDNA was synthesized from 1.5 ug DNase-treated total RNA using the Superscript III kit (Invitrogen Carlsbad CA). Gene-specific primer sets were designed using IDT's primerquest software program (Additional file 3), and their sequence uniqueness confirmed with a nucleotide BLAST search against the tomato genome database. Primer pairs were tested for specificity and efficiency with serial dilution reactions and dissociation curve analysis post-amplification. Real-time PCR reactions were run on the Stratagene MX3000 PCR machine using Sybr Green chemistry (Invitrogen Platinum Sybr Green II master mix, 400 nM primer concentration, ROX reference dye, and 1:150 diluted cDNA). Multiple reference control genes were tested against all samples to identify control genes whose expression was not affected by the NH_4^+ treatments. *LeACT* and *LeUBI* were similarly expressed across the samples while *LeTubulin* was differentially expressed. *LeACT* was subsequently used as the reference control gene, and the relative expression of the various target genes was analyzed according to the $\Delta\Delta\text{CT}$ method [69]. Standard error was computed from the average of the ΔCT values for each biological sample [70].

Additional material

Additional file 1 Microarray analysis of differentially expressed genes across all treatments including the Affymetrix probeset ID, Genbank public ID, linear fold changes, FDR P-values, gene annotation information, and probeset redundancy information.

Additional file 2 Genes similarly regulated by high NH_4^+ and low NH_4^+ treatments compared to the water control.

Additional file 3 Primer sequences for genes assayed with qRT-PCR.

Authors' contributions

DRR carried out the microarray and molecular genetic studies, participated in the statistical analyses, and drafted the manuscript. FB-M and NTH participated in the greenhouse experiments and nutrient analyses, and helped write the manuscript. LEJ and DPS designed and coordinated the study, participated in the greenhouse experiments and data analyses, and helped write the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank Lauren McIntyre for her insights and contributions to the microarray analysis, Julie Chou, Kimberly Jacobs, Julien Linares, Walter Lopez, and Annie Young-Matthews for their help in the execution of the experiment and subsequent sample processing, the University of Missouri DNA Core Facility for hybridization and scanning of the Affymetrix GeneChips, and the National Science Foundation for funding support (#0723775 to DPS and LEJ).

Author Details

¹Donald Danforth Plant Science Center, 975 N Warson Rd., St. Louis MO 63132 USA and ²Department of Land, Air, and Water Resources, University of California, 1 Shields Avenue, Davis CA 95616 USA

Received: 3 February 2010 Accepted: 27 April 2010
Published: 27 April 2010

References

1. Burger M, Jackson LE: Microbial immobilization of ammonium and nitrate in relation to ammonification and nitrification rates in organic and conventional cropping systems. *Soil Biology and Biochemistry* 2003, **35**:29-36.
2. Booth MS, Stark JM, Rastetter E: Controls on nitrogen cycling in terrestrial ecosystems: A synthetic analysis of literature data. *Ecological Monographs* 2005, **75**:139-157.
3. Schimel JP, Bennett J: Nitrogen mineralization: Challenges of a changing paradigm. *Ecology* 2004, **85**:591-602.
4. Hodge A: The plastic plant: root responses to heterogeneous supplies of nutrients. *New Phytologist* 2004, **162**:9-24.
5. Granato TC, Raper CD: Proliferation of maize (*Zea mays* L) roots in response to localized supply of nitrate. *Journal of Experimental Botany* 1989, **40**:263-75.
6. Garnett T, Conn V, Kaiser BN: Root based approaches to improving nitrogen use efficiency in plants. *Plant, Cell and Environment* 2009, **32**:1272-1283.
7. Tibbett M: Roots foraging and the exploration of soil nutrient patches: the role of the mycorrhizal symbiosis. *Functional Ecology* 2000, **14**:397-99.
8. Cavagnaro TR, Jackson LE, Six J, Ferris H, Goyal S, Asami D, Scow KM: Arbuscular mycorrhizas, microbial communities, nutrient availability, and soil aggregates in organic tomato production. *Plant and Soil* 2006, **282**:209-225.
9. Lea PJ, Azevedo RA: Nitrogen use efficiency. 1. Uptake of nitrogen from the soil. *Annals of Applied Biology* 2006, **149**:243-247.
10. Lea PJ, Azevedo RA: Nitrogen use efficiency. 2. Amino acid metabolism. *Annals of Applied Biology* 2007, **151**:269-275.
11. Glass ADM, Britto DT, Kaiser BN, Kinghorn JR, Kronzucker HJ, Kumar A, Okamoto M, Rawat S, Siddiqi MY, Unkles SE, et al.: The regulation of nitrate and ammonium transport systems in plants. *Journal of Experimental Botany* 2002, **53**:855-864.
12. Loqué D, von Wirén N: Regulatory levels for the transport of ammonium in plant roots. *Journal of Experimental Botany* 2004, **55**:1293-1305.
13. Glass ADM: Nitrate uptake by plant roots. *Botany* 2009, **87**:659-667.
14. Ludewig U, Neuhäuser B, Dynowski M: Molecular mechanisms of ammonium transport and accumulation in plants. *FEBS Letters* 2007, **581**:2301-2308.
15. Howitt SM, Udvardi MK: Structure, function and regulation of ammonium transporters in plants. *Biochimica Et Biophysica Acta* 2000, **1465**:152-170.
16. Miller AJ, Cramer MD: Root nitrogen acquisition and assimilation. *Plant and Soil* 2005, **274**:1-36.
17. Tsay YF, Chiu CC, Tsai CB, Ho CH, Hsu PK: Nitrate transporters and peptide transporters. *FEBS Letters* 2007, **581**:2290-2300.
18. Loqué D, von Wirén N: Regulatory levels for the transport of ammonium in plant roots. *Journal of Experimental Botany* 2004, **55**:1293-1305.
19. Wang YH, Garvin DF, Kochian LV: Nitrate-induced genes in tomato roots. Array analysis reveals novel genes that may play a role in nitrogen nutrition. *Plant Physiology* 2001, **127**:345-359.
20. Scheible WR, Morcuende R, Czechowski T, Fritz C, Osuna D, Palacios-Rojas N, Schindelasch D, Thimm O, Udvardi MK, Stitt M: Genome-wide reprogramming of primary and secondary metabolism, protein synthesis, cellular growth processes, and the regulatory infrastructure of *Arabidopsis* in response to nitrogen. *Plant Physiology* 2004, **136**:2483-2499.
21. Wang YH, Okamoto M, Xing X, Crawford NM: Microarray analysis of the nitrate response in *Arabidopsis* roots and shoots reveals over 1000 rapidly responding genes and new linkages to glucose, trehalose-6-phosphate, iron, and sulfate metabolism. *Plant Physiology* 2003, **132**:556-567.
22. Lopes MS, Araus JL: Comparative genomic and physiological analysis of nutrient response to NH_4^+ , $\text{NH}_4^+:\text{NO}_3^-$ and NO_3^- in barley seedlings. *Physiologia Plantarum* 2008, **134**:134-150.

23. Nagy R, Karandashov V, Chague V, Kalinkevich K, Tamasloukht M, Xu G, Jakobsen I, Levy AA, Amrhein N, Bucher M: **The characterization of novel mycorrhiza-specific phosphate transporters from *Lycopersicon esculentum* and *Solanum tuberosum* uncovers functional redundancy in symbiotic phosphate transport in solanaceous species.** *Plant Journal* 2005, **42**:236-250.
24. Xu GH, Chague V, Melamed-Bessudo C, Kapulnik Y, Jain A, Raghothama KG, Levy AA, Silber A: **Functional characterization of LePT4: A phosphate transporter in tomato with mycorrhiza-enhanced expression.** *Journal of Experimental Botany* 2007, **58**:2491-2501.
25. Nagy R, Drissner D, Amrhein N, Jakobsen I, Bucher M: **Mycorrhizal phosphate uptake pathway in tomato is phosphorus-repressible and transcriptionally regulated.** *New Phytologist* 2009, **181**:950-959.
26. Javot H, Pumplin N, Harrison MJ: **Phosphate in the arbuscular mycorrhizal symbiosis: transport properties and regulator roles.** *Plant Cell and Environment* 2007, **30**:310-322.
27. Burger M, Jackson LE, Lundquist EJ, Louie DT, Miller RL, Rolston DE, Scow KM: **Microbial responses and nitrous oxide emissions during wetting and drying of organically and conventionally managed soil under tomatoes.** *Biology and Fertility of Soils* 2005, **42**:109-118.
28. Fageria NK, Baligar VC: **Enhancing nitrogen use efficiency in crop plants.** *Advances in Agronomy* 2005, **88**:97-185.
29. Hutchings MJ, de Kroon H: **Foraging in plants: The role of morphological plasticity in resource acquisition.** *Advances in Ecological Research* 1994, **25**:159-238.
30. Hodge A, Robinson D, Griffiths BS, Fitter AH: **Why plants bother: Root proliferation results in increased nitrogen capture from an organic patch when two grasses compete.** *Plant Cell and Environment* 1999, **22**:811-820.
31. Richardson AE, Barea JM, McNeill AM, Prignet-Combaret C: **Acquisition of phosphorus and nitrogen in the rhizosphere and plant growth promotion by microorganisms.** *Plant Soil* 2009, **321**:305-339.
32. Jovanovic M, Lefebvre V, Laporte P, Gonzalez-Rizzo S, Lelandais-Brière C, Frugier F, Hartmann C, Crespi M: **How the environment regulates root architecture in dicots.** *Advances in Botanical Research* 2007, **46**:35-74.
33. Vidal EA, Gutiérrez RA: **A systems view of nitrogen nutrient and metabolite responses in Arabidopsis.** *Current Opinion in Plant Biology* 2008, **11**:521-529.
34. Meshi T, Taoka KI, Iwabuchi M: **Regulation of histone gene expression during the cell cycle.** *Plant Molecular Biology* 2000, **43**:643-657.
35. Shaw P, Dolan L: **Chromatin and Arabidopsis root development.** *Seminars in Cell and Developmental Biology* 2008, **19**:580-585.
36. Menges M, Hennig L, Gruissem W, Murray JAH: **Cell cycle-regulated gene expression in Arabidopsis.** *Journal of Biological Chemistry* 2002, **277**:41987-42002.
37. Inze D, De Veylder L: **Cell cycle regulation in plant development.** *Annual Review of Genetics* 2006, **40**:77-105.
38. Zhong R, Ye ZH: **Regulation of cell wall biosynthesis.** *Current Opinion in Plant Biology* 2007, **10**:564-572.
39. Liu J, Han L, Chen F, Bao J, Zhang F, Mi G: **Microarray analysis reveals early responsive genes possibly involved in localized nitrate stimulation of lateral root development in maize (*Zea mays* L.).** *Plant Science* 2008, **175**:272-282.
40. Schachtman DP, Shin R: **Nutrient sensing and signaling: NPKS.** *Annual Review of Plant Biology* 2007, **58**:47-69.
41. Bi YM, Wang RL, Zhu T, Rothstein SJ: **Global transcription profiling reveals differential responses to chronic nitrogen stress and putative nitrogen regulatory components in Arabidopsis.** *BMC Genomics* 2007, **8**:281.
42. Lian X, Wang S, Zhang J, Feng Q, Zhang L, Fan D, Li X, Yuan D, Han B, Zhang Q: **Expression profiles of 10,422 genes at early stage of low nitrogen stress in rice assayed using a cDNA microarray.** *Plant Molecular Biology* 2006, **60**:617-631.
43. Lauter FR, Ninnemann O, Bucher M, Riesmeier JW, Frommer WB: **Preferential expression of an ammonium transporter and of two putative nitrate transporters in root hairs of tomato.** *PNAS* 1996, **93**:8139-8144.
44. von Wirén N, Lauter FR, Ninnemann O, Gillissen B, Walch-Liu P, Engels C, Jost W, Frommer WB: **Differential regulation of three functional ammonium transporter genes by nitrogen in root hairs and by light in leaves of tomato.** *The Plant Journal* 2000, **21**:167-175.
45. Loqué D, Yuan L, Kojima S, Gojon A, Wirth J, Gazzarrini S, Ishiyama K, Takahashi H, von Wirén N: **Additive contribution of AMT1;1 and AMT1;3 to high-affinity ammonium uptake across the plasma membrane of nitrogen-deficient Arabidopsis roots.** *Plant Journal* 2006, **48**:522-534.
46. Forde BG, Walch-Liu P: **Nitrate and glutamate as environmental cues for behavioural responses in plant roots.** *Plant, Cell and Environment* 2009, **32**:682-693.
47. Zhang H, Forde BG: **An Arabidopsis MADS box gene that controls nutrient-induced changes in root architecture.** *Science* 1998, **279**:407-409.
48. Gan Y, Filleur S, Rahman A, Gotensparre S, Forde BG: **Nutritional regulation of ANR1 and other root-expressed MADS-box genes in Arabidopsis thaliana.** *Planta* 2005, **22**:730-742.
49. Maynard DN, Hochmuth GJ: *Handbook for Vegetable Growers* New York: John Wiley and Sons, Inc; 1997.
50. Zhu GH, Zhuang CX, Wang YQ, Jiang LR, Peng XX: **Differential expression of rice genes under different nitrogen forms and their relationship with sulfur metabolism.** *Journal of Integrative Plant Biology* 2006, **48**:1177-1184.
51. Armengaud P, Breitling R, Amtmann A: **The potassium-dependant transcriptome of Arabidopsis reveals a prominent role of jasmonic acid in nutrient signaling.** *Plant Physiology* 2004, **136**:2556-2576.
52. Armengaud P, Sulpice R, Miller AJ, Stitt M, Amtmann A, Gibon Y: **Multilevel analysis of primary metabolism provides new insights into the role of potassium nutrition for glycolysis and nitrogen assimilation in Arabidopsis roots.** *Plant Physiology* 2009, **150**:772-785.
53. Gniazdowska A, Krawczak A, Mikulska M, Rychter AM: **Low phosphate nutrition alters bean plants' ability to assimilate and translocate nitrate.** *Journal of Plant Nutrition* 1999, **22**:551-563.
54. de Groot CC, Marcelis LFM, van den Boogaard R, Kaiser WM, Lambers H: **Interaction of nitrogen and phosphorus nutrition in determining growth.** *Plant and Soil* 2003, **248**:257-268.
55. Olsson PA, Burleigh SH, van Aarle IM: **The influence of external nitrogen on carbon allocation to *Glomus intraradices* in monoexinc arbuscular mycorrhiza.** *New Phytologist* 2005, **168**:677-686.
56. Liu C, Muchhal US, Uthappa M, Kononowicz AK, Raghothama KG: **Tomato phosphate transporter genes are differentially regulated in plant tissues by phosphorus.** *Plant Physiology* 1998, **116**:91-99.
57. Poulsen KH, Nagy R, Gao LL, Smith SE, Bucher M, Smith FA, Jakobsen I: **Physiological and molecular evidence for Pi uptake via the symbiotic pathway in a reduced mycorrhizal colonization mutant in tomato associated with a compatible fungus.** *New Phytologist* 2005, **168**:445-454.
58. Nagy R, Vasconcelos MJV, Zhao S, McElver J, Bruce W, Amrhein N, Raghothama KG, Bucher M: **Differential regulation of five Pht1 phosphate transporters from maize (*Zea mays* L.).** *Plant Biology* 2006, **8**:186-197.
59. Liu C, Muchhal US, Raghothama KG: **Differential expression of TPS11, a phosphate starvation-induced gene in tomato.** *Plant Molecular Biology* 1997, **33**:867-874.
60. Barker SJ, Stummer B, Gao L, Dispain I, O'Connor PJ, Smith SE: **A mutant in *Lycopersicon esculentum* Mill. with highly reduced VA mycorrhizal colonization: Isolation and preliminary characterisation.** *Plant Journal* 1998, **15**:791-797.
61. Miranda KM, Espey MG, Wink DA: **A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite.** *Nitric Oxide - Biology and Chemistry* 2001, **5**:62-71.
62. Foster JC: **Soil nitrogen.** In *Methods in applied soil microbiology and biochemistry* San Diego: Academic Press; 1995:79-87.
63. McGonigle TP, Miller MH, Evans DG, Fairchild GL, Swan JA: **A new method which gives an objective measure of colonization of roots by vesicular-arbuscular mycorrhizal fungi.** *New Phytologist* 1990, **115**:495-501.
64. Sah RN, Miller RO: **Spontaneous reaction for acid dissolution of biological tissues in closed vessels.** *Analytical Chemistry* 1992, **64**:230-233.
65. Benjamini Y, Hochberg Y: **Controlling the false discovery rate: a practical and powerful approach to multiple testing.** *Journal of the Royal Statistical Society Series B* 1995, **57**:289-300.
66. Verhoeven KJF, Simonsen KL, McIntyre LM: **Implementing false discovery rate control: Increasing your power.** *Oikos* 2005, **108**:643-647.
67. Bevan M, Bancroft I, Bent E, Love K, Goodman H, Dean C, Bergkamp R, Dirske W, Van Staveren M, Stiekema W, et al.: **Analysis of 1.9 Mb of**

contiguous sequence from chromosome 4 of *Arabidopsis thaliana*.
Nature 1998, **391**:485-488.

68. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: **Basic local alignment search tool.** *Journal of Molecular Biology* 1990, **215**:403-410.
69. Pfaffl MW: **A new mathematical model for relative quantification in real-time RT-PCR.** *Nucleic Acids Research* 2001, **29**:
70. Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method.** *Methods* 2001, **25**:402-408.

doi: 10.1186/1471-2229-10-75

Cite this article as: Ruzicka *et al.*, Tomato root transcriptome response to a nitrogen-enriched soil patch *BMC Plant Biology* 2010, **10**:75

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

