



2022 White House Conference on Hunger, Nutrition and Health

Issue: Healthy beverages for WIC participants

Problem: Americans' sugary drink consumption is too high – it is the largest source of added sugars in the American diet. Sugary drinks are loaded with calories and added sugars that deliver few or no nutritional benefits but increase the risk of myriad diet-related chronic diseases. Consumers' beverage behaviors are impaired by lack of knowledge about healthy choices and tap water may be unsafe or perceived as unsafe. A full array of interventions is needed to reduce this unnecessary risk to Americans' health and to enable them to drink plain water instead.

Pillars: *Integrate nutrition and health; Empower all consumers to make and have access to healthy choices*

Recommendations:

1. USDA, in collaboration with other appropriate stakeholders, should develop and disseminate a nutrition education component on drinking water safety and on healthy hydration habits for the Special Supplemental Nutrition Program for Women, Infants and Children (WIC).
2. Develop partnerships to work with local public water utilities to provide no-cost tap water lead testing for WIC households.
 - Model (unevaluated): The San Francisco Lead in Home Tap Water Testing Program, a San Francisco Public Utilities Commission program, provides free tap water lead testing for WIC clients with filter provision when needed.¹

Why does it matter? It is important to lay the foundation for healthy eating habits early in life.^{2,3,4} Over 40% of U.S. infants are in WIC-participating families.⁵

Although the WIC food package excludes sugary beverages and includes only 100% fruit juice, a study of 12–23-month-olds in WIC found that 25% drink no water and 31% consume sugary beverages (71% drink 100% fruit juice).⁶ Further, even in these early years, there are disparities in sugary drink and juice intakes related to income level and race/ethnicity.⁷

The American Heart Association (AHA) recommends that children consume no more than 25 grams (100 calories or about 6 teaspoons) of added sugars per day and that children under 2 years of age should avoid added sugars altogether. AHA states, “Although added sugars most likely can be safely consumed in low amounts as part of a healthy diet, few children achieve such levels, making this an important public health target.”⁸

Sugary drinks remain the largest single source of added sugars in the diets of American children aged 2–8 years; they are also among the top source of calories for US kids.^{9,10,11} Not only are these typically “empty” calories, but they also often displace more nutritious items.¹²

Extensive science shows that excess consumption of added sugars is a risk factor for many diet-related chronic diseases, and further, that independent of calories, added sugars have detrimental metabolic effects that are not due to weight gain and occur even in the absence of weight gain.^{13,14} Detrimental effects include but are not limited to:

- Obesity^{15,16}

- The 2015 Dietary Guidelines Advisory Committee stated, “Strong and consistent evidence shows that intake of added sugars from food and/or sugar-sweetened beverages are associated with excess body weight in children and adults. The reduction of added sugars and sugar-sweetened beverages in the diet reduces body mass index (BMI) in both children and adults.”¹⁷
- The rate of child obesity surged during the COVID-19 pandemic^{18,19}
- Pre-pandemic modelling showed that if then-current trends continued, the prevalence of obesity in the U.S. adult population will rise to 48.9% by 2030, while 24.2% of US adults will have severe obesity.²⁰
- Metabolic diseases including type 2 diabetes and fatty liver disease²¹
 - Non-alcoholic fatty liver disease is rising among children.²²
- Cardiovascular diseases²³
 - Even children can develop dyslipidemia and hypertension.²⁴
- Dental decay^{25,26}
 - Tooth decay is one of the most common chronic diseases of children and adolescents²⁷

These diet-related chronic conditions are also rife with disparities^{28,29, 30,31,32} and exacerbate economic inequities in the U.S.³³ For example, 44.3% of California children in families under 100% of federal poverty level (FPL) are overweight or obese while 21.2% of California children in families over 400% of FPL are overweight or obese.³⁴

Reducing drinking water lead exposure is also important for WIC families. Lead is a proven toxin, particularly for infants and young children. Infants are more vulnerable to adverse outcomes of lead exposure owing to high volume of water intake per body weight, increased lead absorption and rapid neuro-cognitive development.³⁵ Even low exposure to lead can reduce child IQ and is associated with attention deficit and problem behaviors;^{36,37} modelling shows that even low-level lead exposure reduces population IQ.^{38,39} Infants and children in WIC have been observed to be three times more likely to have elevated blood lead levels.⁴⁰

Homes in low-income communities are more apt to have lead exposures including lead service lines and lead in antiquated plumbing parts. There is limited data on lead in tap water in residential settings because public utilities are required regularly to test only a fraction of taps to monitor for lead in drinking water.⁴¹ A 2018 study of tap water in high-risk homes in New Orleans, found that though only 1% of all home water samples were above 15 ppb of lead, 12% were above 5 ppb (the FDA limit for bottled water) and 60% were above the American Academy of Pediatrics recommended level of 1 ppb.⁴² Nonetheless, it is estimated that 6.5-10 million American homes have lead service lines.⁴³

Lead ingested through tap water is a particular risk for infants fed powdered formula reconstituted with tap water. Sixty percent (60%) of U.S. infants 0-11 months are formula fed⁴⁴ and through formula they consume about 4 cups of water per day which means that 40-100% of their exposure to lead is through water used to mix formula.⁴⁵ Infants in WIC may be more likely to consume formula than non-WIC participants.⁴⁶

Risks of lead exposure during pregnancy and lactation can include elevated lead in the fetal brain and adverse outcomes of pregnancy; while lead can be detected in breastmilk, breastfed infants are generally at lower risk of lead exposure than are infants fed formula.⁴⁷ Lead exposure in US women of childbearing age is generally low yet identifying high-risk women (increased maternal age, race/ethnicity, poverty, immigrant) remains a public health need.⁴⁸

WIC education provides a vehicle on which to capitalize for teaching about healthy hydration for both children and parents, as well as a small number of basic water safety practices (see for example,⁴⁹ or⁵⁰) can make a difference before lead is mitigated through infrastructure improvements.

This recommendation is endorsed in the recent National Clinical Care Commission (NCCC) Report to Congress, “Leveraging Federal Programs to Prevent and Control Diabetes and Its Complications,”⁵¹

NCCC Recommendation 4.4: The National Clinical Care Commission recommends that all relevant federal agencies promote the consumption of water and reduce the consumption of sugar-sweetened beverages in the U.S. population, and that they employ all the necessary tools to achieve these goals, including education, communication, accessibility, water infrastructure, and sugar-sweetened beverage taxation.

- 4.4b. Child nutrition programs should be a conduit for education to promote consumption of water and reduce consumption of sugar-sweetened beverages. **USDA should encourage hydrating with water instead of sugar-sweetened beverages and provide safe water education in WIC nutrition education and in childcare settings.** Congress should harness the Child Nutrition Reauthorization Act to strengthen existing water provisions for school nutrition programs.

Who can act? Congress, EPA, USDA, DHSS; state and local health and social service departments; state departments overseeing tap water lead testing programs and childhood lead testing and lead poisoning prevention programs; federal, state and public-private partnerships, e.g., with public water utilities, university research teams, advocacy groups.

Prepared by Christina Hecht, PhD, Nutrition Policy Institute, University of California, Division of Agriculture and Natural Resources, ceahecht@ucanr.edu, for the White House Executive Office of the President, for the 2022 White House Conference on Hunger, Nutrition, and Health, with appreciation for members of the [National Drinking Water Alliance](#) who have developed the evidence base. 15 June 2022.

¹ San Francisco Public Utilities Commission. Contact: Andrew DeGraca, SFPUC, adegraca@sfwater.org, Kenneth Payne, SFPUC, kpayne@sfwater.org, June Weintraub SCD, MS, Manager, Water Regulatory Programs, SFDPH, oversees the WIC home testing program with SFPUC.

² Murray RD. Savoring Sweet: Sugars in Infant and Toddler Feeding. *Ann Nutr Metab* 2017. **70** Suppl 3:38-46.

³ Mennella, JA. Ontogeny of taste preferences: basic biology and implications for health. *Am J Clin Nutr* 2014. **99**, 704S-711S.10.3945/ajcn.113.067694.

⁴ Giddings SS, Mennella JA. 2016. *Has the world become too sweet?* American Heart Association Commentary. Available at <https://professional.heart.org/en/science-news/Added-Sugars-and-Cardiovascular-Disease-Risk-in-Children/commentary>

⁵ USDA FNS. 2022. WIC 2019 Eligibility and Coverage Rates. At, <https://www.fns.usda.gov/wic/2019-eligibility-coverage-rates>

⁶ Hamner HC, Paolicelli C, Casavale KO, Haake M, Bartholomew A. 2019. Food and Beverage Intake From 12 to 23 Months by WIC Status. *Pediatrics* 143 (3): e20182274. 10.1542/peds.2018-2274

⁷ Grimes CA, Szymlek-Gay EA, Nicklas TA. Beverage Consumption among U.S. Children Aged 0–24 Months: National Health and Nutrition Examination Survey (NHANES). *Nutrients*. 2017; **9**(3):264.

⁸ Vos MB, Kaar JL, Welsh JA, Van Horn LV, Feig DI, et al. American Heart Association Nutrition Committee of the Council on Lifestyle and Cardiometabolic Health; Council on Clinical Cardiology; Council on Cardiovascular Disease in the Young; Council on Cardiovascular and Stroke Nursing; Council on Epidemiology and Prevention; Council on Functional Genomics and Translational Biology; and Council on Hypertension. Added Sugars and Cardiovascular Disease Risk in Children: A Scientific Statement From the American Heart Association. *Circulation*. 2017. **135**(19):e1017-e1034.

⁹ Reedy J, Krebs-Smith SM. Dietary sources of energy, solid fats, and added sugars among children and adolescents in the United States. *J Am Diet Assoc*. 2010 Oct;**110**(10):1477-84.

-
- ¹⁰ Ricciuto L, Fulgoni VL, Gaine PC, Scott MO, DiFrancesco L. 2021. Sources of Added Sugars Intake Among the U.S. Population: Analysis by Selected Sociodemographic Factors Using the National Health and Nutrition Examination Survey 2011–18. *Frontiers in Nutrition* 8. DOI=10.3389/fnut.2021.687643
- ¹¹ Dietary Guidelines Advisory Committee. 2020. *Scientific Report of the 2020 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Agriculture and the Secretary of Health and Human Services*. U.S. Department of Agriculture, Agricultural Research Service, Washington, DC. At https://www.dietaryguidelines.gov/sites/default/files/2020-07/ScientificReport_of_the_2020DietaryGuidelinesAdvisoryCommittee_first-print.pdf
- ¹² Dietary Guidelines Advisory Committee. 2020. *Scientific Report of the 2020 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Agriculture and the Secretary of Health and Human Services*. U.S. Department of Agriculture, Agricultural Research Service, Washington, DC. At https://www.dietaryguidelines.gov/sites/default/files/2020-07/ScientificReport_of_the_2020DietaryGuidelinesAdvisoryCommittee_first-print.pdf
- ¹³ Stanhope KL, Goran MI, Bosy-Westphal A, King JC, Schmidt LA, et al. Pathways and mechanisms linking dietary components to cardiometabolic disease: thinking beyond calories. *Obes Rev*. 2018. 19(9):1205-1235.
- ¹⁴ Lustig R, Schmidt LA & Brindis C. “The Toxic Truth About Sugar.” *Nature*, 2012. 482: 27-9.
- ¹⁵ Malik VS, Popkin BM, Bray GA, Després J-P, Hu FB Sugar-sweetened beverages, obesity, type 2 diabetes mellitus, and cardiovascular disease risk. *Circulation*. 2010. 121:1356–1364
- ¹⁶ Malik VS, Pan A, Willett WC, Hu FB. Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis. *Am J Clin Nutr*. 2013;98(4):1084-102.
- ¹⁷ Dietary Guidelines Advisory Committee. 2015. *Scientific Report of the 2015 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Health and Human Services and the Secretary of Agriculture*. U.S. Department of Agriculture, Agricultural Research Service, Washington, DC.
- ¹⁸ Lange SJ, Kompaniyets L, Freedman DS, et al. 2021. Longitudinal trends in body mass index before and during the covid-19 pandemic among persons aged 2-19 years—United States, 2018-2020. *CDC Morbidity and Mortality Weekly Report*. At, www.cdc.gov/mmwr/volumes/70/wr/mm7037a3.htm.
- ¹⁹ Woolford SJ, Sidell M, Li X, Else V, Young DR, Resnicow K, Koebnick C. Changes in Body Mass Index Among Children and Adolescents During the COVID-19 Pandemic. *JAMA*. 2021 Oct 12;326(14):1434-1436.
- ²⁰ Ward ZJ, et al. Projected U.S. State-Level Prevalence of Adult Obesity and Severe Obesity. *N Engl J Med* 2019. 381:2440-50.
- ²¹ Neuenschwander M, Ballon A, Weber KS, Norat T, Aune D, Schwingshackl L, Schlesinger S. Role of diet in type 2 diabetes incidence: umbrella review of meta-analyses of prospective observational studies. *BMJ*. 2019. 366:12368.
- ²² Uppal V, Mansoor S, Furuya KN. Pediatric Non-alcoholic Fatty Liver Disease. *Curr Gastroenterol Rep*. 2016. 18(5):24.
- ²³ Yang Q, Zhang Z, Gregg EW, Flanders WD, Merritt R, Hu FB. Added sugar intake and cardiovascular diseases mortality among US adults. *JAMA Intern Med*. 2014. 174(4):516-24.
- ²⁴ Vos MB, Kaar JL, Welsh JA et al. Added sugars and cardiovascular disease risk in children: A scientific statement from the American Heart Association. *Circulation*. 2017. 135: e1017-e1034.
- ²⁵ Chi DL, Scott JM. Added Sugar and Dental Caries in Children: A Scientific Update and Future Steps. *Dent Clin N Am*. 2019. 63:17-33.
- ²⁶ Bleich S, Vercammen K. The negative impact of sugar-sweetened beverages on children’s health: an update of the literature. *BMC Obes* 2018; 5:6.
- ²⁷ National Institutes of Health. *Oral Health in America: Advances and Challenges*. Bethesda, MD: US Department of Health and Human Services, National Institutes of Health, National Institute of Dental and Craniofacial Research, 2021. At, <https://www.nidcr.nih.gov/sites/default/files/2021-12/Oral-Health-in-America-Advances-and-Challenges.pdf>
- ²⁸ Hales CM, Carroll MD, Fryar CD & Ogden CL. (2017, October). Prevalence of obesity among adults and youth, United States, 2015-2016 (Data Brief No. 288). Retrieved from National Center for Health Statistics website: <https://www.cdc.gov/nchs/data/databriefs/db288.pdf>
- ²⁹ Taveras EM, Gillman MW, Kleinman KP, Rich-Edwards JW, Rifas-Shiman SL. Reducing Racial/Ethnic Disparities in Childhood Obesity: The Role of Early Life Risk Factors. *JAMA Pediatr*. 2013. 167(8):731–738.
- ³⁰ Centers for Disease Control and Prevention. (2018). Diabetes report card 2017. Retrieved from <https://www.cdc.gov/diabetes/pdfs/library/diabetesreportcard2017-508.pdf> .
- ³¹ Centers for Disease Control and Prevention. *Health, United States, 2015: With Special Feature on Racial and Ethnic Health Disparities*. Hyattsville, MD; 2015.
- ³² Muth ND, Dietz WH, Magge SN, et al. AAP American Academy of Pediatrics, AAP Section on Obesity, AAP Committee on Nutrition, AAP American Heart Association. Public Policies to Reduce Sugary Drink Consumption in Children and Adolescents. *Pediatrics*. 2019. 143(4):e20190282.

-
- ³³ NASEM (National Academies of Sciences, Engineering, Medicine). 2017. *Communities in Action: Pathways to Health Equity*. Washington, DC: The National Academies Press. At, <https://www.nap.edu/catalog/24624/communities-in-action-pathways-to-health-equity>
- ³⁴ California State Fact Sheet. At <https://www.childhealthdata.org/docs/nsch-docs/california-pdf.pdf>
- ³⁵ Levallois P, Barn P, Valcke M, Gauvin D, Kosatsky T. Public Health Consequences of Lead in Drinking Water. *Curr Environ Health Rep*. 2018;5(2):255-262.
- ³⁶ National Toxicology Program. *Monograph on Health Effects of Low-Level Lead*. Research Triangle Park, NC: National Institute of Environmental Health Sciences; 2012:xiii, xv-148.
- ³⁷ Lanphear BP, Dietrich K, Auinger P, Cox C. Cognitive deficits associated with blood lead concentrations <10 microg/dL in US children and adolescents. *Public Health Rep*. 2000;115(6):521-529.
- ³⁸ Bellinger DC. A strategy for comparing the contributions of environmental chemicals and other risk factors to neurodevelopment of children. *Environ Health Perspect*. 2012;120(4):501-507.
- ³⁹ Lanphear BP, Hornung R, Khoury J, et al. Low-level environmental lead exposure and children's intellectual function: an international pooled analysis. *Environ Health Perspect*. 2005;113(7):894-899.
- ⁴⁰ Aoki Y, Brody DJ. WIC Participation and Blood Lead Levels among Children 1-5 Years: 2007-2014. *Environ Health Perspect*. 2018;126(6):067011.
- ⁴¹ Environmental Law Institute. *Preventing Toxic Lead Exposure Through Drinking Water Using Point-of-Use Filtration*. Available at: <https://www.eli.org/sites/default/files/elr/featuredarticles/48.11074.pdf>. Accessed on September 29, 2019.
- ⁴² Katner A, Pieper K, Brown K, Lin HY, Parks J, Wang X, Hu CY, Masters S, Mielke H, Edwards M. Effectiveness of Prevailing Flush Guidelines to Prevent Exposure to Lead in Tap Water. *Int J Environ Res Public Health*. 2018 Jul 20;15(7):1537.
- ⁴³ EPA. 2016. *Lead and Copper Rule Revisions White Paper*. At, https://www.epa.gov/sites/default/files/2016-10/documents/508_lcr_revisions_white_paper_final_10.26.16.pdf
- ⁴⁴ Grimes CA, Szymlek-Gay EA, Nicklas TA. Beverage Consumption among U.S. Children Aged 0-24 Months: National Health and Nutrition Examination Survey (NHANES). *Nutrients*. 2017; 9(3):264
- ⁴⁵ Triantafyllidou S, Gallagher D, Edwards M. 2014. Assessing risk with increasingly stringent public health goals: the case of water lead and blood lead in children. *J Water Health* 12(1):57-68,
- ⁴⁶ Jun S, Catellier DJ, Eldridge AL, Dwyer JT, Eicher-Miller HA, Bailey RL. Usual Nutrient Intakes from the Diets of US Children by WIC Participation and Income: Findings from the Feeding Infants and Toddlers Study (FITS) 2016. *J Nutr*. 2018;148(9S):1567S-1574S
- ⁴⁷ American College of Obstetricians and Gynecologists, Committee on Obstetric Practice. *Lead Screening During Pregnancy and Lactation*. 2019. Available at https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2012/08/lead-screening-during-pregnancy-and-lactation?utm_source=redirect&utm_medium=web&utm_campaign=otn Accessed 3/23/21
- ⁴⁸ Ettinger AS, Egan KB, Homa DM, Brown MJ. Blood Lead Levels in U.S. Women of Childbearing Age, 1976-2016. *Environ Health Perspect*. 2020 128:1
- ⁴⁹ Minnesota Department of Health. *Let it run ... and get the lead out!* At, https://www.health.state.mn.us/communities/environment/water/factsheet/letitrun_english.html
- ⁵⁰ American Academy of Pediatrics. *Lead in Tap Water & Household Plumbing: Parent FAQs - Simple Safety Steps*. At, <https://www.healthychildren.org/English/safety-prevention/at-home/Pages/Lead-in-Tap-Water-Household-Plumbing.aspx>
- ⁵¹ National Clinical Care Commission. 2021. *Report to Congress on Leveraging Federal Programs to Prevent and Control Diabetes and Its Complications*. At, <https://health.gov/about-odphp/committees-workgroups/national-clinical-care-commission/report-congress>