HEALTHY FOOD SYSTEMS . HEALTHY ENVIRONMENTS

University of California

Agriculture and Natural Resources

Making a Difference for California



WEEKLY SOIL MOISTURE LOSS IN INCHES (Estimated Crop Evapotranspiration or ET_c)

	(Esti	•	Evapoulan	•	$\mathbf{E}(\mathbf{C})$						
		04/14/2	3 through (04/20/23							
Crops (Leafout Date)	#	4148 Merce	d				#258 Lemon Cove				
	4/14 - 4/20	Accum'd	4/21 - 4/27		4/14 - 4/20	Accum'd	4/21 - 4/27	4/14 - 4/20	Accum'd	4/21 - 4/27	
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc	
Almonds (3/13) *	0.99	3.80	1.15		1.08	4.10	1.19	0.96	3.72	1.18	
Pistachio (NA) * **	0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	
Citrus (2/1)	0.90	5.83	1.00		0.99	6.27	1.02	0.87	5.80	1.02	
Raisin Grapes (3/22) (11 ft. row spacing)	0.11	0.26	0.12		0.11	0.26	0.12	0.10	0.25	0.12	
Winegrapes (3/22) (10 ft. spacing on California Sprawl Trellis)	0.18	0.50	0.22		0.20	0.55	0.23	0.17	0.48	0.23	
Walnuts (NA)	0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	
Stone Fruit (3/19)	0.63	1.82	0.76		0.69	1.92	0.79	0.61	1.77	0.78	
Past 7 days precipitation (inches)		0.00				0.00			0.00		
Accumulated precipitation (inches) (1/1/2023)		15.98				12.32			15.50		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

** Raisin Grapes and Winegrapes Irrigation should hold off until midday leaf water potential drops to -1.0 MPa, before that soil moisture reservoir is sufficient to supply the vine water demand. Update will be sent shortly once the county wide leaf water potential reaches approximate -1.0 MPa. Growers should adjust the irrigation start date based on the individual vineyard location and soil type.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹												
Crops		#148 Merced				#39 Parlier	•		#258 Lemon Cove			
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/13)	1.5	1.3	1.2	1.0	1.7	1.4	1.3	1.1	1.5	1.3	1.1	1.0
Pistachio (NA)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Citrus (2/1)	1.4	1.2	1.1	0.9	1.5	1.3	1.2	1.0	1.3	1.2	1.0	0.9
Raisin Grapes (3/22) (11 ft. row spacing)***	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Winegrapes (3/22) (10 ft. spacing on California Sprawl Trellis) ***	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Walnuts (NA)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Stone Fruit (3/19)	1.0	0.8	0.7	0.7	1.1	0.9	0.8	0.7	0.9	0.8	0.7	0.6

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE															
Crops		#148 Merced #39 Parlier								#258 Lemon Cove					
Almonds 115 Trees/A	354	307	283	236	401	331	307	260	354	307	260	236			
Pistachio 106 Trees/A	0	0	0	0	0	0	0	0	0	0	0	0			
Citrus 110 Trees/A	346	296	272	222	370	321	296	247	321	296	247	222			
Raisin Grapes 566 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0			
Winegrapes 622 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0			
Walnuts 76 Trees/A	0	0	0	0	0	0	0	0	0	0	0	0			
Stonefruit 172 Trees/A	158	126	111	111	174	142	126	111	142	126	111	95			
For further information concerning all counties receiving this report, contac	t the Fresno	Co. Farm Ad	lvisor's offic	e at (559) 2-	41-7526.										

HEALTHY FOOD SYSTEMS . HEALTHY ENVIRONMENTS

University of California

Agriculture and Natural Resources

Making a Difference for California



UCCE/DWR Weekly Crop Wat Report

WEEKLY SOIL MOISTURE LOSS IN INCHES (Estimated Crop Evapotranspiration or ET_{c})

$\frac{04}{14}/23 \text{ through } 04/20/23$													
		04/14/2	.5 through (14/20/23									
Crops (Leafout Date)	#	124 Panocł	ne		#	2 Five Poin	ts		#15 Stratford				
	4/14 - 4/20	Accum'd	4/21 - 4/27		4/14 - 4/20	Accum'd	4/21 - 4/27		4/14 - 4/20	Accum'd	4/21 - 4/27		
	Water	Seasonal	Estimated		Water	Seasonal	Estimated		Water	Seasonal	Estimated		
	Use	Water Use	ETc		Use	Water Use	ETc		Use	Water Use	ETc		
Almonds (3/13) *	1.01	3.95	1.35		1.08	4.05	1.41		1.11	4.36	1.37		
Pistachio (NA) * **	0.00	0.00	0.00		0.00	0.00	0.00		0.00	0.00	0.00		
Citrus (2/1)	0.91	6.04	1.14		0.96	6.27	1.21		0.98	6.68	1.17		
Raisin Grapes (3/22) (11 ft. row spacing)	0.09	0.24	0.13		0.10	0.26	0.15		0.11	0.27	0.13		
Winegrapes (3/22) (10 ft. spacing on California Sprawl Trellis)	0.18	0.50	0.25		0.20	0.54	0.26		0.20	0.57	0.25		
Walnuts (NA)	0.00	0.00	0.00		0.00	0.00	0.00		0.00	0.00	0.00		
Stone Fruit (3/19)	0.66	1.91	0.87		0.71	1.98	0.90		0.73	2.12	0.88		
Past 7 days precipitation (inches)		0.00				0.00				0.00			
Accumulated precipitation (inches) (1/1/2023)		5.37				7.35				7.62			

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

** Raisin Grapes and Winegrapes Irrigation should hold off until midday leaf water potential drops to -1.0 MPa, before that soil moisture reservoir is sufficient to supply the vine water demand. Update will be sent shortly once the county wide leaf water potential reaches approximate -1.0 MPa. Growers should adjust the irrigation start date based on the individual vineyard location and soil type.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹												
Crops		#124 Panoche				#2 Five Poi	nts		#15 Stratford			
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/13)	1.6	1.3	1.2	1.1	1.7	1.4	1.3	1.1	1.7	1.5	1.3	1.2
Pistachio (NA)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Citrus (2/1)	1.4	1.2	1.1	1.0	1.5	1.3	1.1	1.0	1.5	1.3	1.2	1.0
Raisin Grapes (3/22) (11 ft. row spacing)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Winegrapes (3/22) (10 ft. spacing on California Sprawl Trellis)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Walnuts (NA)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Stone Fruit (3/19)	1.0	0.9	0.8	0.7	1.1	0.9	0.8	0.7	1.1	1.0	0.9	0.8

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE												
Crops		#124 Panoche #2 Five Points										
Almonds 115 Trees/A	378	307	283	260	401	331	307	260	401	354	307	283
Pistachio 106 Trees/A	0	0	0	0	0	0	0	0	0	0	0	0
Citrus 110 Trees/A	346	296	272	247	370	321	272	247	370	321	296	247
Raisin Grapes 566 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0
Winegrapes 622 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0
Walnuts 76 Trees/A	0	0	0	0	0	0	0	0	0	0	0	0
Stonefruit 172 Trees/A	158	142	126	111	174	142	126	111	174	158	142	126
For further information concerning all counties receiving this report, co	ntact the Fresno	Co. Farm Ad	lvisor's offic	e at (559) 2-	41-7526.				•			

r c	ALIF
tor	Use
er	USE

ORNIANS