

Studies on Strawberry Quality

high temperatures that may occur in harvesting and handling of strawberries influence the rate of fruit deterioration

E. C. Maxie, F. Gordon Mitchell, and Arthur Greathead

The following article is the second of two reports on Quality Studies with Strawberries. The first report appeared in the January issue.

Strawberries and the decay organisms that attack them are alive and temperature has a tremendous influence on the rates of their life processes. At temperatures between 32°F and 75°F the rates of these life processes would be expected to increase two to four fold for each 18°F temperature increase. The deterioration of the fruit depends on how fast these processes occur. Thus the rate of deterioration can be expected to be influenced in the same way by temperature.

To measure the effect of temperature on strawberries, studies were started by measuring the respiration rate of the Shasta variety. Rate of respiration closely parallels the rate of other life processes in both fruits and decay organisms.

Fruit was obtained from plantings at Davis. The berries were picked in the early morning and immediately used in the experiment. Duplicate 500 gram samples were prepared for each treatment. The fruit was selected for freedom from defects, weighed, and sealed in

respiration jars. These jars were then placed at five different temperatures—32, 41, 68, 86, and 100°F.

As shown in the lower left graph, temperature has a marked influence on the respiration rate of the Shasta strawberry. The initial respiration rates at the higher temperatures exceeded the rate at 32°F by the following factors: 41°F—1.4 times, 68°F—7.2 times, 86°F—19 times, and 100°F—19.6 times. After about six hours oily, red droplets appeared on the surface of berries held at 86°F and at 100°F, indicating skin damage. This was followed by bleaching of the red in the berries, and still later by mushy breakdown of the tissue. No mold growth was observed at 86°F and 100°F. The tests at the lower temperatures were discontinued when mold appeared on the berries.


These results clearly show the effect of temperature on the two major causes of strawberry deterioration—decay and self-destruction. At 86°F and 100°F the berries were quickly destroyed by the speed of their own life processes. The berries at 68°F showed mold growth

after two and one half days; at 41°F after five days; and, at 32°F, the berries showed mold after 11 days. Under commercial conditions, with picking damage added, it is probable that deterioration would be more rapid than in these tests.


The effect of delayed cooling at high temperatures on the keeping quality of the Shasta variety was next considered. Two large insulated boxes were used. One was fitted for thermostatically controlled heating, the other for forced-air cooling. The warm-air box was preheated to 85°F and held at that temperature for the duration of the experiment. The cool box was iced, and cooled to about 41°F. Sixteen crates of berries were picked by the commercial picking crew on a farm near Salinas. The picking required less than one hour and was completed at 8:30 a.m. Two crates were cooled immediately. The other 14 crates were placed in the warm box and held without air circulation. These crates were transferred to the cold box, two at a time, at the end of one half, one, two, three, four, six, and eight hours. All fruit was quickly

Concluded on page 16

The influence of temperature on the respiration rate of variety Shasta strawberries.



The effect of delayed cooling at high temperatures on keeping quality of Shasta strawberries.



STRAWBERRIES

Continued from page 11

cooled using the forced air principle, and then held for seven days at 41°F. It was then removed and sorted immediately into four classes: sound or undamaged, badly bruised but not decayed, slight decay, and severe decay. Sound fruit and bruised but not decayed berries were considered marketable.

As shown in the lower right graph on page 11, the quality of strawberries drops sharply after two hours at 85°F. The fruits represented by the graph were evaluated while still cool.

All badly bruised berries could be expected to be lost within a short time in commercial marketing channels. Under these conditions only the sound fruit would be marketable.

Another important trend is the increase in decay with increasing time at this high temperature. It seems likely that berries held for more than three hours would have been an almost complete loss in retail channels within 24 hours.

These experiments indicate that strawberries must be cooled very soon after harvest if satisfactory quality is to be

maintained. While the temperature used may seem high for some strawberry growing districts, the information obtained in the studies indicates that such temperatures may be quite commonly reached or exceeded under actual field conditions.

The grower often can not move his berries to the cooler as quickly as would be desirable. However, every effort should be made to give maximum protection to the fruit. It should be placed in the shade if possible, hauled in a covered closed truck, moved to the cooler as soon

as practical, cooled rapidly and kept cool thereafter. These experiments show clearly that strawberries would quickly destroy themselves at high temperatures, even if they were sterile of all decay organisms.

E. C. Maxie is Assistant Pomologist, University of California, Davis.

F. Gordon Mitchell is Extension Marketing Technologist, University of California, Davis.

Arthur Greathead is Farm Advisor, Monterey County, University of California.

The above progress report is based on Research Project No. 1656.

DONATIONS FOR AGRICULTURAL RESEARCH

Contributions to the University of California, Division of Agricultural Sciences

BERKELEY

Butte County	\$500.00
For timber taxation study	
California Cedar Products Company	\$900.00
For research in wood machining	
Union Carbide Chemical Company5 gals. Carbowax
For research on forest products	

DAVIS

Abbott Laboratories	30 grams Gibberellic Acid
For research on dormancy in pears	
American Cyanamid Company	\$3,500.00
For research on infectious diseases of poultry	
American Poultry and Hatchery Federation	\$1,500
For research on host tumor responses in poultry	
Bou-Matic Milkers	Portable pipeline milking machine
For research on the relationship of mechanical milking to mastitis problem	
Bush Berry Advisory Board	1,950 lbs. boysenberries
For research in food technology	
California Bull Association	4 purebred Jersey bulls
For research on genetics of dairy cattle	
California Cattle Feeders Association	\$666.66
For progeny testing program	
California Committee on the Relationship of Electricity to Agriculture	\$7,187.50
For research in the use of electricity in agriculture (Fourth quarterly payment on total of \$28,750.00)	
California Spray-Chemical Corporation	\$2,000.00
For research on insecticides	
Fibreboard Paper Products Corp.	600 experimental fibreboard boxes
For studies on the shipping of apples and pears	
Sugar Research Foundation, Inc.	
For research on the effect of varying sugar acid ratio on the palatability of wine	
	\$1,250.00
(Second quarterly payment on total of \$5,000.00)	
For research on the effect of sweetness on the consumers acceptance of apricots, pears, and peaches	
	\$2,500.00
Syntex Animal Products	\$1,500.00
For research on hormones in heifers and cows of the beef cattle breeds	
U. S. Public Health Service	\$4,053.00
For research on the use of enzymes for tenderizing meat	
Western States Meat Packers Assn., Inc.	\$666.66
For research on economic and quality beef	

LOS ANGELES

Davison Chemical Company	2 Verifax machines
	1 Waring blender
	3 desiccators
	4 dust sieves
For research on silica aerogel SG 67	

RIVERSIDE

California Spray-Chemical Corporation	\$1,000.00
For research on fungicides	

STATEWIDE

Chemical and Pigment Company	600 lbs. zinc sulfate
For micro-nutrient studies in Colusa County and zinc deficiency studies elsewhere in the state	
Dow Chemical Company	150 lbs. powdered Trolene
	200 lbs. Trolene boluses
For studies on the effectiveness of the materials as systemic grubicides in Imperial County	
E. I. du Pont de Nemours & Company	300 lbs. Uramite
For studies on turf variety plots in Los Angeles County	
The Garland Company	200 lbs. Garland Shading Compound Purple
For experiments on greenhouse flowers in San Mateo County	
Jack Wilson Chemical Company	15 lbs. DIXA Weed Killer
For experimental work on morning glory in Imperial County	
Monsanto Chemical Company	
For studies on control of weeds in lettuce, Imperial County	
	10 gals. Vegadex
For studies on control of weeds in onions, Imperial County	
	Randex
The Mountain Copper Company	100 lbs. copper sulfate
For micro-nutrient studies in Colusa County	
Naugatuck Chemical Co.	2 qts. 3YD
For test plots on eradication of nutgrass and morning glory in Imperial County	
Oasis Chemical Company, Inc.	50 lbs. Chloro IPC 10
For test plot work in Imperial County	
Pennsalt Manufacturing Co.	30 gals. Endothal
For establishing defoliation test plots on seed flax and alfalfa	
Shell Chemical Corporation	1 gal. aldrin
For test plot on cricket control in field and vegetable crops in Imperial County	
Stauffer Chemical Company	
For exploratory trials on phosphate fertilization of annual clovers on open ranges in northern California	
	5 tons superphosphate
For fumigation of turfgrass variety plots in Los Angeles County	
	25 gals. Vapam



Penalty for private use to avoid payment of postage, \$300

University of California College of Agriculture, Agricultural Experiment Station, Berkeley 4, California

Paul J. Sharp
Director

Free—Annual Report or Bulletin or Report of Progress
Permit No. 1127