



Synthesis Paper: Targeted Livestock Grazing: Prescription for Healthy Rangelands

Derek W. Bailey ^{a,*}, Jeffrey C. Mosley ^b, Richard E. Estell ^c, Andres F. Cibils ^d, Marc Horney ^e, John R. Hendrickson ^f, John W. Walker ^g, Karen L. Launchbaugh ^h, Elizabeth A. Burritt ⁱ

^a New Mexico State University, Las Cruces, NM 88003, USA

^b Montana State University, Department of Animal and Range Sciences, Bozeman, MT 59717, USA

^c Jornada Experimental Range, USDA-ARS, Las Cruces, NM 88003, USA

^d New Mexico State University, Las Cruces, NM 88003, USA

^e California Polytechnic State University, San Luis Obispo, San Luis Obispo, CA 93407, USA

^f Northern Great Plains Research Laboratory, USDA-ARS, Mandan, ND 58554, USA

^g Texas A&M AgriLife Research and Extension Center, San Angelo, TX 76901, USA

^h University of Idaho, Moscow, ID 83844, USA

ⁱ Utah State University, Logan, UT 84322

ARTICLE INFO

Article history:

Received 24 December 2018

Received in revised form 14 June 2019

Accepted 17 June 2019

Key Words:

focused grazing

livestock

prescriptive grazing

rangelands

targeted grazing

vegetation manipulation

ABSTRACT

Targeted livestock grazing is a proven tool for manipulating rangeland vegetation, and current knowledge about targeted livestock grazing is extensive and expanding rapidly. Targeted grazing prescriptions optimize the timing, frequency, intensity, and selectivity of grazing (or browsing) in combinations that purposely exert grazing/browsing pressure on specific plant species or portions of the landscape. Targeted grazing differs from traditional grazing management in that the goal of targeted grazing is to apply defoliation or trampling to achieve specific vegetation management objectives, whereas the goal of traditional livestock grazing management is generally the production of livestock commodities. A shared aim of targeted livestock grazing and traditional grazing management is to sustain healthy soils, flora, fauna, and water resources that, in turn, can sustain natural ecological processes (e.g., nutrient cycle, water cycle, energy flow). Targeted grazing prescriptions integrate knowledge of plant ecology, livestock nutrition, and livestock foraging behavior. Livestock can be focused on target areas through fencing, herding, or supplement placement. Although practices can be developed to minimize the impact of toxins contained in target plants, the welfare of the animals used in targeted grazing must be a priority. Monitoring is needed to determine if targeted grazing is successful and to refine techniques to improve efficacy and efficiency. Examples of previous research studies and approaches are presented to highlight the ecological benefits that can be achieved when targeted grazing is applied properly. These cases include ways to suppress invasive plants and ways to enhance wildlife habitat and biodiversity. Future research should address the potential to select more adapted and effective livestock for targeted grazing and the associated animal welfare concerns with this practice. Targeted livestock grazing provides land managers a viable alternative to mechanical, chemical, and prescribed fire treatments to manipulate rangeland vegetation.

© 2019 The Authors. Published by Elsevier Inc. on behalf of The Society for Range Management. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

INTRODUCTION

Rangeland managers have a wide range of practices available to maintain rangeland health and productivity or restore degraded rangelands. Most of these practices fit in the categories of axe, plow, fire, cow, and gun that Leopold (1933) suggested were the keys to

managing wildlife. Managers can successfully use traditional livestock grazing management, prescribed fire, mechanical interventions, and herbicide application alone or in combination to manage rangeland vegetation (Masters and Sheley, 2001; Briske et al., 2011). However, tradeoffs exist for every rangeland management practice. Mechanical treatments usually require relatively gentle terrain and are typically expensive (Herbel, 1983; DiTomaso, 2000). Herbicides can have undesirable impacts on nontarget vegetation, elicit concerns for human health, be expensive, and are often not feasible without cost-share programs (DiTomaso, 2000; Shepard et al., 2004; Torell et al.,

* Correspondence: Derek W. Bailey, New Mexico State University, PO Box 30003, MSC 3-1, Las Cruces, NM 88003, USA.

E-mail address: dwbailey@nmsu.edu (D.W. Bailey).

2005). Prescribed fire requires specific conditions to be safely and effectively applied (Fuhlendorf and Engle, 2004), and there is a risk that it can lead to catastrophic wildfires (Yoder, 2004; Yoder et al., 2004). Targeted grazing is an alternative and/or complementary vegetation management tool. Further, targeted livestock grazing is an “environmentally friendly” alternative to traditional methods because it can be applied to extensive inaccessible areas, leaves no chemical herbicide residue, can be removed whenever necessary, and often improves biodiversity. Plus, in the process of removing plant biomass, grazing animals convert it into saleable products—meat and fiber. A survey in Ontario, Canada found that target grazing was the second most preferred method of vegetation control behind direct removal by cutting (Wagner et al., 1998).

Targeted grazing is the application of a specific kind of livestock at a determined season, duration, and intensity to accomplish defined vegetation or landscape goals (Frost and Launchbaugh, 2003). Targeted grazing has been referred to elsewhere, especially in 20th century research literature, as *prescribed grazing*, *prescriptive grazing*, or similar terms (Walker, 1995; Frost and Launchbaugh, 2003). Targeted grazing continues to be a focus of rangeland managers and scientists. The best evidence of the value and growth of targeted grazing is the development and success of targeted grazing businesses. Entrepreneurs are paid to use livestock to control noxious weeds and other invasive plants, provide fire breaks, and for other contemporary vegetation management challenges (Frost et al., 2012). Although the field is growing, additional research and outreach are needed to ensure that targeted grazing becomes a tool that rangeland managers can readily employ to resolve land management issues. The objectives of this paper are to define the practice of “targeted grazing”; describe the information, resources, and skills it requires; and present evidence for specific ecological benefits that can be achieved when targeted livestock grazing is properly used as an element of a management plan.

ORIGIN AND USE OF THE TERM “TARGETED GRAZING”

The use of livestock grazing to accomplish defined vegetation or landscape goals has been referred to as prescribed, targeted, or conservation grazing. Currently this type of grazing management is usually referred to as *targeted grazing*. The term *conservation grazing* is also in common use but typically refers to a less intensive form of management used to maintain and increase the biodiversity of natural or seminatural ecosystems. We searched the Clarivate Analytics Web of Science database using a topic search for the terms “conservation grazing,” “prescribed grazing,” or “targeted grazing” to identify journal articles containing those terms. The articles are provided in Supplement 1 (available online at <https://doi.org/10.1016/j.rama.2019.06.003>) and were examined to ensure that the use of the term referred to livestock grazing management. Before 2006, most writings about using livestock to manage invasive plants or improve wildlife habitat referred to prescribed grazing, but the National Range and Pasture Handbook (Butler et al., 2003) used the term prescribed grazing to describe any type of planned grazing. The term *targeted grazing* came into widespread use after publication of the Targeted Grazing Handbook (Launchbaugh and Walker, 2006). Targeted grazing was used in that publication to differentiate grazing management for a specific vegetation management goals from the broader *prescribed grazing* term. The term targeted grazing was first used in a peer-reviewed journal in 1996 (Holst and Allan, 1996), but the citation is not in the Web of Science database and is not included in Figure 1. A total of 89 publications were found using one of these terms, and the earliest paper used prescribed grazing and was published in 1991. Nine of these articles used both targeted and prescribed grazing, but with one exception the second term was only used as a key word and the paper was classified for the term that was most used. Figure 1 shows that since 2010, targeted grazing has been the preferred term compared with prescribed grazing. Journal articles

that used the terms prescribed, targeted, or conservation grazing were found in 51 different journals with nearly half (43%) of the articles in *Rangeland Ecology & Management*, in which the term targeted grazing was most prevalent (55%).

TARGETED GRAZING PRINCIPLES

Managers can manipulate livestock impacts to vegetation using the principles of grazing management (Valentine, 2001): stocking rate, distribution, species of livestock, and season (or timing) of grazing. The overall intensity of defoliation or forage utilization level can be increased by increasing stocking rate (Table 1). Stocking density can also play a role on the use of targeted plants. Utsumi et al. (2010) found that utilization of one-seed juniper (*Juniper monosperma* [Englem.] Sarg.) saplings was greater at high stocking densities than low densities at similar stocking rates. Grazing can be focused or redirected by applying distribution practices such as fencing, water developments, and herding. Diet selection can be modified by changing the species of livestock (e.g., use of forbs can be increased by switching from cattle to sheep). Changing the season of grazing can affect livestock preferences for vegetation and correspondingly diet selection, which can affect the impact of defoliation on plant vigor. The grazing management principles are the same for both traditional, production-based management and targeted grazing (Table 1), but the goals and priorities differ (Frost et al., 2012). Left unchecked, livestock grazing can lead to deterioration in rangeland health and reduced livestock productivity as animals focus on preferred plants and areas near water (Valentine, 2001). Managers use grazing management principles to prevent overuse of preferred plants and promote livestock gains and reproductive performance. Similarly, targeted grazers can use grazing management principles so that livestock can be used to increase use of undesirable plants at a time in their life cycle when they are most vulnerable to defoliation. For example, a rancher might develop a new water source to redirect grazing pressure to areas near the new water location and improve uniformity of grazing across the pasture, while implementation of targeted grazing might require temporary electric fencing or herding to focus livestock to areas with higher densities of invasive weeds.

The application of grazing management principles to targeted grazing depends on the issue being addressed (Launchbaugh and Walker, 2006). For example, species of livestock is critical for control of leafy spurge (*Euphorbia esula* L.), but timing is not as critical. Leafy spurge is not palatable to cattle, and sheep or goats must be used for control of leafy spurge. For cheatgrass (a.k.a., downy brome) (*Bromus tectorum* L.), the species of livestock is less important, but the timing of grazing is more critical. Most livestock species will consume this invasive annual grass early in the spring when it is green and growing but usually avoid it during the summer after it matures and other perennial grasses are still green and growing (Cook and Harris, 1952; Mosley, 1996).

PLANT ECOPHYSIOLOGY AND SUCCESSION CONSIDERATIONS FOR TARGETED GRAZING

The relationship between grazing animals and plants is similar to predator-prey relationships in which grazing animals are predators and plants are prey. Because plants can't run away from their predators, they use other strategies to resist grazing that Briske (1996) described as grazing avoidance or grazing tolerance. The basic difference between grazing avoidance and grazing tolerance is that avoidance mechanisms reduce the chance of a plant being grazed while tolerance mechanisms allow plants to grow more rapidly after defoliation (Briske, 1996). Understanding grazing resistance mechanisms in a target species is critical to applying targeted grazing. To achieve their landscape objectives, targeted grazers must alter the competitive interactions among plant species to favor desired plants at the expense of the undesired.




Figure 1. Annual frequency of the use of conservation grazing, prescribed grazing, or targeted grazing in peer-reviewed journal articles as determined by the literature using Web of Science. The yr 2006 includes all articles before that date with the earliest citation occurring in 1991. Supporting citations are listed in Supplemental Information 1.

Grazing avoidance is the capacity for a plant to avoid being grazed and involves mechanisms that reduce the probability and/or severity of grazing (Briske, 1996). These defense mechanisms can be divided into plant architectural attributes, mechanical deterrents, and biochemical compounds. Plant secondary compounds are biochemical compounds often used by plants to avoid herbivory (Bennett and Wallsgrove, 1994). Although recent literature has questioned the link between secondary compounds and plant defense, there is still strong evidence that secondary compounds play a major role in grazing resistance (Agrawal and Weber, 2015). Plant architectural attributes, such as plant height (Detling and Painter, 1983), as well as mechanical deterrents, such as spines or thorns (Karban et al., 1999) can reduce the probability of grazing. Finally, plant nutritive value characteristics including lignin and neutral

detergent fiber can limit digestibility and reduce intake, thereby reducing the probability of being grazed.

Compensatory growth after defoliation is an often-cited attribute contributing to grazing tolerance (Briske, 1996), but other traits such as resource allocation patterns (Fornoni, 2011), meristematic number (Briske, 1996), and a high sheath-stem ratio (Cullen et al., 2006) are also important. Expression of grazing tolerance can be driven by both genetics and the environment (Damhoureyeh and Hartnett, 2002). Although presented separately here, many plant species may have developed a mixed avoidance-tolerance response to defoliation (Zheng et al., 2015).

Timing of defoliation (i.e., phenological stage) is an important factor that affects plant response. For example, mowing in the fall, when spotted knapweed (*Centaurea stoebe* L.) was flowering or in seed set, reduced its tiller densities more than mowing in spring when moisture and temperature conditions were more favorable for spotted knapweed regrowth (Rinella et al., 2001). Annual plants need to be grazed in ways to reduce their seed output. Hempy-Mayer and Pyke (2008) found that grazing treatments which defoliated cheatgrass multiple times before seeds began to mature would have the best potential for impacting seed production and reducing cheatgrass populations in successive years. Determining the most appropriate phenological stage to graze requires grazers to understand how target plants reproduce and when they are most susceptible to grazing.

Another complicating factor is the relative palatability differences between the desired and target plants. Plant palatability often changes over the course of the growing season because of advancing leaf age and senescence (Anderson, 1985; Hendrickson et al., 1997), which can shift the relative preference from maturing targeted plants to species growing intermixed with them.

Clonal plants, or plants that spread through roots and rhizomes, can be especially challenging for targeted grazers. Clonal grasses, for example, often have tiller populations containing a variety of growth stages (Hendrickson et al., 1998). The variety of phenological stages makes it difficult to graze when target plants are most vulnerable. Resource sharing among subunits of clonal plants can also propagate new growing points (Derner et al., 2012), making it difficult for a single grazing event to damage the target plant. Because clonal plants can share resources, they often have

Table 1
Application of grazing management principles (Valentine, 2001) to targeted grazing and traditional grazing.

Principle	Targeted Grazing	Traditional Grazing
Stocking rate	Designed heavy use of target plants may result in moderate to heavy use of non-target plants	Low to moderate levels to help ensure plant vigor
Season of use	Timing is often a critical factor <ul style="list-style-type: none"> - Graze when plant secondary metabolites are low to improve palatability and/or reduce adverse post-ingestive feedback - Graze before seed set to reduce seed production - Graze when non-target plants are less palatable and/or more resistant to grazing 	Timing is an important factor <ul style="list-style-type: none"> - Graze when plants are dormant to improve vigor - Use pasture rotation to avoid grazing at the same time each year - Limit grazing period to avoid multiple defoliations of the same plant without rest
Distribution	Livestock are usually focused with fencing or herding at high stocking densities to target vegetation.	Livestock are often dispersed using attractants to avoid heavy use and to encourage even utilization or they are rotated among pastures to avoid prolonged heavy grazing
Species, type and kind of livestock	Livestock are selected based on their preference for targeted vegetation	Livestock are usually selected based on market factors and susceptibility to predation

advantages in heterogeneous environments, such as grazed ecosystems (Hutchings, 1999).

SHAPING AND USING ANIMAL PREFERENCE TO TARGET PLANTS

Success of targeted grazing is dependent on the use of adapted livestock. Ruminants have been classified into three groups on the basis of body size, mouth structure, and digestive capabilities: grazers, browsers, and intermediate feeders (Hofmann, 1989). These differences make some livestock species more capable of ingesting and digesting various vegetation types. For example, goats are browsers and have small, narrow mouths that readily consume leaves and small twigs of shrubs (Shipley, 1999). Goats also have relatively large livers based on their body size, which allows them to process secondary compounds more efficiently than other livestock species (Hofmann, 1989). In contrast, cattle are roughage eaters and have relatively large, broad mouths and greater rumen size than other livestock species, which allows them to use grasses efficiently. Although morphological and physiological differences exist between livestock species, early experiences with forages can modify both morphological and physiological characteristics of livestock's digestive systems so that they are more adapted to efficiently consume forages they were exposed to at a young age (Distel and Provenza, 1991).

Individuals vary greatly in their ability to consume and digest plants that contain large quantities of secondary compounds. Part of this variation is the inherent ability of animals to tolerate and/or process toxins, which may be inherited (Launchbaugh et al., 1999). Another determinant of diet selection is learning and experience. The palatability of forages is a result of postingestive feedback (Provenza, 1995). Livestock prefer forages that provide needed nutrients and do not contain excessive toxins and avoid plants with few nutrients and/or excessive toxin levels. Postingestive feedback is not a conscious decision, but rather a consequence of automatic processing of taste-feedback interactions (Provenza, 1996). Many weeds and invasive plants contain high levels of secondary compounds, which can limit intake of these species unless management interventions are applied (Provenza and Papachristou, 2009).

Lambs, kids, and calves learn about foods from their mother. Young animals are exposed to flavors in utero and through the milk, which can increase their likelihood of consuming the foods that their mother ate (Nolte and Provenza, 1992; Nolte et al., 1992). Exposing young livestock to the forages that managers desire to target should increase intake of those plants later in life (Wiedmeier et al., 2002). Although learning to eat foods from the mother is usually the most effective, young livestock can also learn what to eat from their peers. For example, heifers that were averted from eating larkspur (*Delphinium* spp.) started consuming this poisonous plant when they grazed with heifers that were not averted (Lane et al., 1990). In addition to learning which foods to consume, young livestock must learn how to forage. Young goats with experience eating blackbrush (*Coleogyne ramosissima* Torr.) not only consumed blackbrush more efficiently than inexperienced animals but also consumed live oak (*Quercus turbinella* Greene) more effectively (Ortega-Reyes and Provenza, 1993a, 1993b). In most cases, targeted grazers should consider raising their own female replacement animals and expose their young livestock to the forages they plan to target.

Diet training (a.k.a., diet conditioning) is another strategy that may increase livestock consumption of target plants. Diet training increased use of leafy spurge by sheep (Walker et al., 1992) and use of redberry juniper (*Juniperus pinchotii* Sudw.) by goats (Dietz et al., 2010). However, diet training did not increase cattle use of either broom snakeweed (*Gutierrezia sarothrae* [Pursh] Britton & Rusby; Ralphs and Wiedmeier, 2004) or spotted knapweed (Mosley et al., 2017b).

Supplementation with activated charcoal or polyethylene glycol (PEG) has increased livestock use of target plants. Supplementation of activated charcoal increased sheep use of big sagebrush (*Artemisia*

tridentata Nutt.; Banner et al., 2000; Villalba et al., 2002) and bitterweed (*Hymenoxys odorata* DC.; Poage et al., 2000); however, activated charcoal had little to no effect on goat use of redberry juniper or Ashe juniper (*Juniperus ashei* Buchholz; Bisson et al., 2001). To date, supplementation with activated charcoal is impractical for range livestock due to difficulty finding an appropriate medium for supplementation (Poage et al., 2000). Supplementation with PEG increased goat use of one-seed juniper by sheep and goats (Utsumi et al., 2013), but PEG supplementation on rangelands is currently cost-prohibitive (Ben Salem et al., 1999). Supplemental protein and energy increased sagebrush browsing by sheep (Villalba et al., 2002; Dziba et al., 2007; Guttery, 2011), and supplemental protein increased intake of one-seed juniper by goats (Utsumi et al., 2013). However, supplemental protein-energy did not increase sheep intake of sulfur cinquefoil (Mosley et al., 2017a). Protein-supplemented cattle readily browsed big sagebrush when targeted cattle grazing was applied at a high stock density within small, 0.2-ha experimental pastures (Petersen et al., 2014) but not when targeted cattle grazing was applied at a low stock density within large, 625-ha pastures (Payne, 2018; Payne et al., 2018).

CHANGING ANIMAL USE OF LANDSCAPES

In many cases, one of the keys for effective targeted grazing is to focus the livestock in a specific area where vegetation treatments are needed. Small enclosures facilitate focusing livestock use (Kott et al., 2006). For example, managers often use temporary electric fence to construct small enclosures around patches of noxious weeds or invasive plants, which limits livestock's ability to graze nontarget plants. Small enclosures are also useful for direct defoliation in specific areas to produce fuel breaks, reducing vegetation levels under power lines, defoliating salt cedar (*Tamarix* spp.) along riparian areas, and similar treatments. Temporary fencing can be cost-effective and feasible depending on the availability of stock water. However, it may not always be feasible to construct small pens with livestock water, especially in mountainous and arid areas. In addition, enclosures should provide a variety of forages to keep animals from overingesting toxins.

Livestock grazing can be focused without fencing. Attractants can be used to lure livestock to desired areas (Bailey, 2004). Low-moisture block and other protein supplements are effective tools to attract cattle to underutilized areas (Martin and Ward, 1973; Bailey and Welling, 1999). For example, low-moisture block protein supplement was used effectively to concentrate cattle trampling and reduce dense sagebrush cover in southwestern Montana (Payne, 2018; Payne et al., 2018). Salt and mineral mixes can be effective tools to attract cattle when forages are green and growing (Pittarello et al., 2016), but protein-based low-moisture block supplements are more effective than salt when the forage is dormant (Bailey et al., 2008). Cattle grazing can be focused without fences using a combination of low-stress stockmanship and strategic supplement placement. This combination of distribution practices focused cattle grazing in underutilized areas of pastures > 2 km from water and in rugged terrain in New Mexico (Pollak, 2007; Stephenson et al., 2016) and in Arizona (Bruegger et al., 2016). Herding can also work well for focusing sheep grazing (Kott et al., 2006). Typically, whiteface sheep breeds are more suited to herding because they are usually more gregarious and have a stronger flocking instinct than blackface breeds (Olson and Launchbaugh, 2006).

High stocking densities can be created by placing relatively large numbers of livestock in small paddocks. High stocking densities may be helpful in ensuring good soil seed contact in rangeland restoration efforts (Winkel and Roundy, 1991; Mosley and Roselle, 2006), increasing mortality of undesirable shrubs through browsing and trampling (Marquardt et al., 2009), and modifying shrub structure and growth (Ganskopp et al., 2004). When livestock are spatially concentrated (high stocking densities on areas near water sources or supplements), animals can mechanically damage vegetation and impact soil structure through trampling (Warren et al., 1986; Abdel-Magid et al., 1987). Soil

bulk density increases and water holding capacity and infiltration decreases after trampling, but soil structure naturally recovers after livestock exclusion (i.e., rest from grazing) through freeze–thaw and wetting and drying cycles (Drewry, 2006). Ultrahigh stocking density or “mob grazing” requires > 200 000 kg of cattle liveweight ha⁻¹ and has been purported to increase harvest efficiency, plant productivity, plant species diversity and soil carbon (Gompert, 2010). Ultrahigh stocking densities can result in two to three times the extent of trampling compared with four pasture rotation systems (Johnson, 2012). An 8-yr study comparing mob grazing and four pasture rotation systems in Nebraska meadows did not detect any improvement in harvest efficiency, steer gain, plant productivity, species composition, ground cover, soil bulk density, and soil carbon (Johnson, 2012; Lindsey, 2016; Shropshire, 2018). Although the Nebraska studies did not support the purported benefits of ultrahigh stocking densities, further studies in other locations and vegetation types are needed.

OBSTACLES TO ADOPTION AND EFFECTIVE USE OF TARGETED GRAZING

Plant Secondary Chemistry

Plant secondary metabolites (PSMs) are a major factor driving diet selection and preferences of domestic livestock and wildlife (Kronberg and Walker, 2007). Encroaching species targeted for removal are often shrubs that maintain a competitive edge because of their PSM content. Thousands of PSM exist, representing several structural classes, and their presence and concentration in a given plant are influenced by many factors, including genetics, phenology, and a host of biotic and abiotic environmental factors. Numerous physiological (e.g., season, plant age, leaf age, plant part, regrowth) and environmental (soil properties, moisture, nutrient availability, competition, temperature, humidity, light, herbivory, microbial attack, mechanical damage, CO₂, ozone) factors affect PSM concentration (Kuokkanen et al., 2003; Llusà et al., 2006; Thines et al., 2007). Adding to this complexity is the fact that PSM concentrations vary from plant to plant, day to day, within day/diurnal cycles, and within plant (Heyworth et al., 1998; Komenda and Koppmann, 2002). Furthermore, secondary compounds can be present constitutively or induced in response to stress (e.g., herbivory or drought; Holopainen, 2004). Induced responses can occur rapidly in response to biotic stresses such as herbivory (i.e., within minutes, hours, or a few days; Kost and Heil, 2006) and are often short-lived (Faeth, 1992). Plant volatiles can even signal neighboring plants to increase defense levels before damage (Kost and Heil, 2006). Thus, for a given targeted species, the number of permutations of PSM and nutrient profiles and concentrations are infinitely large in time and space. At a larger scale, chemical variability may be driven by factors such as ecological site (i.e., soil characteristics), induction due to localized outbreaks of insects or peak populations of small mammals, and localized water pulses. This variability in PSM poses serious challenges to developing grazing prescriptions that are consistently effective. The more information available about the concentrations of these chemicals, the greater the chance of successfully implementing a targeted grazing program.

Various management options are available to increase intake of PSM-laden plants. Approaches that rely on animal genetics (e.g., selection for browsers), behavior (e.g., diet training), nutrition (e.g., supplementation, additives), and management (e.g., manipulating timing of browsing, plane of nutrition, body condition, stocking rate) (Shaw et al., 2006; Dziba et al., 2007; Papachristou et al., 2007; Frost et al., 2008; Rogosic et al., 2008; Utsumi et al., 2009; Waldron et al., 2009; Campbell et al., 2010; Utsumi et al., 2010, 2013) can be incorporated into targeted grazing programs. However, the effectiveness of these practices is influenced by variations in PSM concentrations. Conversely, understanding this variability can help to identify times and locations when a target plant is particularly vulnerable. For example, Utsumi et al.

(2009) observed intake of one-seed juniper by small ruminants in the fall when PSM concentrations were highest was about half that during summer, winter, or spring. In that same study, protein supplementation increased juniper consumption by approximately twofold; however, during the fall when terpenoid concentrations were greatest, effects of protein were negligible. In a follow-up study under field conditions, Utsumi et al. (2010) observed that browsing intensity by small ruminants was greater during summer than spring and for short versus medium or tall saplings. Total terpenoid concentrations in juniper were inversely related to defoliation intensity and were lower in summer versus spring and in short versus medium or tall saplings (Estell et al., 2014). Consequently, targeted browsing of short juniper saplings during summer may be more effective than in the fall.

Seed Dispersal or Endozoochory

Endozoochory (i.e., dispersal of seeds by animals after passage through the animals' digestive system) is a valid concern when using targeted livestock grazing to suppress undesired plants. Although livestock that consume immature seed-heads of plants do not ingest viable seeds, livestock sometimes do ingest viable seeds if they consume mature seed-heads. Exposure of ingested seeds to the digestive tract of livestock significantly reduces their viability, but livestock can excrete viable seeds in their feces (Lacey et al., 1992; Wallander et al., 1995; Olson and Wallander, 2002; Frost et al., 2013). If livestock consume viable seeds of invasive or otherwise undesired plants, they should be confined in a corral before they are moved to a new area (Kott et al., 2006). Several studies indicate that livestock should remain confined for 3 or 4 d to provide sufficient time for them to excrete undesirable plant seeds (Lacey et al., 1992; Wallander et al., 1995; Olson and Wallander, 2002; Frost et al., 2013). An exception is halogeton (*Halogeton glomeratus* [M. Bieb] C.A. Mey.). Lehrer and Tisdale (1956) documented that most viable seeds of halogeton were excreted by the fourth day after consumption, but a few viable halogeton seeds were not excreted until 9 d post consumption. On the other hand, Goodman et al. (2014) reported zero germination of white locoweed (*Oxytropis sericea* Nutt.) seeds recovered from fecal pellets of sheep that had grazed infested plots for 5 d during the seed set stage.

Animal Welfare

Targeted grazing is sometimes accomplished via transient confining of animals in relatively small, fenced grazing areas (Goodman et al., 2014; Utsumi et al., 2010) or herding (Bruegger et al., 2016; Stephenson et al., 2016) to induce consumption of target (unwanted) plants that frequently contain significant concentrations of varied PSMs (see earlier). Transient confinement and, perhaps more importantly, consumption of potentially toxic feeds are management practices likely to raise concerns about animal welfare. Management of animals in targeted grazing programs usually involves modifying the expression of “natural” behaviors (Kilgour, 2012; Špinka, 2006) which are considered essential to animal welfare (Bracke and Hopster, 2006) and are a central tenet of organic animal farming (Vaarst and Alrøe, 2012). Perceived suppression of natural behaviors could pose challenges to the broad adoption of this technique, particularly among implementers seeking more natural means of managing rangeland vegetation. Ingestion of target plants containing varying levels of toxins, as is often the case in targeted grazing programs, can induce transient to chronic animal malaise (Foley et al., 1999; Estell, 2010), which, even if subclinical, could be perceived as causing unnecessary animal distress. Potential implementers of targeted grazing programs might be deterred by the risk of jeopardizing animal well-being.

Although the focus of most targeted grazing research to date has naturally been placed on measuring the efficacy of this technique to suppress unwanted plant populations (Wallace et al., 2008; Goehring et al., 2010; Utsumi et al., 2010; Mosley et al., 2016; Probo et al., 2016),

researchers have recently begun quantifying the effects of target plant PSMs on animal health (Goodman et al., 2014; Arviv et al., 2016). Goodman et al. (2014), for example, tested two targeted grazing treatments to identify a prescription that would avoid alkaloid intoxication in sheep without compromising the efficacy of the targeted grazing treatment. They reported that by applying intermittent targeted grazing of white locoweed-infested rangeland (5 d on locoweed followed by 3 d off), subclinical symptoms of alkaloid toxicity were significantly reduced relative to sheep that grazed locoweed continuously for 25 d. Interestingly, locoweed suppression efficacy was not different among targeted grazing treatments. Alkaline phosphatase levels in plasma (an indicator of liver activity) remained within normal ranges in ewes in the intermittent targeted grazing treatment, suggesting that this prescription likely spared animals from potential adverse effects of locoweed PSMs.

PLANNING AND MONITORING

The Society for Range Management defines monitoring as “the orderly collection, analysis, and interpretation of resource data to evaluate progress toward meeting management objectives ...” (SRM, 1998). Good monitoring practices and effective use of the information they produce are keys to successfully and consistently accomplishing objectives.

Successfully managing grazing animals and plant communities with targeted grazing requires 1) clear objectives (preferably measurable) for the desired plant community and soil surface characteristics being manipulated; 2) some reasonable estimate of those same characteristics at the start of the project; 3) the right species, class, and number of animals to impact vegetation in the way intended in the time available; and 4) a strategy for managing the animals in a way that will obtain the desired vegetation and/or soil surface effects. Ongoing monitoring provides feedback on the rate and direction of change in the site characteristics. All 5 of those elements (1–4 plus the monitoring plan) should be detailed in a targeted grazing prescription.

Monitoring is important because targeted grazing projects entail many moving parts and, therefore, a great deal of uncertainty. Animal demand; available forage; relative palatability of plants (influenced by season, stage of growth, species composition, residue levels, etc.); elapsed time during the grazing period; weather conditions; vulnerability of nontarget plants; and water quality and availability can all change continuously over time. The timeliness and appropriateness of the manager’s response to variations in these factors will influence the success of the project, and it is the combination of continuous casual observations and appropriate measurements that make such a response possible. This kind of monitoring was a component of what came to be described as “adaptive management” by Holling (1978) and Walters and Hilborn (1978). This is a management approach suited to dealing with uncertainty (i.e., situations where the best practice(s) for the intended objectives can only be approximately known in advance). In this process, management decisions are adjusted iteratively by evaluating information in a continuous feedback loop (Allen et al., 2017). Such management requires a commitment to using the feedback to change management actions and requires a plan for collecting specific feedback and a process for coupling it with management actions (“learning by doing” or “the art of range management”) (Stoddart et al., 1975; Mosley, 1985).

Sharrow and Seefeldt (2006) outlined two basic functions of monitoring in targeted grazing projects: 1) using progress toward objectives to evaluate and refine the grazing prescription (the adaptive management component) and 2) generating documentation of practices and results that may be required under grazing agreements. With respect to the first point, they emphasize the importance of selecting objectives (goals) that are clearly stated, measurable, include a timeline for completion, and are realistically achievable. Proper goal/objective

selection can seem trivial, but it usually is not. For example, “Reduce target plant density” is a simple, clear goal, but by itself is not useful for making management decisions. What’s missing? The size of the desired change, the time allowed for the change to occur, the method(s) for measuring the change, and the prescription to accomplish the change. These comprise the basic elements of a monitoring plan. By identifying responsible personnel and a timeline of events, the monitoring process is complete.

There is no single, straightforward guide that covers the wide range of monitoring protocols that might be useful for the variety of targeted grazing projects that are possible. However, many good materials exist that describe monitoring practices for many common environmental and ecological objectives (Elzinga et al., 1998; Coulloudon et al., 1999; Winward, 2000; Pellant et al., 2005; Herrick et al., 2017). Modifications of these for specific circumstances and applications are continually being created and refined. University Extension programs, agricultural experiment stations, and many conservation organizations have been developing streamlined protocols of particular value for land managers and livestock operators (Lewis et al., 2000; Ward et al., 2003; Mosley et al., 2018). These resources can help livestock and land managers select appropriate monitoring methods for their specific objectives and circumstances. The goal should be to select or develop monitoring methods that can detect changes of the size anticipated and that can be completed in the time available seasonally to do the field work. This difficulty is as real for federal land management agencies as it is for conservation organizations, private landowners, and livestock operators.

APPLICATION OF TARGETED LIVESTOCK GRAZING ON RANGELANDS

It is important to recognize that targeted livestock grazing is not always the most appropriate tool to select from the vegetation management methods options. Other methods may be more practical or appropriate in some situations (Mosley, 1997). Even when targeted livestock grazing is an appropriate option, its effectiveness will depend on the skill of the person applying this technology (Table 2). Those lacking experience with targeted livestock grazing should apply targeted grazing cautiously at first and seek guidance from more experienced practitioners (Mosley, 1997). With these guiding principles in mind, opportunities abound for using targeted livestock grazing to manage rangeland vegetation. Some of the possible applications of targeted grazing are described as follows.

Suppress Undesired Plant Species

To suppress undesired plant species and increase desired ones, targeted livestock grazing should be applied when 1) targeted plants are susceptible to defoliation and relatively palatable and 2) when desired plants are less susceptible to defoliation and relatively less palatable (see Table 2). Clipping studies can efficiently and economically identify the phenotypic stages when targeted plants are susceptible to defoliation. Benzel et al. (2009), for example, hand-clipped spotted knapweed plants at seven different timings and frequencies to identify opportunities for reducing viable seed production by spotted knapweed. Similarly, Frost and Mosley (2012) hand-clipped sulfur cinquefoil (*Potentilla recta* L.) plants at seven timings and two intensities to identify potential defoliation strategies for reducing the yield and viable seed production of sulfur cinquefoil. These clipping studies were then followed with grazing experiments that examined the relative palatability of the targeted weeds and the efficacy of targeted grazing applied during the phenotypic stages identified as most promising by the clipping studies. Results of these experiments indicated that targeted grazing by sheep or cattle can suppress spotted knapweed if targeted grazing is applied when spotted knapweed is in its late bud–early flower phenotypic stage (late July) or full-flower

Table 2
Examples of targeted grazing applications and comparisons to alternative management

Goal of Targeted Grazing	Targeted Grazing Approach	Targeted Grazing Feasibility	Alternative Management Feasibility
Suppress undesired plants			
- Cheatgrass	Grazed when green to reduce seed production	Effective if sufficient livestock are available More flexible than spring grazing and easier to apply	Herbicides are expensive Seeding perennial vegetation is expensive and has low probability of success unless cheatgrass abundance is reduced
- Spotted knapweed	Winter grazing can reduce cheatgrass abundance Grazed sheep or cattle during late bud to flower stage	Can suppress spotted knapweed at a moderate cost	Herbicides are effective but expensive
- One-seed juniper	Browse saplings with goats or goats and sheep	Mortality of browsed saplings is 5 to 15%. Dependent on goat and sheep availability. Not effective on large trees	Mechanical treatments are effective but expensive Herbicides are effective but expensive
Enhance wildlife forage quantity and quality	Grazed cattle or sheep to remove decadent vegetation	Increases use of big game and can be achieved using attractants (e.g., supplements) or minor changes in traditional grazing management	Prescribed fire is effective but has risks and moderate costs
Modify shrub structure and rejuvenate productivity	Lightly graze bitterbrush in the spring with cattle to stimulate twig growth	Cattle grazing and periodic deferment increases height and diameter of bitterbrush plants	Mechanical pruning would be impractical and expensive
Fire breaks	Grazed cattle or sheep to reduce fine fuel loads	Difficult to reduce standing crop below moderate levels without fencing	Mechanical treatments are effective but expensive and may result in erosion Prescribed fire can be effective but has risks and moderate costs

phenotypic stage (mid–August) (Thrift et al., 2008; Surber et al., 2011; Henderson et al., 2012; Mosley et al., 2016). To suppress sulfur cinquefoil, targeted livestock grazing can be applied effectively when sulfur cinquefoil is in either its early flowering stage (late June) or late flowering–early seedset stage (mid–July) (Mosley et al., 2017a). For both spotted knapweed and sulfur cinquefoil, targeted livestock grazing was best applied when the targeted weed remained green during mid to late summer, but desirable grasses and forbs were largely dormant. These conditions limited potential adverse effects to desirable plants and encouraged livestock to consume the targeted weeds. Also, targeted grazing applied during mid to late summer stressed the targeted weeds when moisture was depleted and insufficient for plant recovery (Wooley et al., 2011). In addition, targeted grazing was applied before spotted knapweed or sulfur cinquefoil plants had produced viable seeds.

A targeted grazing prescription for suppressing cheatgrass was synthesized from the results of clipping studies and historical observations (Mosley, 1996). Hulbert (1955) and (Tausch et al., 1994) determined that hand-clipping cheatgrass plants twice during late spring (late April to mid–May) reduced cheatgrass plant density and biomass. Clipping occurred when cheatgrass plants were in the boot stage, before cheatgrass produced viable seeds. A subsequent study by Hempy-Mayer and Pyke (2008) also determined that clipping during the boot stage reduced cheatgrass biomass and reduced viable seed production by cheatgrass. Conclusions from the clipping studies were supported by historical observations of cheatgrass response to sheep grazing (Megee, 1938; Daubenmire, 1940), and a contemporary grazing experiment by Diamond et al. (2009) confirmed that targeted cattle grazing can suppress cheatgrass if targeted grazing is applied when cheatgrass is in the boot stage (typically April or May). In many areas cheatgrass requires a second grazing in spring because it can regrow and produce new viable seeds about 3–4 wk after the first defoliation (Hulbert, 1955). Cheatgrass can be suppressed by targeted livestock grazing that significantly limits its production of viable seeds for 2 or 3 consecutive yr (Daubenmire, 1940; Diamond et al., 2009). Targeted sheep or cattle grazing during late fall or winter also can be used to reduce the biomass of cheatgrass mulch, helping to suppress cheatgrass and enhancing establishment of perennial plants (Hull and Pechanec, 1947; Schmelzer et al., 2014).

Enhance Wildlife Forage Quantity and Quality

Forage quantity and quality often determine whether rangeland wildlife live or die. Targeted livestock grazing, browsing, and trampling can be used to purposely enhance forage yield, accessibility, and nutritive quality for many wildlife species (Mosley, 1994; Severson and Urness, 1994; Mosley and Brewer, 2006). For example, targeted livestock grazing, browsing, and trampling can be used to create preferred foraging habitat for northern bobwhite quail (*Colinus virginianus*) (Hernández and Guthery, 2012) and greater sage-grouse (*Centrocercus urophasianus*) (Crawford et al., 2004; Guttery, 2011; Payne, 2018; Payne et al., 2018). Targeted sheep browsing can increase the abundance and accessibility of nutritious winter browse for mule deer (*Odocoileus hemionus*), white-tailed deer (*Odocoileus virginianus*), elk (*Cervus elaphus*), and moose (*Alces alces*) (Jensen et al., 1972; Alpe et al., 1999), and targeted grazing by sheep or cattle can improve accessibility to nutritious herbaceous forage for mule deer, pronghorns (*Antilocapra americana*), and elk (Smith et al., 1979; Willms et al., 1979; Clark et al., 2000; Short and Knight, 2003; Pollak, 2007). Crane et al. (2016) demonstrated on a large landscape scale that elk in spring avoided foraging in areas not grazed by cattle during the previous summer – early fall. Rather, elk preferred to graze where cattle had grazed lightly (11–30% forage use) or moderately (31–60% forage use), and preference by elk was stronger for moderately grazed sites. Both moderate and light cattle grazing intensity had more influence on elk foraging site selection than distance to security cover, distance to roads, aspect, or slope.

Targeted grazing can also be used to modify shrub structure and increase productivity. In the absence of grazing, bitterbrush (*Purshia tridentata* [Pursh] DC) productivity can decline by 70% (Tueller and Tower, 1979). Cattle grazing can modify the structure of bitterbrush shrubs (Ganskopp et al., 1999), and light cattle grazing in the spring can rejuvenate plants and improve productivity (Ganskopp et al., 2004).

Management of vegetation diversity is critical for wildlife habitat management. For example, grassland birds inhabit a gradient of vegetative structure from heavily grazed grasslands to undisturbed tall vegetation (Derner et al., 2009). For instance, Collins et al. (1998) demonstrated that vegetation diversity of tall grass prairie can be maintained through grazing. Livestock grazing is also critical for maintaining vegetation diversity of ephemeral wetlands (Pyke and Marty, 2005). Vegetation diversity of vernal pools in California

declined after 10 yr when grazing was excluded (Marty, 2015). Targeted grazing can be designed to specifically address the needs of individual species or develop diverse habitat conditions. In locations with appreciable forage regrowth, continuous grazing at low stocking densities and stocking rates can be used to create patches and diversify vegetation structure (Cid and Brizuela, 1998; Teague and Dowhower, 2003; Bailey and Brown, 2011). Supplements can be used to temporally create areas of heavy grazing and low vegetation levels near placement sites (Bailey and Welling, 1999). Distribution management tools such as fencing and herding, discussed earlier, can also be used to increase or decrease forage defoliation in desired areas to enhance diversity.

Fire Breaks

By focusing livestock in specific areas, managers can use targeted grazing to produce fire breaks to reduce the fuel continuity and the potential for catastrophic wildfire (see Table 2). Controlled and repeated grazing during the growing season can be used to reduce standing crop and regrowth to create firebreak strips (Taylor, 2006). Diamond et al. (2009) demonstrated that targeted cattle grazing could reduce fuel loads of cheatgrass sufficiently to reduce flame lengths and fire spread. Similarly, Bruegger et al. (2016) showed that cattle targeted grazing using low-stress herding and strategic supplement placement could reduce fuel loads sufficiently to reduce flame lengths and fire spread (Varelas, 2012). The combination of targeted grazing and prescribed fire can be an effective tool to produce firebreaks (Taylor, 2006; Diamond et al., 2012).

Integrated Management

Targeted livestock grazing can be integrated with other vegetation management options, including disking and seeding, herbicides, biological control insects, and prescribed fire. For example, cheatgrass seeds often germinate after disking, burning, or herbicide applications to control broad-leaved weeds. Targeted livestock grazing can defoliate the newly established cheatgrass plants and prevent them from producing viable seed (Mosley et al., 1999; Mosley and Roselle, 2006). After targeted livestock grazing is applied, sites can be drill- or broadcast-seeded. Another option is to broadcast seed immediately before the targeted grazing treatment and use the livestock to trample the seed into the ground (Havstad, 1994; Mosley et al., 1999), preferably with a high livestock density for a brief period when soils are moist (Mosley and Roselle, 2006).

Herbicides combined with targeted livestock grazing often suppress targeted plant species more than herbicides or targeted livestock grazing alone. To suppress leafy spurge, picloram plus 2,4-D can be applied in the fall after applying targeted livestock grazing during the summer (Lym et al., 1997). For spotted knapweed, 2,4-D can be applied in the spring to remove adult spotted knapweed plants, followed by targeted livestock grazing during summer to suppress seedling and juvenile spotted knapweed plants (Sheley et al., 2004).

Targeted livestock grazing and biological control insects complement each other, together stressing targeted plants more than either method applied alone. For example, after 4 yr of treatment, spotted knapweed plant density was 86% less in areas treated with biological control insects plus targeted sheep grazing applied during late July, versus areas treated with biological control insects alone (Mosley et al., 2016). Furthermore, combined herbivory by targeted sheep grazing and biological control insects reduced adult plant density and prevented compensatory recruitment of spotted knapweed seedlings and juvenile plants, but treatment with biological control insects alone did not. Targeted sheep grazing was timed to be compatible with the life cycles of the biological control insects.

Prescribed fire can be integrated with targeted livestock grazing to either suppress invasive plants or to alter the structural and spatial heterogeneity of vegetation to benefit wildlife. Diamond et al. (2012)

combined targeted cattle grazing in May with prescribed burning in October to suppress cheatgrass. The combined targeted grazing—prescribed fire treatment was more effective than either treatment applied alone. Patch-burn grazing is another way that prescribed fire can be combined with pyric herbivory. Patch-burn grazing divides pastures into several patches using firebreaks but no cross-fences. One or two patches in each pasture are burned on a rotational basis each year, and livestock are allowed to move freely among the patches in a pasture. The most recently burned patch typically receives the greatest grazing pressure, whereas reduced grazing in unburned areas leads to greater vegetative cover. Thus, patch-burn grazing creates a mosaic of heterogeneous vegetation structure and cover (Fuhlendorf and Engle, 2004), which provides preferred habitat for many grassland birds such as mountain plover (*Charadrius montanus*), long-billed curlew (*Numenius americanus*), lesser prairie chicken (*Tympanuchus pallidicinctus*), greater prairie-chicken (*Tympanuchus cupido*), and upland sandpiper (*Bartramia longicauda*) (Derner et al., 2009; Sandercock et al., 2015; Winder et al., 2017). Patch-burn grazing also increases grassland butterfly diversity (Delaney et al., 2016).

FUTURE RESEARCH

Livestock producers, land managers, and entrepreneurs have been using the principles described in the Launchbaugh and Walker (2006) handbook for > 10 yr to apply targeted grazing. However, ongoing research is improving and refining targeted grazing practices (Figure 2). Improving managers' ability to concentrate livestock into designated areas will improve the efficacy of targeted grazing. Low-stress stockmanship developed by Bud Williams and described by Hibbard (2012) may be one potential method to focus livestock grazing, and virtual fencing (Anderson et al., 2014) is a promising technique to constrain livestock. Our understanding of secondary compounds is expanding and allowing practitioners to develop more effective targeted grazing protocols (Estell, 2010). However, the priorities for future research are animal welfare, genetic selection, and supplementation. The impact of targeted grazing on animal well-being is a critical issue that researchers have not fully considered. Consumers and other stakeholders are concerned with animal welfare (Verbeke, 2009; Lagerkvist and Hess, 2010), and some applications of targeted grazing may potentially impact animal well-being. Diet preference in sheep (Snowder et al., 2001) and goats (Waldron et al., 2009) is heritable for sagebrush and juniper, respectively, and for the latter both a physiological (Campbell et al., 2010) and genomic (Walker et al., 2019) basis has been demonstrated. The other important and exciting research area for targeted grazing is the use of supplements and pharmaceutical enhancements to minimize adverse impacts of secondary compounds on livestock. Additional research on these priority topics has great promise to increase the application of targeted grazing in the future.

Animal Welfare

Addressing potential animal welfare issues should become a priority in targeted grazing studies, particularly in those cases where the target plant species contains high concentrations of PSMs or where natural animal behavior is severely altered. Implementing targeted grazing programs at times when PSM concentrations are lowest should not only enhance chance of success (i.e., increase consumption of targeted species) but also minimize negative effects on animal well-being. The research community could anticipate obstacles to adoption of this technique by routinely addressing animal well-being in all targeted grazing studies.

Genetic Selection

A fundamental principle of targeted livestock grazing is to match the vegetation management task with the appropriate




Figure 2. Grazing management tools (stocking rate, timing of grazing, species of herbivore, and distribution; Vallentine, 2001) used to achieve targeted grazing goals (change plant species composition, remove plant biomass, improve forage quality, and mechanical impacts) and ongoing research to improve the efficacy of these tools. Arrows show approaches to improve the efficacy of the grazing management tools and where additional research may be helpful.

animal. Finding a good animal match is relatively easy when targeted plant species are highly palatable and located on gentle terrain. However, few good matches may be available if the vegetation management task requires livestock to consume phytotoxin-containing plants or plants growing in rugged terrain (see Figure 2). Selective breeding has great potential to mitigate phytotoxicosis in range livestock (Launchbaugh et al., 1999; Campbell et al., 2010; Estell et al., 2012, 2014), and selective breeding also may be able to increase rugged terrain use by livestock (Bailey et al., 2015). Previous studies confirm the practicality and relevance of pursuing selective breeding. For example, resistance to ergot alkaloids that cause fescue toxicosis was developed by selective breeding for hepatic biotransformation enzymes in sheep (Morris et al., 1989, 1995) and cattle (Morris et al., 1998). Snowden et al. (2001) and Waldron et al. (2009) demonstrated heritable differences among individual sheep and goats in their ability to cope with phytotoxins in big sagebrush and Ashe juniper, respectively, and Cook et al. (2011) identified interindividual variation to larkspur toxicosis in cattle. Furthermore, Walker et al. (2019) found genetic markers for juniper consumption in goats divergently selected for increased or decreased juniper consumption. Ultimately it may be possible for livestock producers to identify suitable animals by submitting animal blood or hair samples for genetic testing.

Supplements and Pharmaceutical Enhancements

Several research projects have identified the value of dietary supplement to enhance detoxification and increase consumption of a targeted plant. For example, as mentioned earlier, feeding supplemental protein and polyethylene glycol may increase intake of oneseed juniper by goats (Utsumi et al., 2013) and supplementing sheep with barley and activated charcoal increased intake of big sagebrush (Banner et al., 2000). As we learn more about PSMs and physiological mechanisms of detoxification, it is likely that specific supplements will be designed to entice animals to consume target plants. In this way, individual animals will be able to consume greater amounts of target plants and fewer animals will be needed for targeted grazing projects, thereby increasing the economic and management feasibility of targeted grazing.

CONCLUSIONS

Targeted livestock grazing differs from other grazing management schemes in that its objective is to accomplish defined land management goals rather than livestock productivity. Managers use a specific kind of livestock at defined timing, intensity, duration, and location of grazing to resolve specific vegetative issues as an alternative or in combination with other practices such as herbicide applications,

mechanical treatments, or prescribed burning. Like other management, targeted grazing should be approached as an adaptive process, where applications are monitored and evaluated so that the techniques can be improved and refined. Targeted grazing must overcome challenges associated with secondary compounds, endozoochory, and animal welfare. However, ongoing and future research is addressing issues that limit application of targeted grazing. Targeted livestock grazing is a valuable option in land managers' tool boxes.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.rama.2019.06.003>.

Acknowledgments

The development of this paper was an activity of the Targeted Grazing Committee of the Society for Range Management.

References

- Abdel-Magid, A.H., Trlica, M., Hart, R.H., 1987. Soil and vegetation responses to simulated trampling. *Journal of Range Management* 40, 303–306.
- Agrawal, A.A., Weber, M.G., 2015. On the study of plant defence and herbivory using comparative approaches: how important are secondary plant compounds. *Ecology Letters* 18, 985–991.
- Allen, C.R., Angeler, D.G., Fontaine, J.J., Garmestani, A.S., Hart, N.M., Pope, K.L., Twidwell, D., 2017. Adaptive management of rangeland systems. In: Briske, D. (Ed.), *Rangeland systems*. Springer Nature, Cham, Switzerland, pp. 373–394.
- Alpe, M.J., Kingery, J.L., Mosley, J.C., 1999. Effects of summer sheep grazing on browse nutritive quality in autumn and winter. *Journal of Wildlife Management* 63, 346–354.
- Anderson, B., 1985. The influence of aging on forage quality of individual switchgrass leaves and stems. In: 15th International Grassland Congress; Nishinasuno, Tochigiken. National Grassland Research Institute, Japan, pp. 947–949.
- Anderson, D.M., Estell, R.E., Holecchek, J.L., Ivey, S., Smith, G.B., 2014. Virtual herding for flexible livestock management review. *The Rangeland Journal* 36, 205–221.
- Arviv, A., Muklada, H., Kigel, J., Voet, H., Glasser, T., Dvash, L., Ungar, E., Landau, S., 2016. Targeted grazing of milk thistle (*Silybum marianum*) and Syrian thistle (*Notobasis syriaca*) by goats: preference following preconditioning, generational transfer, and toxicity. *Applied Animal Behaviour Science* 179, 53–59.
- Bailey, D.W., 2004. Management strategies for optimal grazing distribution and use of arid rangelands. *Journal of Animal Science* 82, E147–E153.
- Bailey, D.W., Brown, J.R., 2011. Rotational grazing systems and livestock grazing behavior in shrub-dominated semi-arid and arid rangelands. *Rangeland Ecology & Management* 64, 1–9.
- Bailey, D.W., Lunt, S., Lipka, A., Thomas, M.G., Medrano, J.F., Cánovas, A., Rincon, G., Stephenson, M.B., Jensen, D., 2015. Genetic influences on cattle grazing distribution: association of genetic markers with terrain use in cattle. *Rangeland Ecology & Management* 68, 142–149.
- Bailey, D.W., VanWagoner, H.C., Weinmeister, R., Jensen, D., 2008. Comparison of low-moisture blocks and salt for manipulating grazing patterns of beef cows. *Journal of Animal Science* 86, 1271–1277.
- Bailey, D.W., Welling, G.R., 1999. Modification of cattle grazing distribution with dehydrated molasses supplement. *Journal of Range Management* 52, 575–582.
- Banner, R.E., Rogosic, J., Burritt, E.A., Provenza, F.D., 2000. Supplemental barley and charcoal increase intake of sagebrush by lambs. *Journal of Range Management* 53, 415–420.
- Ben Salem, H., Nefzaoui, A., Ben Salem, L., Tisserand, J.L., 1999. Different means of administering polyethylene glycol to sheep: effect on the nutritive value of *Acacia cyanophylla* Lindl. foliage. *Animal Science* 68, 809–818.
- Bennett, R., Wallsgrove, R., 1994. Secondary metabolites in plant defence mechanisms. *New Phytologist* 127, 617–633.
- Benzel, K.R., Mosley, T.K., Mosley, J.C., 2009. Defoliation timing effects on spotted knapweed seed production and viability. *Rangeland Ecology & Management* 62, 550–556.
- Bisson, M.G., Scott, C.B., Taylor Jr., C.A., 2001. Activated charcoal and experience affect intake of juniper by goats. *Journal of Range Management* 54, 274–278.
- Bracke, M.B., Hopster, H., 2006. Assessing the importance of natural behavior for animal welfare. *Journal of Agricultural and Environmental Ethics* 19, 77–89.
- Briske, D., 1996. Strategies of plant survival in grazed systems: a functional interpretation. In: Hodgson, J., Illius, A.W. (Eds.), *Ecology and management of grazing systems*. CAB International, Wallingford, UK, pp. 37–67.
- Briske, D.D., Derner, J.D., Milchunas, D.G., Tate, K.W., 2011. An evidence-based assessment of prescribed grazing practices. Conservation benefits of rangeland practices: assessment, recommendations, and knowledge gaps. USDA-NRCS, Washington, DC, USA, pp. 21–74.
- Bruegger, R.A., Varelas, L.A., Howery, L.D., Torell, L.A., Stephenson, M.B., Bailey, D.W., 2016. Targeted grazing in southern Arizona: using cattle to reduce fine fuel loads. *Rangeland Ecology & Management* 69, 43–51.
- Butler, L., Cropper, J., Johnson, R., Norman, A., Peacock, G., Shaver, P., Spaeth, K., 2003. National range and pasture handbook. USDA National Resources Conservation Service, Washington, DC, USA, p. 214.
- Campbell, E., Frost, R., Mosley, T., Mosley, J., Lupton, C., Taylor, C., Walker, J., Waldron, D., Musser, J., 2010. Pharmacokinetic differences in exposure to camphor after intraruminal dosing in selectively bred lines of goats. *Journal of Animal Science* 88, 2620–2626.
- Cid, M.S., Brizuela, M.A., 1998. Heterogeneity in tall fescue pastures created and sustained by cattle grazing. *Journal of Range Management* 51, 644–649.
- Clark, P.E., Krueger, W.C., Bryant, L.D., Thomas, D.R., 2000. Livestock grazing effects on forage quality of elk winter range. *Journal of Range Management* 53, 97–105.
- Collins, S.L., Knapp, A.K., Briggs, J.M., Blair, J.M., Steinauer, E.M., 1998. Modulation of diversity by grazing and mowing in native tallgrass prairie. *Science* 280, 745–747.
- Cook, C.W., Harris, L.E., 1952. Nutritive value of cheatgrass and crested wheatgrass on spring ranges of Utah. *Journal of Range Management* 5, 331–337.
- Cook, D., Green, B.T., Welch, K.D., Gardner, D.R., Pfister, J.A., Panter, K.E., 2011. Comparison of the toxic effects of two duncecap larkspur (*Delphinium occidentale*) chemotypes in mice and cattle. *American Journal of Veterinary Research* 72, 706–714.
- Coulloudon, B., Eshelman, K., Gianola, J., Habich, N., Hughes, L., Johnson, C., Pellant, M., Podborny, P., Rasmussen, A., Robles, B., Shaver, P., Spehar, J., Willoughby, J., 1999. Utilization studies and residual measurements: interagency technical reference, second revision. U.S. Department of the Interior, Bureau of Land Management Technical Reference 1734-3, Denver, CO, USA, 165 p.
- Crane, K.K., Mosley, J.C., Mosley, T.K., Frost, R.A., Smith, M.A., Fuller, W.L., Tess, M.W., 2016. Elk foraging site selection on foothill and mountain rangeland in spring. *Rangeland Ecology & Management* 69, 319–325.
- Crawford, J.A., Olson, R.A., West, N.E., Mosley, J.C., Schroeder, M.A., Whitson, T.D., Miller, R.F., Gregg, M.A., Boyd, C.S., 2004. Ecology and management of sage-grouse and sage-grouse habitat. *Journal of Range Management* 57, 2–19.
- Cullen, B., Chapman, D., Quigley, P., 2006. Comparative defoliation tolerance of temperate perennial grasses. *Grass and Forage Science* 61, 405–412.
- Damhoureyeh, S.A., Hartnett, D.C., 2002. Variation in grazing tolerance among three tallgrass prairie plant species. *American Journal of Botany* 89, 1634–1643.
- Daubenmire, R.F., 1940. Plant succession due to overgrazing in the *Agropyron* bunchgrass prairie of southeastern Washington. *Ecology* 21, 55–64.
- Delaney, J.T., Moranz, R.A., Debinski, D.M., Engle, D.M., Miller, J.R., 2016. Exotic-dominated grasslands show signs of recovery with cattle grazing and fire. *PLoS One* 11, e0165758.
- Derner, J.D., Briske, D.D., Polley, H.W., 2012. Tiller organization within the tussock grass *Schizachyrium scoparium*: a field assessment of competition–cooperation tradeoffs. *Botany* 90, 669–677.
- Derner, J.D., Lauenroth, W.K., Stapp, P., Augustine, D.J., 2009. Livestock as ecosystem engineers for grassland bird habitat in the western Great Plains of North America. *Rangeland Ecology & Management* 62, 111–118.
- Detling, J., Painter, E., 1983. Defoliation responses of western wheatgrass populations with diverse histories of prairie dog grazing. *Oecologia* 57, 65–71.
- Diamond, J.M., Call, C.A., Devoe, N., 2009. Effects of targeted cattle grazing on fire behavior of cheatgrass-dominated rangeland in the northern Great Basin, USA. *International Journal of Wildland Fire* 18, 944–950.
- Diamond, J.M., Call, C.A., Devoe, N., 2012. Effects of targeted grazing and prescribed burning on community and seed dynamics of a downy brome (*Bromus tectorum*)-dominated landscape. *Invasive Plant Science and Management* 5, 259–269.
- Dietz, T.H., Scott, C.B., Campbell, E.J., Owens, C., Taylor Jr., C.A., Brantley, R., 2010. Feeding redberry juniper (*Juniperus pinchotii*) at weaning increases juniper consumption by goats on pasture. *Rangeland Ecology & Management* 63, 366–372.
- Distel, R.A., Provenza, F.D., 1991. Experience early in life affects voluntary intake of blackbrush by goats. *Journal of Chemical Ecology* 17, 431–450.
- DiTomaso, J.M., 2000. Invasive weeds in rangelands: species, impacts, and management. *Weed Science* 48, 255–265.
- Drewry, J.J., 2006. Natural recovery of soil physical properties from treading damage of pastoral soils in New Zealand and Australia: a review. *Agriculture, Ecosystems & Environment* 114, 159–169.
- Dziba, L.E., Provenza, F.D., Villalba, J.J., Atwood, S.B., 2007. Supplemental energy and protein increase use of sagebrush by sheep. *Small Ruminant Research* 69, 203–207.
- Elzinga, C.L., Salzer, D.W., Willoughby, J.W., 1998. Measuring & monitoring plant populations. Denver, CO, USA, Bureau of Land Management, p. 477.
- Estell, R., Utsumi, S., Cibils, A., Anderson, D., 2014. Is differential use of *Juniperus monosperma* by small ruminants driven by terpenoid concentration? *Journal of Chemical Ecology* 40, 285–293.
- Estell, R.E., 2010. Coping with shrub secondary metabolites by ruminants. *Small Ruminant Research* 94, 1–9.
- Estell, R.E., Havstad, K.M., Cibils, A.F., Fredrickson, E.L., Anderson, D.M., Schrader, T.S., James, D.K., 2012. Increasing shrub use by livestock in a world with less grass. *Rangeland Ecology & Management* 65, 553–562.
- Faeth, S.H., 1992. Interspecific and intraspecific interactions via plant responses to folivory: an experimental field test. *Ecology* 73, 1802–1813.
- Foley, W.J., Iason, G.R., McArthur, C., 1999. Role of plant secondary metabolites in the nutritional ecology of mammalian herbivores: how far have we come in 25 years? In: Jung, H.J.C., Fahey Jr., J.R. (Eds.), *Nutritional Ecology of Herbivores*, Fifth International Symposium on the Nutrition of Herbivores. American Society of Animal Science, Savoy, pp. 130–209.
- Fornoni, J., 2011. Ecological and evolutionary implications of plant tolerance to herbivory. *Functional Ecology* 25, 399–407.

- Frost, R., Walker, J., Madsen, C., Holes, R., Lehfeldt, J., Cunningham, J., Voth, K., Welling, B., Davis, T.Z., Bradford, D., 2012. Targeted grazing: applying the research to the land. *Rangelands* 34, 2–10.
- Frost, R.A., Launchbaugh, K.L., 2003. Prescription grazing for rangeland weed management. *Rangelands* 25, 43–47.
- Frost, R.A., Launchbaugh, K.L., Taylor Jr., C.A., 2008. Age and body condition of goats influence consumption of juniper and monoterpene-treated feed. *Rangeland Ecology & Management* 61, 48–54.
- Frost, R.A., Mosley, J.C., 2012. Sulfur cinquefoil (*Potentilla recta*) response to defoliation on foothill rangeland. *Invasive Plant Science and Management* 5, 408–416.
- Frost, R.A., Mosley, J.C., Roeder, B.L., 2013. Recovery and viability of sulfur cinquefoil seeds from the feces of sheep and goats. *Rangeland Ecology & Management* 66, 51–55.
- Fuhlendorf, S., Engle, D., 2004. Application of the fire–grazing interaction to restore a shifting mosaic on tallgrass prairie. *Journal of Applied Ecology* 41, 604–614.
- Ganskopp, D., Svejcar, T., Taylor, F., Farstvedt, J., 2004. Can spring cattle grazing among young bitterbrush stimulate shrub growth? *Journal of Range Management* 57, 161–168.
- Ganskopp, D., Svejcar, T., Taylor, F., Farstvedt, J., Paintner, K., 1999. Seasonal cattle management in 3 to 5 year old bitterbrush stands. *Journal of Range Management* 52, 166–173.
- Goehring, B.J., Launchbaugh, K.L., Wilson, L.M., 2010. Late-season targeted grazing of yellow starthistle (*Centaurea solstitialis*) with goats in Idaho. *Invasive Plant Science and Management* 3, 148–154.
- Gompert, T., 2010. The power of stock density. *Proceedings of Nebraska Grazing Conference*. University of Nebraska–Lincoln, Kearney, NE, USA, pp. 14–17.
- Goodman, L.E., Cibils, A.F., Lopez, S.C., Steiner, R.L., Graham, J.D., McDaniel, K.C., Abbott, L.B., Stegelmeier, B.L., Hallford, D.M., 2014. Targeted grazing of white locoweed: short-term effects of herbivory regime on vegetation and sheep. *Rangeland Ecology & Management* 67, 680–692.
- Guttery, M.R., 2011. Ecology and management of a high elevation southern range greater sage-grouse population: vegetation manipulation, early chick survival, and hunter motivations [dissertation]. Utah State University, Logan, UT, USA, 119 p.
- Havstad, K., 1994. Sheep grazing as a range improvement tool. *Sheep Research Journal* 10, 72–78.
- Hempy-Mayer, K., Pyke, D.A., 2008. Defoliation effects on *Bromus tectorum* seed production: implications for grazing. *Rangeland Ecology & Management* 61, 116–123.
- Henderson, S.L., Mosley, T.K., Mosley, J.C., Kott, R.W., 2012. Spotted knapweed utilization by sequential cattle and sheep grazing. *Rangeland Ecology & Management* 65, 286–291.
- Hendrickson, J., Moser, L.E., Moore, K., Waller, S.S., 1997. Leaf nutritive value related to tiller development in warm-season grasses. *Journal of Range Management* 50, 116–122.
- Hendrickson, J., Moser, L.E., Moore, K., Waller, S.S., 1998. Morphological development of 2 warm-season grasses in the Nebraska Sandhills. *Journal of Range Management* 51, 456–462.
- Herbel, C.H., 1983. Principles of intensive range improvements. *Journal of Range Management* 36, 140–144.
- Hernández, F., Guthery, F.S., 2012. Beef, brush, and bobwhites: quail management in cattle country. Texas A&M University Press, Kingsville, TX, USA, 288 p.
- Herrick, J.E., Van Zee, J.W., McCord, S.E., Courtright, E.M., Karl, J.W., Burkett, L.M., 2017. Monitoring manual for grassland, shrubland, and savanna ecosystems. In: Volume I: core methods, 2nd ed. USDA-ARS Jornada Experimental Range, Las Cruces, NM, USA. 88003-8003. ISBN 0-9755552-0-0. Available at: <https://jornada.nmsu.edu/monit-assess/manuals/monitoring>. Accessed July 23, 2019.
- Heyworth, C., Iason, G., Temperton, V., Jarvis, P., Duncan, A., 1998. The effect of elevated CO₂ concentration and nutrient supply on carbon-based plant secondary metabolites in *Pinus sylvestris* L. *Oecologia* 115, 344–350.
- Hibbard, W., 2012. Bud Williams' low stress livestock handling. *Stockmanship Journal* 1, 6–163.
- Hofmann, R., 1989. Evolutionary steps of eophysiological adaptation and diversification of ruminants: a comparative view of their digestive system. *Oecologia* 78, 443–457.
- Holling, C.S., 1978. Adaptive environmental assessment and management. John Wiley & Sons, Chichester, UK, 377 p.
- Holopainen, J.K., 2004. Multiple functions of inducible plant volatiles. *Trends in Plant Science* 9, 529–533.
- Holst, P.J., Allan, C.J., 1996. Targeted grazing of thistles using sheep and goats. *Plant Protection Quarterly* 11 (SUP2), 271–273.
- Hulbert, L.C., 1955. Ecological studies of *Bromus tectorum* and other annual bromegrasses. *Ecological Monographs* 25, 181–213.
- Hull, A., Pechanec, J.F., 1947. Cheatgrass—a challenge to range research. *Journal of Forestry* 45, 555–564.
- Hutchings, M., 1999. Clonal plants as cooperative systems: benefits in heterogeneous environments. *Plant Species Biology* 14, 1–10.
- Jensen, C.H., Smith, A.D., Scotter, G.W., 1972. Guidelines for grazing sheep on rangelands used by big game in winter. *Journal of Range Management* 25, 346–352.
- Johnson, J.R., 2012. Stocking density affects trampling and use of vegetation on Nebraska sandhills meadow [thesis]. University of Nebraska, Lincoln, NE, USA, 79 p.
- Karban, R., Agrawal, A.A., Thaler, J.S., Adler, L.S., 1999. Induced plant responses and information content about risk of herbivory. *Trends in Ecology & Evolution* 14, 443–447.
- Kilgour, R.J., 2012. In pursuit of “normal”: a review of the behaviour of cattle at pasture. *Applied Animal Behaviour Science* 138, 1–11.
- Komenda, M., Koppmann, R., 2002. Monoterpene emissions from Scots pine (*Pinus sylvestris*): field studies of emission rate variabilities. *Journal of Geophysical Research: Atmospheres* 107 (D13), 4161.
- Kost, C., Heil, M., 2006. Herbivore-induced plant volatiles induce an indirect defence in neighbouring plants. *Journal of Ecology* 94, 619–628.
- Kott, R., Faller, T., Knight, J., Nudell, D., Roeder, B., 2006. Animal husbandry of sheep and goats for vegetative management. In: Launchbaugh, K., Walker, J. (Eds.), *Targeted grazing: a natural approach to vegetation management and landscape enhancement*. American Sheep Industry Association, Centennial, Colorado, USA, pp. 22–31.
- Kronberg, S.L., Walker, J.W., 2007. Learning through foraging consequences: a mechanism of feeding niche separation in sympatric ruminants. *Rangeland Ecology & Management* 60, 195–198.
- Kuokkanen, K., Yan, S., Niemelä, P., 2003. Effects of elevated CO₂ and temperature on the leaf chemistry of birch *Betula pendula* (Roth) and the feeding behaviour of the weevil *Phyllobius maculicornis*. *Agricultural and Forest Entomology* 5, 209–217.
- Lacey, J.R., Wallander, R., Olson-Rutz, K., 1992. Recovery, germinability, and viability of leafy spurge (*Euphorbia esula*) seeds ingested by sheep and goats. *Weed Technology* 6, 599–602.
- Lagerkvist, C.J., Hess, S., 2010. A meta-analysis of consumer willingness to pay for farm animal welfare. *European Review of Agricultural Economics* 38, 55–78.
- Lane, M.A., Ralphs, M.H., Olsen, J.D., Provenza, F.D., Pfister, J.A., 1990. Conditioned taste aversion: potential for reducing cattle loss to larkspur. *Journal of Range Management* 43, 127–131.
- Launchbaugh, K., Walker, J., Taylor Jr., C., 1999. Foraging behavior: experience or inheritance. In: Launchbaugh, K., Sanders, K., Mosley, J. (Eds.), *Proceedings—grazing behavior of livestock and wildlife symposium*. University of Idaho Forest, Wildlife and Range Experiment Station #70, Moscow, ID, USA, pp. 28–35.
- Launchbaugh, K.L., Walker, J.W., 2006. Targeted grazing—a new paradigm for livestock management. Targeted grazing: a natural approach to vegetation management and landscape enhancement. American Sheep Industry Association, Centennial, CO, USA, pp. 2–8.
- Lehrer Jr., W., Tisdale, E., 1956. Effect of sheep and rabbit digestion on the viability of some range plant seeds. *Journal of Range Management* 9, 118–122.
- Leopold, A., 1933. *Game management*. Charles Scribner's Sons, New York, NY, USA, 481 p.
- Lewis, D.G., Tate, K.W., Harper, J.M., 2000. *Sediment delivery and monitoring: a method of water quality management in rangeland watersheds*, Publication 8014. University of California, Davis, CA, USA, p. 14.
- Lindsey, T., 2016. Grazing method effects on forage production, utilization, animal performance and animal activity on Nebraska sandhills meadow [thesis]. University of Nebraska, Lincoln, NE, USA, 62 p.
- Lusia, J., Peñuelas, J., Alessio, G.A., Estiarte, M., 2006. Seasonal contrasting changes of foliar concentrations of terpenes and other volatile organic compound in four dominant species of a Mediterranean shrubland submitted to a field experimental drought and warming. *Physiologia Plantarum* 127, 632–649.
- Lym, R. G., Sedivec, K. K., and Kirby, D. R. 1997. Leafy spurge control with angora goats and herbicides. *Journal of Range Management* 50:123–128.
- Marquardt, S., Marquez, A., Bouillot, H., Beck, S.G., Mayer, A.C., Kreuzer, M., Alzérrec, H.A., 2009. Intensity of browsing on trees and shrubs under experimental variation of cattle stocking densities in southern Bolivia. *Forest Ecology and Management* 258, 1422–1428.
- Martin, S. C., and Ward, D. E. 1973. Salt and meal-salt help distribute cattle use on semidesert range. *Journal of Range Management* 26:94–97.
- Marty, J.T., 2015. Loss of biodiversity and hydrologic function in seasonal wetlands persists over 10 years of livestock grazing removal. *Restoration Ecology* 23, 548–554.
- Masters, R. A., and Sheley, R. L. 2001. Principles and practices for managing rangeland invasive plants. *Journal of Range Management* 54:502–517.
- Megee, C., 1938. Wild oats or downy brome: troublesome weed on sandy land. Michigan Agricultural Experiment Station Occasional Bulletin 20, 153–156.
- Morris, C., Burton, L., Towers, N., Cullen, N., Rendel, J., Johnson, D., 1998. Genetics of susceptibility to facial eczema in Friesian and Jersey cattle. *New Zealand Journal of Agricultural Research* 41, 347–357.
- Morris, C., Towers, N., Campbell, A., Meyer, H., Wesselink, C., Wheeler, M., 1989. Responses achieved in Romney flocks selected for or against susceptibility to facial eczema, 1975–87. *New Zealand Journal of Agricultural Research* 32, 379–388.
- Morris, C., Towers, N., Wheeler, M., Wesselink, C., 1995. Selection for or against facial eczema susceptibility in Romney sheep, as monitored by serum concentrations of a liver enzyme. *New Zealand Journal of Agricultural Research* 38, 211–219.
- Mosley, J., Frost, R., Barta, S., Standley, E., Schuldt, C., Caquelin, R., Jensen, H., Thompson, F., McCauley, A., 2018. Monitoring for success: official handbook for the Montana Rangeland Monitoring Program. Montana Department of Natural Resources and Conservation, Helena, MT, USA.
- Mosley, J.C., 1985. Let's not forget the art in range management. *Rangelands* 7, 154–155.
- Mosley, J.C., 1994. Prescribed sheep grazing to enhance wildlife habitat on North American rangelands. *Sheep Research Journal* 10, 79–91.
- Mosley, J.C., 1996. Prescribed sheep grazing to suppress cheatgrass: a review. *Sheep and Goat Research Journal* 12, 74–80.
- Mosley, J.C., 1997. Prescribed livestock grazing to sustain and enhance riparian habitat. In: Crane, K., Mosley, J. (Eds.), *Proceedings—ecology and management*

- of grazing by large herbivores, 1997 Montana/Wyoming Range Management Workshop. University of Wyoming Cooperative Extension, Laramie, WY, USA.
- Mosley, J.C., Brewer, T.K., 2006. Targeted livestock grazing for wildlife habitat improvement. In: Launchbaugh, K., Walker, J. (Eds.), Targeted grazing: a natural approach to vegetation management and landscape enhancement. American Sheep Industry Association, Centennial, CO, USA, pp. 115–128.
- Mosley, J.C., Bunting, S.C., Manoukian, M.E., 1999. Cheatgrass. In: Sheley, R.L., Petroff, J.K. (Eds.), Biology and management of noxious rangeland weeds. Oregon State University Press, Corvallis, OR, USA, pp. 175–188.
- Mosley, J.C., Frost, R.A., Roeder, B.L., Kott, R.W., 2017. Targeted sheep grazing to suppress sulfur cinquefoil (*Potentilla recta*) on northwestern Montana rangeland. Rangeland Ecology & Management 70, 560–568.
- Mosley, J.C., Frost, R.A., Roeder, B.L., Mosley, T.K., Marks, G., 2016. Combined herbivory by targeted sheep grazing and biological control insects to suppress spotted knapweed (*Centaurea stoebe*). Invasive Plant Science and Management 9, 22–32.
- Mosley, J.C., Roeder, B.L., Kittle, R., Pauley, J.L., Mangold, J.M., Mosley, T.K., Lucas, D.E., Marks, G., 2017. Targeted cattle grazing to suppress spotted knapweed: effects of diet conditioning. In: Proceedings of the 70th Society for Range Management Annual Meeting, pp. 21–22. Available at: <http://rangelands.org/wp-content/uploads/2018/06/2017-SRM-Annual-Meeting-Abstracts.pdf>. (Accessed 10 December 2018).
- Mosley, J.C., Roselle, L., 2006. Targeted livestock grazing to suppress invasive annual grasses. In: Launchbaugh, K., Walker, J. (Eds.), Targeted grazing: a natural approach to vegetation management and landscape enhancement. American Sheep Industry Association, Centennial, CO, USA, pp. 68–77.
- Nolte, D.L., Provenza, F.D., 1992. Food preferences in lambs after exposure to flavors in milk. Applied Animal Behaviour Science 32, 381–389.
- Nolte, D.L., Provenza, F.D., Callan, R., Panter, K.E., 1992. Garlic in the ovine fetal environment. Physiology & Behavior 52, 1091–1093.
- Olson, B., Launchbaugh, K., 2006. Managing herbaceous broadleaf weeds with targeted grazing. In: Launchbaugh, K., Walker, J. (Eds.), Targeted grazing: a natural approach to vegetation management and landscape enhancement. American Sheep Industry Association, Centennial, CO, USA, pp. 58–67.
- Olson, B.E., Wallander, R.T., 2002. Does ruminal retention time affect leafy spurge seed of varying maturity? Journal of Range Management 55, 65–69.
- Ortega-Reyes, L., Provenza, F.D., 1993. Amount of experience and age affect the development of foraging skills of goats browsing blackbrush (*Coleogyne ramosissima*). Applied Animal Behaviour Science 36, 169–183.
- Ortega-Reyes, L., Provenza, F.D., 1993. Experience with blackbrush affects ingestion of shrub live oak by goats. Journal of Animal Science 71, 380–383.
- Papachristou, T.G., Dziba, L.E., Villalba, J.J., Provenza, F.D., 2007. Patterns of diet mixing by sheep offered foods varying in nutrients and plant secondary compounds. Applied Animal Behaviour Science 108, 68–80.
- Payne, J.M., 2018. Can targeted cattle grazing increase abundance of forbs or arthropods in sage-grouse brood-rearing habitat [thesis]? Montana State University, Bozeman, MT, USA, 59 p.
- Payne, J.M., Mosley, J.C., Litt, A.R., Roeder, B.L., Mosley, T.K., McNew, L.B., Goosey, H.B., 2018. Targeted cattle grazing to enhance sage-grouse brood-rearing habitat. In: Proceedings of the 71st Society for Range Management Annual Meeting, p. 459. Available at: <http://rangelands.org/wp-content/uploads/2018/01/2018-Abstracts.pdf>. (Accessed 20 December 2018).
- Pellant, M., Shaver, P., Pyke, D., Herrick, J., 2005. Interpreting indicators of rangeland health, Version 4. Technical Reference 1734-6. Denver, CO, USA, USDI Bureau of Land Management, p. 136.
- Petersen, C.A., Villalba, J.J., Provenza, F.D., 2014. Influence of experience on browsing sagebrush by cattle and its impacts on plant community structure. Rangeland Ecology & Management 67, 78–87.
- Pittarello, M., Probo, M., Lonati, M., Bailey, D.W., Lombardi, G., 2016. Effects of traditional salt placement and strategically placed mineral mix supplements on cattle distribution in the Western Italian Alps. Grass and Forage Science 71, 529–539.
- Poage, G.W., Scott, C.B., Bisson, M.G., Hartmann, F.S., 2000. Activated charcoal attenuates bitterweed toxicosis in sheep. Journal of Range Management 53, 73–78.
- Pollak, E.R., 2007. Evaluation of low-stress herding and supplement placement to modify cattle distribution and improve pronghorn habitat [thesis]. New Mexico State University, Las Cruces, NM, USA, 123 p.
- Probo, M., Pittarello, M., Lonati, M., Lombardi, G., 2016. Targeted grazing for the restoration of sub-alpine shrub-encroached grasslands. Italian Journal of Agronomy 11, 268–272.
- Provenza, F., Papachristou, T., 2009. Behavior-based management of ecosystems. Options Méditerranéennes, Series A 85, 13–28.
- Provenza, F.D., 1995. Postingestive feedback as an elementary determinant of food preference and intake in ruminants. Journal of Range Management Archives 48, 2–17.
- Provenza, F.D., 1996. Acquired aversions as the basis for varied diets of ruminants foraging on rangelands. Journal of Animal Science 74, 2010–2020.
- Pyke, C.R., Marty, J., 2005. Cattle grazing mediates climate change impacts on ephemeral wetlands. Conservation Biology 19, 1619–1625.
- Ralphs, M.H., Wiedmeier, R.D., 2004. Conditioning cattle to graze broom snakeweed (*Gutierrezia sarothrae*). Journal of Animal Science 82, 3100–3106.
- Rinella, M.J., Jacobs, J.S., Sheley, R.L., Borkowski, J.J., 2001. Spotted knapweed response to season and frequency of mowing. Journal of Range Management 54, 52–56.
- Rogosic, J., Estell, R., Ivankovic, S., Kezic, J., Razov, J., 2008. Potential mechanisms to increase shrub intake and performance of small ruminants in Mediterranean shrubby ecosystems. Small Ruminant Research 74, 1–15.
- Sandercock, B.K., Alfaro-Barrios, M., Casey, A.E., Johnson, T.N., Mong, T.W., Odom, K.J., Strum, K.M., Winder, V.L., 2015. Effects of grazing and prescribed fire on resource selection and nest survival of upland sandpipers in an experimental landscape. Landscape Ecology 30, 325–337.
- Schmelzer, L., Perryman, B., Bruce, B., Schultz, B., McAdoo, K., McCuin, G., Swanson, S., Wilker, J., Conley, K., 2014. Case study: reducing cheatgrass (*Bromus tectorum* L.) fuel loads using fall cattle grazing. The Professional Animal Scientist 30, 270–278.
- Severson, K.E., Urness, P.J., 1994. Livestock grazing: a tool to improve wildlife habitat. In: Vavra, M., Laycock, W.A., Pieper, R.D. (Eds.), Ecological implications of livestock herbivory in the West. Society for Range Management, Denver, CO, USA, pp. 232–249.
- Sharrow, S.H., Seefeldt, S.S., 2006. Monitoring for success. In: Launchbaugh, K., Walker, J. (Eds.), Targeted grazing: a natural approach to vegetation management and landscape enhancement. American Sheep Industry Association, Centennial, CO, USA, pp. 40–49.
- Shaw, R.A., Villalba, J.J., Provenza, F.D., 2006. Influence of stock density and rate and temporal patterns of forage allocation on the diet mixing behavior of sheep grazing sagebrush steppe. Applied Animal Behaviour Science 100, 207–218.
- Sheley, R.L., Jacobs, J.S., Martin, J.M., 2004. Integrating 2, 4-D and sheep grazing to rehabilitate spotted knapweed infestations. Journal of Range Management 57, 371–375.
- Shepard, J.P., Creighton, J., Duzan, H., 2004. Forestry herbicides in the United States: an overview. Wildlife Society Bulletin 32, 1020–1027.
- Shiple, L.A., 1999. Grazers and browsers: how digestive morphology affects diet selection. In: Launchbaugh, K., Sanders, K., Mosley, J. (Eds.), Proceedings—Grazing Behavior of Livestock and Wildlife Symposium. University of Idaho Forest, Wildlife and Range Experiment Station #70, Moscow, ID, USA, pp. 20–27.
- Short, J.J., Knight, J.E., 2003. Fall grazing affects big game forage on rough fescue grasslands. Journal of Range Management 56, 213–217.
- Shropshire, A., 2018. Grazing strategy effects on utilization, animal performance, aboveground production, species composition, and soil properties on Nebraska sandhills meadow [thesis]. University of Nebraska, Lincoln, NE, USA, 118 p.
- Smith, M.A., Malechek, J. C., and Fulgham, K. O. 1979. Forage selection by mule deer on winter range grazed by sheep in spring. Journal of Range Management 32: 40–45.
- Snowder, G., Walker, J., Launchbaugh, K., and Van Vleck, L. D. 2001. Genetic and phenotypic parameters for dietary selection of mountain big sagebrush (*Artemisia tridentata* Nutt. ssp. *vaseyana* [Rydb] Beetle) in Rambouillet sheep. Journal of Animal Science 79:486–492.
- Spinka, M., 2006. How important is natural behaviour in animal farming systems? Applied Animal Behaviour Science 100, 117–128.
- SRM, 1998. Glossary of terms used in range management, fourth edition. Society for Range Management, Denver, CO.
- Stephenson, M.B., Bailey, D.W., Howerly, L.D., Henderson, L., 2016. Efficacy of low-stress herding and low-moisture block to target cattle grazing locations on New Mexico rangelands. Journal of Arid Environments 130, 84–93.
- Stoddart, L.A., Smith, A.D., Box, T.W., 1975. Range management. McGraw-Hill, New York, NY, USA, 532 p.
- Surber, L.M., Rude, M.E., Roeder, B.L., Mosley, T.K., Grove, A.V., Walker, J.W., Kott, R.W., 2011. Percent spotted knapweed (*Centaurea stoebe*) in the diets of grazing sheep. Invasive Plant Science and Management 4, 95–101.
- Tausch, R.J., Nowak, R.S., Bruner, A.D., Smithson, J., 1994. Effects of simulated fall and early spring grazing on cheatgrass and perennial grass in western Nevada. Proceedings of Ecology and Management of Annual Rangelands. USDA Forest Service, Intermountain Research Station. General Technical Report INT-GTR-313. Ogden, UT, USA. Intermountain Research Station, USDA Forest Service, pp. 113–119.
- Taylor Jr., C.A., 2006. Targeted grazing to manage fire risk. In: Launchbaugh, K., Walker, J. (Eds.), Targeted grazing: a natural approach to vegetation management and landscape enhancement. American Sheep Industry Association, Centennial, CO, USA, pp. 107–112.
- Teague, W., Dowhower, S., 2003. Patch dynamics under rotational and continuous grazing management in large, heterogeneous paddocks. Journal of Arid Environments 53, 211–229.
- Thines, N.J., Shipley, L.A., Bassman, J.H., Fellman, J.K., Mattison, D.S., Slusser, J.R., Gao, W., 2007. Effects of enhanced UV-B radiation on plant chemistry: nutritional consequences for a specialist and generalist lagomorph. Journal of Chemical Ecology 33, 1025–1039.
- Thrift, B.D., Mosley, J.C., Brewer, T.K., Roeder, B.L., Olson, B.E., Kott, R.W., 2008. Prescribed sheep grazing to suppress spotted knapweed on foothill rangeland. Rangeland Ecology & Management 61, 18–25.
- Torell, L.A., McDaniel, K.C., Ochoa, C.G., 2005. Economics and optimal frequency of Wyoming big sagebrush control with tebuthiuron. Rangeland Ecology & Management 58, 77–84.
- Tueller, P.T., Tower, J.D., 1979. Vegetation stagnation in three-phase game enclosures. Journal of Range Management 32, 258–263.
- Utsumi, S.A., Cibils, A.F., Estell, R.E., Baker, T.T., Walker, J.W., 2010. One-seed juniper sapling use by goats in relation to stocking density and mixed grazing with sheep. Rangeland Ecology & Management 63, 373–386.
- Utsumi, S.A., Cibils, A.F., Estell, R.E., Soto-Navarro, S.A., Chen, L., Hallford, D.M., 2013. Effects of adding protein, condensed tannins, and polyethylene glycol to diets of

- sheep and goats fed one-seed juniper and low quality roughage. *Small Ruminant Research* 112, 56–68.
- Utsumi, S.A., Cibils, A.F., Estell, R.E., Soto-Navarro, S.A., Van Leeuwen, D., 2009. Seasonal changes in one seed juniper intake by sheep and goats in relation to protein and plant secondary metabolites. *Small Ruminant Research* 81, 152–162.
- Vaarst, M., Alrøe, H.F., 2012. Concepts of animal health and welfare in organic livestock systems. *Journal of Agricultural and Environmental Ethics* 25, 333–347.
- Vallentine, J.F., 2001. *Grazing management*. Academic Press, San Diego, CA, USA, 659 p.
- Varelas, L.A., 2012. Effectiveness and costs of using targeted grazing to alter fire behavior [thesis]. New Mexico State University, Las Cruces, NM, USA, 101 p.
- Verbeke, W., 2009. Stakeholder, citizen and consumer interests in farm animal welfare. *Animal Welfare* 18, 325–333.
- Villalba, J.J., Provenza, F.D., Banner, R.E., 2002. Influence of macronutrients and activated charcoal on intake of sagebrush by sheep and goats. *Journal of Animal Science* 80, 2099–2109.
- Wagner, R., Flynn, J., Gregory, R., 1998. Public perceptions of risk and acceptability of forest vegetation management alternatives in Ontario. *The Forestry Chronicle* 74, 720–727.
- Waldron, D.F., Taylor Jr., C.A., Walker, J.W., Campbell, E.S., Lupton, C.J., Willingham, T.D., Landau, S.Y., 2009. Heritability of juniper consumption in goats. *Journal of Animal Science* 87, 491–495.
- Walker, J.W., 1995. Viewpoint: grazing management and research now and in the next millennium. *Journal of Range Management* 48, 350–357.
- Walker, J.W., Hemenway, K.G., Hatfield, P.G., Glimp, H.A., 1992. Training lambs to be weed eaters: studies with leafy spurge. *Journal of Range Management* 45, 245–249.
- Walker, J.W., Waldron, D.F., Taylor Jr., C.A., Wang, S., Brun, M., Hill, J., Stull, M.A., Mertz, R.P., Schulze, E., Gautam, P., Johnson, C.D., 2019. Breeding goats for juniper (*Juniperus* sp.) consumption using fNIR determinations for phenotypic data. *In: Gold Coast, Australia: 18th International Conference on Near Infrared Spectroscopy*, September 15–20, 2019.
- Wallace, J.M., Wilson, L.M., Launchbaugh, K.L., 2008. The effect of targeted grazing and biological control on yellow starthistle (*Centaurea solstitialis*) in canyon grasslands of Idaho. *Rangeland Ecology & Management* 61, 314–320.
- Wallander, R.T., Olson, B.E., Lacey, J.R., 1995. Spotted knapweed seed viability after passing through sheep and mule deer. *Journal of Range Management* 145–149.
- Walters, C.J., Hilborn, R., 1978. Ecological optimization and adaptive management. *Annual Review of Ecology and Systematics* 9, 157–188.
- Ward, T.A., Tate, K.W., Atwill, E.R., 2003. Guidelines for monitoring the establishment of riparian grazing systems, Publication 8094. University of California, Davis, CA, USA, 37 p.
- Warren, S., Thurow, T., Blackburn, W., Garza, N., 1986. The influence of livestock trampling under intensive rotation grazing on soil hydrologic characteristics. *Journal of Range Management* 39, 491–495.
- Wiedmeier, R.D., Provenza, F.D., Burritt, E.A., 2002. Exposure to ammoniated wheat straw as suckling calves improves performance of mature beef cows wintered on ammoniated wheat straw. *Journal of Animal Science* 80, 2340–2348.
- Willms, W., McLean, A., Tucker, R., Ritcey, R., 1979. Interactions between mule deer and cattle on big sagebrush range in British Columbia. *Journal of Range Management* 32, 299–304.
- Winder, V.L., McNew, L.B., Pitman, J.C., Sandercock, B.K., 2017. Space use of female Greater Prairie-Chickens in response to fire and grazing interactions. *Rangeland Ecology & Management* 70, 165–174.
- Winkel, V.K., Roundy, B.A., 1991. Effects of cattle trampling and mechanical seedbed preparation on grass seedling emergence. *Journal of Range Management* 44, 176–180.
- Winward, A.H., 2000. Monitoring the vegetation resources in riparian areas. *In: Gen. Tech. Rep. RMRS-GTR-47*. USDA Forest Service Rocky Mountain Research Center, Ogden, UT, USA, 49 p.
- Wooley, S.C., Smith, B., King, C., Seastedt, T.R., Knoche, D.G., 2011. The lesser of two weevils: physiological responses of spotted knapweed (*Centaurea stoebe*) to above- and belowground herbivory by *Larinus minutus* and *Cyphocleonus achates*. *Biocontrol Science and Technology* 21, 153–170.
- Yoder, J., 2004. Playing with fire: endogenous risk in resource management. *American Journal of Agricultural Economics* 86, 933–948.
- Yoder, J., Engle, D., and Fuhlendorf, S. 2004. Liability, incentives, and prescribed fire for ecosystem management. *Frontiers in Ecology and the Environment* 2: 361–366.
- Zheng, S., Li, W., Lan, Z., Ren, H., Wang, K., 2015. Functional trait responses to grazing are mediated by soil moisture and plant functional group identity. *Scientific Reports* 5, 18163.