WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ETc) 04/24/15 through 04/30/15

Crops (Leafout Date)	Tehama County - Gerber South			Butte County - Durham			Colusa County - Colusa		
	Past Week	Accum'd	Next Week's	Past Week	Accum'd	Next Week's	Past Week	Accum'd	Next Week's
	of Water	Seasonal	Estimated	of Water	Seasonal	Estimated	of Water	Seasonal	Estimated
	Use	Water Use	ETc	Use	Water Use	ETc	Use	Water Use	ETc
Pasture	1.69	12.31	1.46	1.43	11.20	1.42	1.64	11.91	1.50
Olives Table *	1.28	9.32	1.11	1.08	8.48	1.07	1.23	9.01	1.15
Olives High Density *	1.01	7.36	0.90	0.86	6.69	0.86	0.97	7.11	0.91
Citrus *	1.10	8.03	0.97	0.93	7.33	0.93	1.07	7.76	0.98
Almonds (2/15) *	1.47	8.47	1.33	1.24	7.69	1.29	1.42	8.22	1.37
Prunes (3/6) *	1.58	7.42	1.39	1.34	6.72	1.35	1.53	7.32	1.43
Walnuts (4/1) *	0.95	3.32	0.98	0.81	2.93	0.98	0.92	3.31	1.01
Urban Turf Grass	1.35	9.32	1.18	1.15	8.50	1.14	1.32	9.07	1.22
Past 7 days precipitation (inches)		0.22			0.43			0.23	
Accumulated precipitation (inches)		1.44			2.27			1.34	

Accumulations started on February 10, 2015 or on the approximate leafout date for a specific orchard crop as indicated in parentheses. Criteria for beginning this report are based on the season's last significant rainfall event where the soil moisture profile is estimated to be near its highest level for the new season.

* Estimates are for orchard floor conditions where vegetation is managed by come combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY 1

Crops	Tehama County - Gerber South						Butte	County - D	urham		Colusa County - Colusa					
System Efficiency >>	50%	60%	70%	80%	90%	50%	60%	70%	80%	90%	50%	60%	70%	80%	90%	
Olives Table	2.6	2.1	1.8	1.6	1.4	2.2	1.8	1.5	1.4	1.2	2.5	2.1	1.8	1.5	1.4	
Olives High Density	2.0	1.7	1.4	1.3	1.1	1.7	1.4	1.2	1.1	1.0	1.9	1.6	1.4	1.2	1.1	
Citrus	2.2	1.8	1.6	1.4	1.2	1.9	1.6	1.3	1.2	1.0	2.1	1.8	1.5	1.3	1.2	
Almonds (2/15)	2.9	2.5	2.1	1.8	1.6	2.5	2.1	1.8	1.6	1.4	2.8	2.4	2.0	1.8	1.6	
Prunes (3/6)	3.2	2.6	2.3	2.0	1.8	2.7	2.2	1.9	1.7	1.5	3.1	2.6	2.2	1.9	1.7	
Walnuts (4/1)	1.9	1.6	1.4	1.2	1.1	1.6	1.4	1.2	1.0	0.9	1.8	1.5	1.3	1.2	1.0	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

For further information concerning all counties receiving this report, contact the Tehama Co. Farm Advisor's office at (530) 527-3101 or the Glenn Co. Farm Advisor's office at (530) 865-1153.

^{*} Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.