

Chart 1. Registered Herbicides and Application Timings For Selected Dry Bean Types in California

	Baby Lima	Blackeye	Garbanzo	Kidney	Large Lima
<i>Fallow ground or preformed beds before planting</i>					
carfentrazone (Shark®)	Yes: to 1 day after planting	Yes: to 1 day after planting	Yes: to 1 day after planting	Yes: to 1 day after planting	Yes: to 1 day after planting
glyphosate (Roundup®, etc.)	Yes: to before crop emerge.	Yes: to before crop emerge.	Yes: to before crop emerge.	Yes: to before crop emerge.	Yes: to before crop emerge.
oxyfluorfen (Goal®, etc.)	Yes	Yes	Yes	Yes	Yes
paraquat (Gramoxone Inteon®)	Yes	Yes	Yes	Yes	Yes
<i>Preplant - mechanically incorporated</i>					
eptc (Eptam®)	No	No	No	Yes	No
ethafluralin (Sonalan®)	Yes	Yes	No	Yes	Yes
imazethapyr (Pursuit®)	No	No	Yes	No	No
metribuzin (Sencor®)	No	No	Yes	No	No
pendimethalin (Prowl H2O and EC®)	Yes	Yes	Yes	Yes	Yes
s-metolachlor (Dual Magnum®)	Yes	Yes	Yes	Yes	Yes
trifluralin (Treflan®)	Yes	Yes	Yes	Yes	Yes
<i>After planting - before crop and weed emergence</i>					
flumioxazin (Chateau®)	No	No	Yes	No	No
imazethapyr (Pursuit®)	No	No	Yes	No	No
metribuzin (Sencor®)	No	No	Yes	No	No
oxyfluorfen (Goal®, etc.)	No	No	Yes	No	No
pendimethalin (Prowl H2O and EC®)	No	No	Yes	No	No
<i>After planting - after crop and weed emergence</i>					
bentazon (Basagran®)	Yes	No	No	Yes	Yes
carfentrazone (Shark®)	Yes: shielded in row middles	Yes: shielded in row middles	Yes: shielded in row middles	Yes: shielded in row middles	Yes: shielded in row middles
clethodim (Select Max®)	Yes	Yes	Yes	Yes	Yes
fluazifop (Fusilade DX®)	Yes	No	Yes	Yes	Yes
sethoxydim (Poast®)	Yes	Yes	Yes	Yes	Yes
<i>Lay-by before weed emergence</i>					
eptc (Eptam®)	No	No	No	Yes	No
<i>Pre-harvest aid desiccant</i>					
carfentrazone (Shark®)	Yes	Yes	Yes	Yes	Yes
flumioxazin (Chateau®)	Yes	Yes	Yes	Yes	Yes
<p><i>This is not an endorsement for any of the trade names listed, nor does the omission of specific trade names reflect the view of the author. Please refer to your local dealer or chemical representative for specific herbicide products available but not listed. Always read and follow the label directions carefully before using any pesticide. Refer to the appropriate herbicide label for specific bean types products are registered for use in. Kurt Hembree, UCCE, Fresno County. July 2012. http://cefresno.ucdavis.edu</i></p>					

Chart 2. Selectivity of Annual Broadleaf Weeds to Herbicides Registered in Dry Beans in California

	Preemergent herbicides									Postemergent herbicides								
	eptc (Eptam®)	ethafluralin (Sonalan®)	flumioxazin (Chateau®)	imazethapyr (Pursuit®)	metrinuzin (Sencor®, Metribuzin®)	oxyflurofen (Goal®, etc.)	pendimethalin (Prowl H2O®, EC®)	s-metolachlor (Dual Magnum®)	trifluralin (Treflan®)	bentazon (Basagran®)	carfentrazone (Shark®)	clethodim (Select Max®)	fluzafop (Fusilade DX®)	glyphosate (Roundup®, etc.)	oxyflurofen (Goal®, etc.)	paraquat (Gramoxone Inteon®)	sethoxydim (Poast®)	
Annual Broadleaves																		
Cheeseweed	N	P	C	C	P	C	P	P	N	--	C	N	N	P	C	P	N	
Chickweed	C	C	C	C	C	N	C	C	C	N	C	N	N	C	N	C	N	
Clovers	N	P	--	N	C	P	N	--	N	N	P	N	N	P	P	P	N	
Cocklebur	N	P	N	P	P	C	N	N	N	C	P	N	N	C	C	C	N	
Cudweed	P	P	P	P	N	N	N	N	N	N	P	N	N	C	N	N	N	
Fiddleneck	C	C	C	P	C	C	C	N	C	--	C	N	N	C	C	P	N	
Filaree spp.	P	--	P	P	C	C	N	N	P	--	C	N	N	P	C	P	N	
Goosefoot	C	C	C	P	C	C	C	P	C	N	C	N	N	C	C	C	N	
Groundcherry	C	P	C	C	P	C	N	C	P	C	C	N	N	C	C	C	N	
Groundsel, common	C	C	C	P	P	C	N	N	N	--	C	N	N	C	C	C	N	
Hairy fleabane	C	P	P	N	N	P	N	N	N	N	P	N	N	P	P	P	N	
Henbit	C	P	C	P	--	C	C	--	P	N	C	N	N	C	C	C	N	
Horseweed	C	P	C	N	N	P	N	N	N	N	P	N	N	P	P	P	N	
Knotweed, common	P	P	N	C	N	O	C	N	C	--	P	N	N	P	O	P	N	
Lambsquarters	C	C	C	P	C	C	C	P	C	C	C	N	N	C	C	C	N	
London rocket	C	N	P	C	C	C	P	N	N	P	C	N	N	C	C	C	N	
Morningglory, annual	C	--	C	P	N	P	P	N	P	P	C	N	N	P	P	P	N	
Mustard spp.	N	P	C	C	C	C	P	N	N	C	P	N	N	C	C	C	N	
Nettle	C	P	C	P	C	C	N	--	P	--	C	N	N	N	C	P	N	
Nightshade, black	C	P	C	C	N	C	N	C	N	C	C	N	N	C	C	C	N	
Nightshade, hairy	P	P	C	C	P	C	N	C	N	C	C	N	N	C	C	C	N	
Pigweed spp.	C	C	C	C	C	C	C	C	C	C	C	N	N	C	C	C	N	
Prickly lettuce	C	C	C	N	C	C	N	N	N	--	C	N	N	C	C	P	N	
Puncturevine	N	N	C	C	--	C	P	--	P	N	P	N	N	C	C	C	N	
Purslane, common	C	C	C	C	P	P	C	C	P	C	N	N	N	C	P	C	N	
Redmaids	C	C	C	C	--	C	C	--	C	N	--	N	N	C	C	C	N	
Russian thistle	P	C	C	P	C	P	P	P	P	--	P	N	N	C	P	C	N	
Shepherd's-purse	P	P	C	C	C	P	P	P	N	C	P	N	N	C	P	P	N	
Sowthistle, annual	C	C	P	N	C	C	N	P	N	--	N	N	N	C	C	P	N	
Sunflower, common	N	--	P	P	P	C	N	N	N	C	P	N	N	C	C	P	N	
Velvetleaf	P	N	--	C	P	P	N	P	N	C	C	N	N	P	P	C	N	
Wild radish	N	P	C	P	C	C	N	N	N	P	P	N	N	C	C	C	N	

C = control, P = partial control, N = no control, -- = no information

This is not an endorsement for any of the trade names listed, nor does the omission of specific trade names reflect the view of the author. Please refer to your local dealer or chemical representative for specific herbicide products available but not listed. Always read and follow the label directions carefully before using any pesticide. Ratings reflect appropriate timing and dose according to label recommendations. Refer to the appropriate herbicide label and Chart 1 for specific bean types products listed are registered for use in.

Kurt Hembree, UCCE, Fresno County. July 2012. <http://cefresno.ucdavis.edu>

Chart 3. Selectivity of Annual Grass and Perennial Weeds to Herbicides Registered in Dry Beans in California

	Preemergent herbicides									Postemergent herbicides								
	eptc (Eptam®)	ethafluralin (Sonalan®)	flumioxazin (Chateau®)	imazethapyr (Pursuit®)	metrinuzin (Sencor®, Metribuzin®)	oxyflurofen (Goal®, etc.)	pendimethalin (Prowl H2O®, EC®)	s-metolachlor (Dual Magnum®)	trifluralin (Treflan®)	bentazon (Basagran®)	carfentrazone (Shark®)	clethodim (Select Max®)	fluzafop (Fusilade DX®)	glyphosate (Roundup®, etc.)	oxyflurofen (Goal®, etc.)	paraquat (Gramoxone Inteon®)	sethoxydim (Poast®)	
Annual Grasses																		
Annual bluegrass	C	C	C	N	C	P	C	C	C	N	N	C	N	C	P	P	N	
Barnyardgrass	C	C	C	P	P	P	C	C	C	N	N	C	C	C	P	P	C	
Bromegrass spp.	C	C	C	N	C	N	C	P	C	N	N	P	P	C	N	P	P	
Canarygrass	C	C	N	P	C	P	C	C	C	N	N	C	C	C	P	P	C	
Crabgrass, large	C	C	C	P	C	N	C	C	C	N	N	C	C	C	N	C	C	
Fescue spp.	C	C	N	P	--	N	C	--	C	N	N	P	P	C	N	P	P	
Foxtail spp.	C	C	C	C	P	N	C	C	C	N	N	C	C	C	N	C	C	
Junglerice	C	C	P	P	P	P	C	C	C	N	N	C	C	C	P	P	C	
Lovegrass	C	--	C	N	P	C	C	C	C	N	N	C	C	C	C	P	C	
Ryegrass, Italian	C	N	N	P	C	N	C	C	C	N	N	C	C	C	N	P	C	
Sandbur	C	--	N	P	C	N	C	C	C	N	N	C	C	C	N	P	C	
Sprangletop	C	--	N	N	N	N	C	C	C	N	N	C	C	C	N	N	C	
Wild barley	C	C	N	P	C	P	P	P	C	N	N	C	C	C	P	P	C	
Wild oat	C	C	N	P	P	P	P	N	P	N	N	C	C	C	P	P	C	
Witchgrass	C	--	N	--	--	P	C	C	C	N	N	C	C	C	P	P	C	
Perennials (seed)																		
Bermudagrass	C	--	N	P	N	P	C	N	C	N	N	C	C	C	P	P	C	
Dallisgrass	C	--	N	P	N	N	C	C	C	N	N	C	C	C	N	N	C	
Johnsongrass	C	C	N	P	N	N	C	C	C	N	N	C	C	C	N	C	C	
Field bindweed	N	N	N	--	N	N	P	N	P	P	C	N	N	C	N	P	N	
Perennials (established)																		
Bermudagrass	P	N	N	N	N	N	N	N	N	N	N	P	P	P	N	N	P	
Dallisgrass	N	N	N	N	N	N	N	N	N	N	N	P	P	P	N	N	P	
Johnsongrass	N	N	N	P	N	N	P	N	P	N	N	P	P	P	N	N	P	
Field bindweed	N	N	N	N	N	N	P	N	P	N	P	N	N	P	N	N	N	
Nutsedge, purple	P	P	N	P	N	N	N	N	N	P	N	N	N	P	N	P	N	
Nutsedge, yellow	P	P	N	P	N	N	N	C	N	P	N	N	N	P	N	P	N	

C = control, P = partial control, N = no control, -- = no information

This is not an endorsement for any of the trade names listed, nor does the omission of specific trade names reflect the view of the author. Please refer to your local dealer or chemical representative for specific herbicide products available but not listed. Always read and follow the label directions carefully before using any pesticide. Ratings reflect appropriate timing and dose according to label recommendations. Refer to the appropriate herbicide label and Chart 1 for specific bean types products listed are registered for use in.

Kurt Hembree, UCCE, Fresno County. July 2012. <http://cefresno.ucdavis.edu>

Chart 4. Herbicide and Application Information of Selected Dry Bean Types in California

Preemergent Herbicide	Use Rate/A	Mode-of-Action	PHI	Application Information
<i>Fallow ground or preformed beds before planting</i>				
carfentrazone (Shark®)	1 - 2 fl oz (6.1 fl oz/crop)	Group 14 (WSSA)	0 days	Up to 1 day after planting; ground or air; 5 to 40 gpa; surfactant required; good spray coverage; small annual broadleaf weeds.
glyphosate (Roundup®, etc.)	1 - 4 lb a.i.	Group 9 (WSSA)	0 days	Up to before crop emergence; ground or air; 3 to 30 gpa; AMS added to tank first; weeds <4"; wait 7 days to cultivate perennials; mix with oxyfluorfen for broader control.
oxyfluorfen (Goal®, etc.)	1 - 2 pt (2XL) 0.5 - 1 pt (4F)	Group 14 (WSSA)	0 days	At least 60 days before planting; ground or air; 5 to 40 gpa; mix with glyphosate for broad control; work beds 2.5" deep before planting.
paraquat (Gramoxone Inteon®)	1.5 - 4 pt	Group 22 (WSSA)		By ground or air; 5 to 40 gpa; NIS required; good spray coverage; small annual weeds; Restricted Use permit required.
<i>Preplant - mechanically incorporated</i>				
eptc (Eptam®)	3.5 - 4.5 pt (9 pt/crop)	Group 8 (WSSA)	0 days	Only in kidney beans; mechanically incorporate 2" deep; can be metered through irrigation water immediately after planting.
ethafluralin (Sonalan®)	1.5 - 4.5 pt	Group 3 (WSSA)	0 days	Don't use in garbanzos; 3 to 4.5 pt/acre if nightshade and/or groundcherry are present; at least 5 gpa; mechanically incorporate 2 to 3" deep within 24 hours of treatment.
imazethapyr (Pursuit®)	3 fl oz (1 app./crop)	Group 2 (WSSA)	60 days	Only in garbanzos; up to 7 days before planting; ground or air; mechanically incorporate 1 to 2" deep; can mix with pendimethalin, trifluralin, or s-metolachlor.
metribuzin (Sencor® or Metribuzin 75®)	0.33 - 0.5 lb	Group 5 (WSSA)	0 days	Only in garbanzos; by ground only; 10 to 40 gpa; mechanically incorporate 1 to 2" deep; crop injury can occur under rainy conditions.
pendimethalin (Prowl H2O®, Prowl 3.3 EC®)	1.5 - 3 pt (H2O) 1.8 - 3.6 pt (EC)	Group 3 (WSSA)	0 days	Up to 60 days before planting; mechanically incorporate within 7 days after treatment.
s-metolachlor (Dual Magnum®)	1 - 2 pt (2 pt/crop)	Group 15 (WSSA)	0 days	Mechanically incorporate 2 to 4" deep within 14 days of planting; don't cut for hay within 120 days of treatment; don't use for forage within 60 days of treatment; can cause crop injury of planting deep in cold, wet soil.
trifluralin (Treflan®)	1 - 2 pt	Group 3 (WSSA)	0 days	Mechanically incorporate 2 to 4" deep within 24 hours; can mix with eptc or s-metolachlor.
<i>After planting - before crop and weed emergence</i>				
flumioxazin (Chateau®)	1.5 oz (3 oz/crop)	Group 14 (WSSA)	5 days	Only in garbanzos; within 2 days following planting; activated with ½" of rainfall, sprinkler, or furrow irrigation.
imazethapyr (Pursuit®)	3 fl oz (1 app./crop)	Group 2 (WSSA)	60 days	Only in garbanzos; up to 3 days after planting; ground or air; activate with sprinkler irrigation to wet top 2"; can mix with pendimethalin.
metribuzin (Sencor® or Metribuzin 75®)	0.33 - 0.5 lb	Group 5 (WSSA)	0 days	Only in garbanzos; by ground only; 10 to 40 gpa; shallow mechanical incorporation or with <½" rainfall or sprinkler irrigation; crop injury can occur under rainy, wet conditions.
oxyfluorfen (Goal®, etc.)	1 pt (2XL) 0.5 pt (4F)	Group 14 (WSSA)	0 days	Only in garbanzos; within 2 days after planting; at least 20 gpa; activated with ½" of rainfall, sprinkler, or furrow irrigation.
pendimethalin (Prowl H2O®, Prowl 3.3 EC®)	1.5 - 3 pt (H2O)	Group 3 (WSSA)	0 days	Up to 2 days after planting; activate with sprinkler or furrow irrigation.

Chart 4 (continued). Herbicide and Application Information of Selected Dry Bean Types in California

Preemergent Herbicide	Use Rate/A	Mode-of-Action	PHI	Application Information
<i>After planting - after crop and weed emergence</i>				
bentazon (Basagran®)	1 - 2 pt (4 pt/crop)	Group 6 (WSSA)	30 days	Beans with at least 1 fully expanded trifoliolate leaf; COC, UAN, or AMS can improve weed control, but may also increase crop injury; ground or air; 5 to 20 gpa; thorough spray coverage; weeds not stressed for moisture; does not control grasses; don't cultivate 5 days before or 7 days after treatment split treatment 10 to 14 days apart for nutsedge control.
carfentrazone (Shark®)	0.5 - 2 fl oz (6.1 fl oz/crop)	Group 14 (WSSA)	0 days	Hooded or shielded spray to row middles; at least 10 gpa; surfactant needed; keep spray off crop foliage; travel speed <5 mph; broadleaf weeds small; good spray coverage is essential.
clethodim (Select Max®)	9 - 32 fl oz (64 fl oz/crop)	Group 1 (WSSA)	30 days	Controls grasses only; AMS mixed in tank first; NIS needed; ground or air; 3 to 40 gpa; within 7 days after irrigating; repeat treatment 14 days apart; cultivate 7 to 14 days later.
fluazifop (Fusilade DX®)	8 - 24 fl oz (48 fl oz/crop)	Group 1 (WSSA)	60 days	Not for blackeyes, controls grasses only; COC or NIS needed; 5 to 40 gpa; within 7 days after irrigating; repeat treatment at 14 days apart; cultivate 7 to 14 days later.
sethoxydim (Poast®)	1.5 - 2.5 pt (4 pt/crop)	Group 1 (WSSA)	30 days	Controls grasses only; COC or MSO needed; ground or air; at least 10 gpa; within 7 days after an irrigating; repeat treatment 14 days apart; can mix with bentazon in kidney and lima beans; cultivate 7 to 14 days later.
<i>Lay-by before weed emergence</i>				
eptc (Eptam®)	2.25 - 4.5 pt (9 pt/crop)	Group 8 (WSSA)	0 days	Use only in kidney beans; directed application before pods form; keep off crop foliage; activate with irrigation water.
<i>Pre-harvest aid desiccant</i>				
carfentrazone (Shark®)	1 to 2 fl oz (6.1 fl oz/crop)	Group 14 (WSSA)	0 days	Ground or air; NIS, COC, or MSO needed; 15 to 30 gpa with good coverage.
flumioxazin (Chateau®)	1 - 3 oz (3 oz/crop)	Group 14 (WSSA)	5 days	Ground or air; COC or MSO needed; 15 to 30 gpa with good coverage.
<p><i>This is not an endorsement for any of the trade names listed, nor does the omission of specific trade names reflect the view of the author. Refer to your local chemical dealer or manufacturer for specific herbicide products available. Numerous factors influence the performance of herbicides, so consult the manufacturer's labels for specific treatment recommendations. Refer to Chart 1 and the appropriate herbicide label for specific bean types products are registered for use in. When combining products always employ the most restrictive limits allowed by the labels.</i></p> <p><i>Kurt Hembree, UCCE, Fresno County. July 2012. http://cefresno.ucdavis.edu</i></p>				