

# Nutrition and Health Info-Sheet

For Health Professionals

Produced by  
Karrie Heneman, PhD  
Sheri Zidenberg-Cherr, PhD  
UC Cooperative Extension  
Center for Health and  
Nutrition Research  
Department of Nutrition  
University of California  
Davis, CA 95616-8669  
December 2007

## *Some Facts About Flavonols*

### *What are flavonols?*

Flavonols are phytochemical compounds found in high concentrations in a variety of plant-based foods and beverages. Based on their structure, flavonols are classified as flavonoids and include the following compounds: quercetin, kaempferol, and myricetin. The specific amounts of flavonols in foods are affected by a range of factors including plant type and growth, season, light, degree of ripeness, food preparation, and processing. High concentrations of flavonols can be found in apples, apricots, beans, broad beans, broccoli, cherry tomatoes, chives, cranberries, kale, leeks, pear, onions, red grapes, sweet cherries, and white currants (1). Below is a list of the flavonol content of various foods:



*The Flavonol Content of Various Foods (2)*

<i>Food</i>	<i>Quercetin Content (mg/100g)</i>	<i>Kaempferol (mg/100g)</i>	<i>Myricetin (mg/100g)</i>
Apples	4.3	0.02	0.0
Apricots	2.1	0.0	0.0
Beans	0.0	31.3	0.0
Broad Beans (Fava Beans)	0.6	0.4	0.0
Broccoli	2.5	0.01	4.0
Cherry Tomatoes	2.8	0.1	0.0
Chives	4.8	10.0	0.0
Cranberries	15.1	0.1	6.8
Kale	7.7	26.7	0.0
Leeks	0.1	3.0	0.0
Pear	4.5	0.0	0.0
Onions	21.4	0.6	0.02
Red Grapes	1.4	0.0	0.01
Sweet Cherries	2.6	0.0	0.0
Tea	2.7	0.9	0.9
White Currant	2.7	0.2	0.2

### *Are there beneficial health effects associated with consumption of flavonols?*

Consumption of flavonols has been associated with a variety of beneficial effects including an increase in erythrocyte superoxide dismutase activity, decrease in lymphocyte DNA damage, decrease in urinary 8-hydroxy-2'-deoxyguanosine, and an increase in plasma antioxidant capacity (3).

### *There seems to be a lot of media hype around apples, broccoli, and cranberries. Are these really “super foods”?*

As shown in the above table, apples, broccoli, and cranberries are all high in flavonols, in addition to a variety of other polyphenolic compounds. Below is a summary of the research that has looked at the effects of consuming these foods:

- **Apples:** Research has investigated the relationship between consumption of apples and a variety of chronic diseases including cancer, cardiovascular disease, asthma, and diabetes (4).



- **Cancer:** A reduced risk of oral, pharynx, esophagus, colon, larynx, breast, ovary and prostate cancer has been observed in those consuming greater than or equal to one apple a day in comparison to those consuming less (5). In support of these findings, previous research in *in vitro* systems has found that apple flavonoids can positively affect carcinogen bioactivation, cell-signaling, cell cycle regulation, angiogenesis, oxidative stress, and inflammation (5).
- **Cardiovascular Disease:** Consumption of apples has been associated with a reduced risk of cardiovascular disease, cardiovascular events, coronary mortality, and thrombotic stroke (4).
- **Asthma:** In adults, a reduced risk of asthma, bronchial sensitivity, and chronic obstructive pulmonary disease in addition to improved lung function (as measured by forced expiratory volume) has been associated with apple consumption (4).
- **Diabetes:** Apple consumption has also been associated with a reduced risk of type 2 diabetes. In comparison to women who did not consume apples, consumption of at least one apple per day was associated with a 28% reduced risk of this chronic disease (6).
- **Conclusions:** Current research supports an inverse relationship between consumption of apples and a variety of chronic diseases. In light of these findings, consuming an apple a day as part of the recommended intake of fruits and vegetables may prove to be beneficial to overall health; however, variety is important and apples should not be the sole fruit consumed by an individual.

- **Broccoli:** Cruciferous vegetables are part of the plant family Brassicaceae, which includes broccoli, in addition to cabbage, cauliflower, Brussels sprouts, turnips, and watercress. Consumption of this group of plant foods has been associated with a reduction in risk of several cancers including lung, breast, colorectal, and prostate (7).



- **Cancer:** Epidemiological evidence supports an inverse association between consumption of broccoli and breast cancer risk in pre-menopausal women (8), prostate cancer (9), bladder cancer in men (10), lung cancer in postmenopausal women, colon cancer in postmenopausal women, cancer of the respiratory tract, stomach cancer in men, cancer of the reproductive organs in women, and thyroid cancer (11).
- **Conclusions:** In light of this research, the American Cancer Society recommends consuming broccoli as part of a balanced diet that includes foods from a variety of plant sources.

- **Cranberries:** Cranberries are commonly touted as a folk remedy for treating urinary tract infections. Current research has also investigated the relationship between consumption of cranberry products and cancer and cardiovascular disease.
  - *Urinary Tract Infections:* Current scientific evidence suggests that use of cranberry products can prevent new urinary tract infections, possibly through inhibition of bacterial adhesion to the urinary tract (12). However, caution is warranted as cranberry juice may interact with warfarin, resulting in bleeding (13).
  - *Cancer:* Despite promising findings in experimental models suggesting that cranberries can protect against breast, colon, prostate, lung, and other tumors (14), current human research does not provide evidence that consumption of cranberry products reduces risk of cancer (15).
  - *Heart Disease:* There is limited clinical evidence showing that consumption of cranberry products can protect against cardiovascular disease possibly by decreasing LDL oxidation and increasing plasma antioxidant capacity (16).
  - *Conclusions:* The Dietary Guidelines for Americans, 2005 recommends limiting consumption of juice to ensure adequate intake of fiber (31g/ 2000 kcals). For women prone to urinary tract infections, consuming ½ cup of cranberry juice daily may help to reduce the number of new infections. As with apples and broccoli, current scientific evidence supports consumption of cranberry products as part of a varied diet.



## REFERENCES

1. Aheme SA, and O'Brien NM. Dietary flavonols: chemistry, food content, and metabolism. *Nutrition*; 2002; 18: 75-81.
2. Nutrient Data Laboratory US Department of Agriculture. USDA Database for the Flavonoid Content of Selected Foods. US Department of Agriculture, Agricultural Research Service, 2007.
3. Williamson G, and Manach C. Bioavailability and bioefficacy of polyphenols in humans. II. Review of 93 intervention studies. *Am J Clin Nutr*; 2005; 81: 243S-255S.
4. Boyer J, Liu RH. Apple phytochemicals and their health benefits. *Nutr J*; 2004; 3: 5.
5. Gallus S, et al. Does an apple a day keep the oncologist away? *Ann Oncol*; 2005;16:1841-4.
6. Song Y, et al. Associations of dietary flavonoids with risk of type 2 diabetes, and markers of insulin resistance and systemic inflammation in women: a prospective study and cross-sectional analysis. *J Am Coll Nutr*; 2005; 24: 376-84.
7. Juge N, et al. Molecular basis for chemoprevention by sulforaphane: a comprehensive review. *Cell Mol Life Sci*; 2007; 64: 1105-27.
8. Ambrosone CB, et al. Breast cancer risk in premenopausal women is inversely associated with consumption of broccoli, a source of isothiocyanates, but is not modified by GST genotype. *J Nutr*; 2004; 134: 1134-8.

9. Kirsh VA, et al. Prospective study of fruit and vegetable intake and risk of prostate cancer. *J Natl Cancer Inst*; 2007; 99: 1200-9.
10. Michaud DS, et al. Fruit and vegetable intake and incidence of bladder cancer in a male prospective cohort. *J Natl Cancer Inst*; 1999; 91: 605-13.
11. Verhoeven DT, et al. Epidemiological studies on brassica vegetables and cancer risk. *Cancer Epidemiol Biomarkers Prev*; 1996; 5: 733-48.
12. Jepson RG, and Craig JC. A systematic review of the evidence for cranberries and blueberries in UTI prevention. *Mol Nutr Food Res*; 2007; 51: 738-45.
13. Pham DQ, and Pham AQ. Interaction potential between cranberry juice and warfarin. *Am J Health Syst Pharm*; 2007; 64: 490-4.
14. Neto CC. Cranberry and its phytochemicals: a review of in vitro anticancer studies. *J Nutr*; 2007; 137: 186S-193S.
15. Duthie SJ, et al. The effects of cranberry juice consumption on antioxidant status and biomarkers relating to heart disease and cancer in healthy human volunteers. *Eur J Nutr*; 2006; 45: 113-22.
16. Ruel G, et al. Changes in plasma antioxidant capacity and oxidized low-density lipoprotein levels in men after short-term cranberry juice consumption. *Metabolism*; 2005; 54: 856-61.

*\* Production of this material was supported by a grant from the Vitamin Cases Consumer Settlement Fund, created as a result of an antitrust class action. One of the purposes of the fund is to improve the health and nutrition of California consumers.*

---

The University of California, in accordance with Federal and State law and University policy, does not discriminate on the basis of race, color, national origin, religion, sex, disability, age, medical condition (cancer-related), ancestry, marital status, citizenship, sexual orientation, or status as a Vietnam-era veteran or special disabled veteran. The University also prohibits sexual harassment. Inquiries regarding the university's nondiscrimination policies may be directed to the Affirmative Action Director, University of California, Agriculture and Natural Resources, 300 Lakeside Drive, 6th floor, Oakland, CA 94612-3560, (510) 987-0096.

---