

WEEKLY SOIL MOISTURE LOSS IN INCHES (Estimated Crop Evapotranspiration or ET _c) 09/13/24 through 09/19/24												
Crops (Leafout Date)	#148 Merced				#39 Parlier				#258 Lemon Cove			
	09/13 - 09/19 Water Use	Accum'd Seasonal Water Use	09/20 - 09/26 Estimated ET _c		09/13 - 09/19 Water Use	Accum'd Seasonal Water Use	09/20 - 09/26 Estimated ET _c		09/13 - 09/19 Water Use	Accum'd Seasonal Water Use	09/20 - 09/26 Estimated ET _c	
Almonds (3/1) *	1.32	42.11	1.16		1.29	43.85	1.12		1.26	42.52	1.02	
Pistachio (4/20) * **	1.26	34.68	1.09		1.23	36.13	1.05		1.20	35.07	0.95	
Citrus (2/1)	0.87	30.89	0.77		0.84	32.31	0.76		0.81	31.30	0.67	
Raisin Grapes (3/11) (11 ft. row spacing)	0.90	24.58	0.81		0.89	25.61	0.78		0.87	24.87	0.75	
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis) ***	1.02	27.09	0.93		0.99	28.28	0.89		0.97	27.44	0.81	
Walnuts (4/20)	1.31	32.74	1.12		1.32	34.22	1.11		1.29	33.21	1.02	
Stone Fruit (3/11)	1.41	34.78	1.30		1.37	36.38	1.26		1.35	35.28	1.13	
Past 7 days precipitation (inches)	0.00				0.07				0.02			
Accumulated precipitation (inches) (1/1/2024)	14.98				9.05				9.73			
Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop * Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth. ** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.												
PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹												
Crops	#148 Merced				#39 Parlier				#258 Lemon Cove			
	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
System Efficiency >>												
Almonds (3/1)	2.0	1.8	1.6	1.4	2.0	1.7	1.5	1.4	1.9	1.7	1.5	1.3
Pistachio (4/20)	1.9	1.7	1.5	1.3	1.9	1.6	1.4	1.3	1.8	1.6	1.4	1.3
Citrus (2/1)	1.3	1.2	1.0	0.9	1.3	1.1	1.0	0.9	1.2	1.1	1.0	0.9
Raisin Grapes (3/11) (11 ft. row spacing)***	Assume all grape			0.9	Assume all grape			0.9	Assume all grape			0.9
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis) ***	irrigation type is drip			1.1	irrigation type is drip			1.0	irrigation type is drip			1.0
Walnuts (4/20)	2.0	1.7	1.5	1.4	2.0	1.8	1.6	1.4	2.0	1.7	1.5	1.4
Stone Fruit (3/11)	2.2	1.9	1.7	1.5	2.1	1.8	1.6	1.4	2.1	1.8	1.6	1.4
¹ The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.												
PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE												
Crops	#148 Merced				#39 Parlier				#258 Lemon Cove			
Almonds 115 Trees/A	472	425	378	331	472	401	354	331	449	401	354	307
Pistachio 106 Trees/A	473	424	374	324	473	399	349	324	448	399	349	324
Citrus 110 Trees/A	321	296	247	222	321	272	247	222	296	272	247	222
Raisin Grapes 566 Vines/A	Assume all grape			43	Assume all grape			43	Assume all grape			43
Winegrapes 622 Vines/A	irrigation type is drip			48	irrigation type is drip			44	irrigation type is drip			44
Walnuts 76 Trees/A	715	607	536	500	715	643	572	500	715	607	536	500
Stonefruit 172 Trees/A	347	300	268	237	332	284	253	221	332	284	253	221
For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.												



WEEKLY SOIL MOISTURE LOSS IN INCHES
(Estimated Crop Evapotranspiration or ET_c)
09/13/24 through 09/19/24

Crops (Leafout Date)	#124 Panoche				#2 Five Points				#15 Stratford		
	09/13- 09/19 Water Use	Accum'd Seasonal Water Use	09/13- 09/19 Estimated ETc		09/13- 09/19 Water Use	Accum'd Seasonal Water Use	09/13- 09/19 Estimated ETc		09/13- 09/19 Water Use	Accum'd Seasonal Water Use	09/13- 09/19 Estimated ETc
Almonds (3/1) *	1.27	45.44	1.24		1.25	46.08	1.35		1.36	44.10	1.31
Pistachio (4/20) * **	1.21	38.01	1.17		1.17	38.39	1.28		1.28	37.02	1.24
Citrus (2/1)	0.81	33.66	0.84		0.78	34.31	0.91		0.84	32.74	0.89
Raisin Grapes (3/11) (11 ft. row spacing)	0.87	26.82	0.88		0.85	27.16	0.97		0.91	26.03	0.94
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	0.99	29.60	0.99		0.94	29.84	1.09		1.04	28.74	1.05
Walnuts (4/20)	1.29	35.94	1.17		1.25	36.23	1.28		1.37	35.20	1.24
Stone Fruit (3/11)	1.34	38.34	1.38		1.32	38.65	1.49		1.43	37.36	1.45
Past 7 days precipitation (inches)	0.00				0.08				0.02		
Accumulated precipitation (inches) (1/1/2024)	6.60				6.94				5.45		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop
 * Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.
 ** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹

Crops	#124 Panoche					#2 Five Points					#15 Stratford			
	65%	75%	85%	95%		65%	75%	85%	95%		65%	75%	85%	95%
System Efficiency >>	65%	75%	85%	95%		65%	75%	85%	95%		65%	75%	85%	95%
Almonds (3/1)	2.0	1.7	1.5	1.3		1.9	1.7	1.5	1.3		2.1	1.8	1.6	1.4
Pistachio (4/20)	1.9	1.6	1.4	1.3		1.8	1.6	1.4	1.2		2.0	1.7	1.5	1.3
Citrus (2/1)	1.2	1.1	1.0	0.9		1.2	1.0	0.9	0.8		1.3	1.1	1.0	0.9
Raisin Grapes (3/11) (11 ft. row spacing)	Assume all grape			0.9		Assume all grape			0.9		Assume all grape			1.0
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	irrigation type is drip			1.0		irrigation type is drip			1.0		irrigation type is drip			1.1
Walnuts (4/20)	2.0	1.7	1.5	1.4		1.9	1.7	1.5	1.3		2.1	1.8	1.6	1.4
Stone Fruit (3/11)	2.1	1.8	1.6	1.4		2.0	1.8	1.6	1.4		2.2	1.9	1.7	1.5

¹ The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

Crops	#124 Panoche					#2 Five Points					#15 Stratford			
	472	401	354	307		449	401	354	307		496	425	378	331
Almonds 115 Trees/A	472	401	354	307		449	401	354	307		496	425	378	331
Pistachio 106 Trees/A	473	399	349	324		448	399	349	299		498	424	374	324
Citrus 110 Trees/A	296	272	247	222		296	247	222	197		321	272	247	222
Raisin Grapes 566 Vines/A	Assume all grape			43		Assume all grape			43		Assume all grape			48
Winegrapes 622 Vines/A	irrigation type is drip			44		irrigation type is drip			44		irrigation type is drip			48
Walnuts 76 Trees/A	715	607	536	500		679	607	536	464		750	643	572	500
Stonefruit 172 Trees/A	332	284	253	221		316	284	253	221		347	300	268	237

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.