

# The Influence of SOD on Fire Behavior



Steven Swain

Environmental Horticulture Advisor

UCCE Marin & Sonoma Counties

# SOD & Fire

- Natural fires
  - Big Sur
    - Slightly different ecosystems
    - Chalk
    - Big Basin
    - Pfeiffer
- Eradication efforts
  - Clearing and burning
    - Oregon



Photo: Rani Sanderson

# Reports of SOD-Fire interactions

**“The oak trees, many of which had been killed by sudden oak death, helped accelerate the spread of the fire...”**


New York Times, July 8, 2008, in “Gains Reported on 2 California Blazes, but Worries Persist” by Felicity Barringer

**“...Hundreds of thousands of oak trees in the area have been killed in recent years by a disease known as sudden oak death, producing fuel that allows flames to spread more quickly through redwoods and other evergreens, [forest experts] said.”**

Los Angeles Times, July 7, 2008, in “Fungus-killed oaks make Basin Complex fire hotter, harder to fight” by Deborah Schoch




# Assumed hypothesis





How to evaluate this hypothesis?

# Did disease increase fire severity?

Metz et al. 2011.  
*Ecological Applications*



# Did disease increase fire severity?



# What is the disconnect?

- “Forest experts” just seeing what they expect?
  - Maybe ...
- Or is SOD & Fire more complex than the regression (or press) accounts for?



# Fuels vary with disease stage

Early...



...Late



Surface, ladder and aerial fuels  
Various stages of fragmentation and decay





# Early stage disease: More crown fires, scorching, torching



15 plots with *P. ramorum* have no recent mortality

# Late stage disease: More (dead) logs, greater soil burn severity



17 plots with *P. ramorum*  
have *no* logs

# Fire & SOD in redwood forests



Tanoak

*Notholithocarpus densiflorus*



Coast redwood  
*Sequoia sempervirens*




California bay laurel  
*Umbellularia californica*

# Species differ in susceptibility

	Fire	SOD
Tanoak	Sensitive	High mortality
Bay Laurel	Sensitive	Negligible impacts
Redwood	Resilient	Negligible impacts

Are *joint* impacts of fire and SOD additive or synergistic?

# Fire & SOD in redwood forests



# Fuels vary with disease stage


Early...

...Late



Surface, ladder and aerial fuels  
Various stages of fragmentation and decay


# Disease stage affects redwood risk



- Mid stage SOD = higher damage to redwoods
- Late stage SOD = lower damage to redwoods

# Disease stage affects redwood risk


Mortality of 35 cm trees  
Without SOD = 20%  
With SOD = 75%




5% Mortality Risk  
Without SOD,  $\geq 48$  cm  
With SOD,  $\geq 72$ cm



# Dead tanoaks carried flames upwards





- No SOD, early-stage SOD
- Middle stage SOD
- Late stage SOD

# Does SOD increase fire risk?

- The Goldilocks story
  - If it does, conditions have to be *just right*
  - Otherwise, the answer is “no” ...
    - ... and even if it is just right, the number of plots this occurs on is relatively few.
- Who’s risk?
  - Pfeiffer fire Dec 2013
  - Fire department saved all homes
  - 30 burned when it doubled and tripled back
    - Long “residence times”
  - No lives lost

# What about the potential for positive effects?

- California has been managed with fire for thousands of years
  - California forests are adapted to periodic low intensity fires
- Can fire be used to eliminate *P. ramorum* from forest stands?



# Eradication Effort


- Decade long study
  - Now abandoned
- Total host removal
- Burning of slash and landscape



Success?  
Could natural fires succeed?

# Pathogen recovery post-fire

- *P. ramorum* found in previously positive sites
  - 20% in 2009
  - 40% in 2010



Beh, Metz et al. 2012.  
*New Phytologist*.










# Pathogen recovery post-fire

- *P. ramorum* found in previously positive sites
  - 20% in 2009
  - 40% in 2010
- Surviving key hosts as pathogen reservoir in patchily burned landscape



# Conclusions

- On the whole, SOD doesn't significantly alter fire behavior
  - The devil's in the details
- Mid stage damage from SOD increases fire severity in a brief window of time
  - Within it, they interact in surprising ways
    - Increased redwood mortality
- SOD isn't eradicated by fire





# Management Recommendations

- Reduce the amount of *standing* fuels
  - Homes
    - CalFire clearances
  - Redwoods
- Lop downed fuels to below knee height
- Risk to who?
  - Increased fuels may have real consequence for homeowners
  - Fire ecologists and fire fighters: two disciplines divided by a common language

# Resources

- [www.suddenoakdeath.org](http://www.suddenoakdeath.org)
- Metz, M. et al. (2013) Unexpected Redwood Mortality from Synergies Between Wildfire and Emerging Infectious Disease. *Ecology* 94:2152–2159 <http://dx.doi.org/10.1890/13-0915.1>
- Metz, M. et al. (2011) Interacting Disturbances: Wildfire Severity Affected by Stage of Forest Disease Invasion. *Ecological Applications* 21(2):313-320 <http://www.esajournals.org/doi/pdf/10.1890/10-0419.1>
- Beh, M. et al. (2012) The Key Host for an Invasive Forest Pathogen Also Facilitates the Pathogen's Survival of Wildfire in California Forests <http://onlinelibrary.wiley.com/doi/10.1111/j.1469-8137.2012.04352.x/abstract>
- CalFire clearances: [http://www.calfire.ca.gov/communications/downloads/fact\\_sheets/DefensibleSpaceFlyer.pdf](http://www.calfire.ca.gov/communications/downloads/fact_sheets/DefensibleSpaceFlyer.pdf)
- This presentation is on line at: <http://ucanr.edu/MarinIPM>
- Steven Swain: [svswain@ucanr.edu](mailto:svswain@ucanr.edu)  
415 473 4226

