WEEKLY ET REPORT

(Estimated Crop Evapotranspiration or ETc) 05/18/18 through 05/24/18

Crops (Leafout Date)	Tehama County - Gerber South			Butt	e County -	Biggs	Butte County - Durham				Colusa County - William		
	Past Week	Accum'd	Next Week's	Past Week	Accum'd	Next Week's	Past Week	Accum'd	Next Week's	Ī	Past Week	Accum'd	Next Weel
	of Water	Seasonal	Estimated	of Water	Seasonal	Estimated	of Water	Seasonal	Estimated		of Water	Seasonal	Estimated
	Use	Water Use	ETc	Use	Water Use	ETc	Use	Water Use	ETc		Use	Water Use	ETc
Pasture [ETo]	1.73	15.49	1.69	1.60	14.95	1.55	1.49	13.94	1.55		1.45	15.09	1.62
Olives Table *	1.31	11.71	1.27	1.21	11.32	1.19	1.14	10.60	1.19		1.11	11.46	1.20
Olives High Density *	1.04	9.35	1.00	0.95	9.00	0.92	0.90	8.39	0.92		0.86	9.07	0.98
Citrus *	1.12	10.11	1.11	1.04	9.66	0.99	0.97	9.08	0.99		0.93	9.80	1.06
Almonds (2/16) *	1.71	12.36	1.69	1.58	11.92	1.55	1.47	11.19	1.55		1.43	11.93	1.62
Cling Peaches (3/1) *	1.36	7.28	1.34	1.25	7.00	1.20	1.15	6.60	1.20		1.12	6.86	1.27
Pistachios (4/16) *	1.59	5.67	1.55	1.46	5.36	1.41	1.39	5.09	1.41		1.34	5.17	1.48
Prunes (3/20) *	1.66	10.96	1.62	1.53	10.51	1.48	1.42	9.88	1.48		1.38	10.40	1.55
Walnuts (4/1) *	1.80	9.28	1.76	1.67	8.89	1.62	1.56	8.33	1.62		1.52	8.61	1.69
Urban Turf Grass	1.39	12.04	1.38	1.28	11.67	1.29	1.21	10.92	1.29		1.18	11.79	1.31
Past 7 days precipitation (inches)		(0.00)			(0.00)			(0.00)				(0.00)	•
Accumulated precipitation (inches)		(5.26)			(5.52)			(6.58)				(3.65)	

Accumulations started on February 16, 2018 or on the approximate leafout date for a specific orchard crop as indicated in parentheses. Criteria for beginning this report are based on the season's last significant rainfall event where the soil moisture profile is estimated to be near its highest level for the new season.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY 1

Crops	Tehama County - Gerber South			Butt	Butte County - Biggs			Butte County - Durham			Colusa County - Williams		
System Efficiency >>	70%	80%	90%	70%	80%	90%	Ī	70%	80%	90%	70%	80%	90%
Olives Table	1.9	1.6	1.5	1.7	1.5	1.3	Ī	1.6	1.4	1.3	1.6	1.4	1.2
Olives High Density	1.5	1.3	1.2	1.4	1.2	1.1		1.3	1.1	1.0	1.2	1.1	1.0
Citrus	1.6	1.4	1.2	1.5	1.3	1.2		1.4	1.2	1.1	1.3	1.2	1.0
Almonds (2/16)	2.4	2.1	1.9	2.3	2.0	1.8		2.1	1.8	1.6	2.0	1.8	1.6
Cling Peaches (3/1)	1.9	1.7	1.5	1.8	1.6	1.4		1.6	1.4	1.3	1.6	1.4	1.2
Pistachios (4/16)	2.3	2.0	1.8	2.1	1.8	1.6		2.0	1.7	1.5	1.9	1.7	1.5
Prunes (3/20)	2.4	2.1	1.8	2.2	1.9	1.7		2.0	1.8	1.6	2.0	1.7	1.5
Walnuts (4/1)	2.6	2.3	2.0	2.4	2.1	1.9		2.2	2.0	1.7	2.2	1.9	1.7

¹ The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

For further information concerning all counties receiving this report, contact the Tehama Co. Farm Advisor's office at (530) 527-3101 or the Glenn Co. Farm Advisor's office at (530) 865-1153.

This same information and source is now available in the ET Reports section of the sacvalleyorchard.com website. Same information, just in a different format.

^{*} Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.