

NITROGEN DYNAMICS IN AN ORGANIC STRAWBERRY PRODUCTION SYSTEM

Joji Muramoto¹, Stephen R. Gliessman¹, Dan Schmida², Robert Stephens³, Carol Shennan¹, and Sean T. Swezey^{1,4}
¹UC Santa Cruz, ²Sandpiper Farms, ³Elkhorn Ranch, ⁴UC-SAREP

ABSTRACT

Optimizing nitrogen (N) management in organic strawberries on the central coastal California is a challenge because of high N sensitivity and the long growth period of strawberries, transplanting immediately followed by the winter rainy season and unpredictability of N mineralization from organic fertilizers and soil organic matter. Moreover, research-based information on N fertility management in organic strawberries is lacking. In a replicated on-farm organic rotation trial in Moss Landing, California, N dynamics in an organic strawberry production system was monitored for two years. In the first year, we examined the grower's current N-management practice, and an improved N-management practice was demonstrated in the second year. Fertility management practices, soil inorganic N content (0-60cm deep), and fruit yield, plant biomass, and N content of strawberries (cv. Aromas) were monitored. Changes in cumulative daily N uptake of strawberries were calculated and N-loss from the root zone (0-30cm deep in the bed area) during the winter rainy season was estimated. We used a Monte Carlo simulation method to express statistical variability of the results. In the winter rainy season of the first year, we observed 214 kg-N ha⁻¹ of inorganic N-loss from the root zone within 20 weeks after planting. Plastic mulch was applied 7 weeks after the planting. Marketable fruit yield was 38.7 tons ha⁻¹ with a total N application rate of 255 kg ha⁻¹. Total N uptake of strawberries was ~120 kg ha⁻¹ and about 80% of this occurred in the later half of the growth period. In the second year, we adopted pre-plant plastic mulch application and adjusted basal/supplemental organic fertilizer-N rates to better synchronize the N supply with the plant N demand. As a result, although the residual soil inorganic-N was much lower than the first year, the N-loss during the rainy season was decreased to 13 kg-N ha⁻¹, while maintaining fruit yield a similar to the first year.

INTRODUCTION

Organic strawberry production in California is on the rise. In 2001, 162 organic strawberry growers planted 757 acres of organic strawberries in California. The acreage increased to 1,279 acres in 2002 (Klonsky, 2003).

Strawberry is a nitrogen (N)-sensitive crop. The yield and quality of strawberry fruits is strongly affected by plant N status

2. Cultural practices and fruit yield survey

Strawberry transplants were planted on Nov. 19, 2001 in the first year and on Nov. 26, 2002 in the second year. Plant population per ha was 52,340 in year 1 and 47,340 in year 2. To suppress *Verticillium dahliae* population in soils, we incorporated broccoli residues as a biofumigant (Koike and Subbarao, 2000). The application rate was 2.2 tons ha⁻¹ and 3.9 tons ha⁻¹ (dry matter each) in year 1 and 2, respectively. Total N application rate was 255 kg-N ha⁻¹ in year 1 (compost 55*, pelletized organic fertilizer 82*, broccoli 86*, liquid organic fertilizer 33 kg-N ha⁻¹. * basal application) and 329 kg-N ha⁻¹ in year 2 (compost 66*, pelletized organic fertilizer 37*, broccoli 151*, liquid organic fertilizer 75 kg-N ha⁻¹). See figure 1a (year 1) and 1b (year 2) for timing of each application. Fresh strawberry fruit yield was measured once to twice per week from 40 designated harvest plants in the middle two beds of each plot from May to September in both years. Marketable and cull fruit yields were weighed separately by experienced harvesters. Mortality of strawberry plants was examined from May to the end of the season.

3. N monitoring

Throughout the growth period, samplings were done 3 to 4 times for plants and once a month for soils from 0-60 cm deep. Biomass and T-N content in each part of the strawberry plant, including total fruit biomass and its N content, were determined. Cumulative daily N uptakes were calculated from regression curves of N content and biomass of plants and fruits (Hunt, 1982). Nitrate and ammonium was extracted from soil with 2M KCl and determined by the flow injection analysis method. Soil bulk density in all plots was determined using the soil core method at the end of season each year. Nitrate plus ammonium content was expressed volumetrically as inorganic N (= NO₃-N + NH₄-N kg ha⁻¹). We estimated N-Loss from the root zone during the rainy season (between 0 and 20 weeks after planting, which is approximately from mid November to early April) using the following formula:

$$\text{N-Loss}_{20\text{wks}} (\text{kg ha}^{-1}) = (\text{Soil inorg. N}_0 - \text{Soil inorg. N}_{20\text{wks}}) - (\text{Plant uptake N}_{20\text{wks}} - \text{Plant input N}_0)$$

where, Soil inorg. N₀; Inorganic N (kg ha⁻¹) in 0-30 cm deep soil on the planting day, Soil inorg. N_{20wks}; Inorganic N (kg ha⁻¹) in 0-30 cm deep soil at 20 weeks after planting, Plant uptake N_{20wks}; Plant uptake N (kg ha⁻¹) at 20 weeks after planting, and Plant input N₀; N input (kg ha⁻¹) from transplants. We defined a 0-30cm soil layer in the bed area as the strawberry root zone, based on literature (UC-IPM Program, 1994) and a preliminary strawberry root profile survey. N-Loss is considered to consist mainly of leached-N and denitrified-N with minor contributions from biological immobilized-N. To demonstrate statistical variability, we used the Monte-Carlo simulation method and expressed the results as a mean and 90% credible interval (5%, 95%) (Morgan and Henrion, 1990; Muramoto et al., 2001).

RESULTS

1. The first year (2001-2002)

Pre-plant soil inorganic N analysis revealed 150 kg-N ha⁻¹ of residual inorganic N from the previous crop (fig. 1a). By adding 222 kg-N ha⁻¹ of total basal N, inorganic N content at the time of planting rose to 260 kg-N ha⁻¹. After receiving 280 mm of precipitation, the grower applied plastic mulch in January 2002. In the meantime, a considerable amount of N was lost since planting. It was estimated that 214 (169, 262) kg ha⁻¹ of inorganic N was lost from the root zone within 20 weeks from planting (fig. 1a).

Strawberries grew well and had no major disease problems. Marketable fruit yield and total fruit yield was 39.8 tons ha⁻¹ and 63.1 tons ha⁻¹, respectively. Mortality of plants at the end of the season was 1.2 %. Cumulative daily N uptake of strawberry plants showed that ~80% of the total N uptake took place in the later half of the growth period (fig. 1a). It also indicated a negative accumulation in the late growth stage due probably to N removal from plants through fruits harvests. Total N uptake by strawberry was ~120 kg-N ha⁻¹.

2. The second year (2002-2003)

Based on the results of the first year, the grower modified his N fertility management for the following year. He adopted: 1) pre-planting plastic mulch application to reduce N-loss during the rainy season, and 2) adjusted the rates of basal/supplemental organic fertilizer applications to better meet the N demand of strawberries. He decreased the rate of pre-plant pelletized organic fertilizers to a half of the previous year and increased the rate of liquid fertilizers through fertigation in the late growth stage (fig. 1b).

As a result, the amount of N-loss during the rainy season (Nov. 2002-April 2003) was reduced to 13 (-16, 51) kg-N ha⁻¹ (fig. 1b). It should be noted, however, that the residual N level at the time of planting in the second year was much lower than the first year and precipitation during 0 and 20 weeks after planting in the second year (318 mm) was 15% less than the first year.

The plant development in the second year was earlier than the first year due to warmer temperatures. Two-spotted spider mite (*Tetranychus urticae*) reduced the plant vigor in May and June resulting in relatively smaller plant biomass. Marketable fruit yield was 38.7 tons ha⁻¹, a similar level with the first year, and total fruit yield was

58.4 tons ha⁻¹. Mortality of plants at the end of season was 0.3%. Unlike the first year, cumulative daily N uptake of strawberry plants in the second year did not show a negative accumulation in the late growth stage (fig. 1b). Total N uptake by strawberry was comparable with the first year.




Fig. 1 N loss from an organic strawberry field in Moss landing, California. cv. Aromas. Estimated N-Loss during the rainy season was as follows: year 1; $N\text{-Loss}_{20\text{wks}} = 214 (169, 262)^*$ kg ha⁻¹, and year 2; $N\text{-Loss}_{20\text{wks}} = 13 (-16, 51)^*$ kg ha⁻¹. * mean and 90% credible interval (5%, 95%).

DISCUSSION

The first year's results in this case study showed that organic strawberry production could have a considerable N-loss and a significant impact on the environment during the rainy season (ATTRA, 2003). On the other hand, the second year's results clearly demonstrate that relatively simple adjustments in N management can make a big difference in reducing N-loss while maintaining the fruit yield.

Use of plastic mulch is a common practice in California organic strawberry production (Gliessman et al., 1996), but time of mulch application varies among growers. Though the initial soil inorganic N level was low, our results from the second year indicate the potential of pre-planting plastic mulch application for minimizing N loss during the winter rainy season. Unlike conventional strawberry systems where use of control-release N fertilizers is widespread, no organic control-release N fertilizers that can synchronize the soil N release with the strawberry N demand are currently available. Therefore, a pre-plant plastic mulch application might help reserve inorganic N that was mineralized in the soil during the early rainy season when the plant roots are not fully developed and hence the system is most susceptible to N leaching. Further studies are needed to figure out if interactions exist between timing of mulch application and soil inorganic N level. The soil N monitoring in the first year also suggests the importance of examining residual inorganic N levels in organic systems. Some conventional practices such as this and irrigation management using soil water sensors should be used to enhance N use efficiency in organic systems.

Considering the 10-month growth period of strawberries, the total N uptake of 120 kg ha⁻¹ appears not so high. Limited information is available on N application rates in organic strawberry production. Our N application rate was similar to an example given for sample costs to produce organic strawberries (Bolda et al., 2003), but we might be able to reduce it by furthering synchrony of N supply and N demand, which is the key to improve N use efficiency (Campbell et al., 1995; Woomeer and Swift, 1994). Total fruit yield of the first year of the experiment was 83% of the average total yield of Monterey County in the same year (Monterey County Agricultural Commissioner, 2003).

The N uptake pattern of strawberry plants (fig. 1) suggests that the peak N demand exists in the mid to later growth stages, whereas less N is utilized in the early stage. N fertility studies for organic long-season vegetable production indicate cover crops or compost alone might be inadequate to fulfill the late N demand of these crops (Gaskell and Klauer, 2001; Gaskell et al., 2000). The same is likely to be true in organic strawberry production. Under the current system, fertigation and foliar N application are the only practical options as supplemental N application measures. In the second year of this case study, to meet the N demand of strawberries, the grower decreased the basal N rate of the organic fertilizer and increased the rate of liquid organic fertilizer for the late stage.

This appears to work fine in supplying sufficient N in the late stage, since, unlike the first year, we did not find a negative plant N accumulation in the late season. However, though we did not examine it in this case study, organic fertigation may have a high risk of N leaching under excess irrigation conditions (Pritts and Handley, 1998). Effectiveness of foliar N application in organic strawberry systems is not well studied.

Finally, in spite of a low uptake in this stage, a sufficient level of N in the early growth stage is necessary for bud formation (May and Pritts, 1990). A preliminary study showed a positive correlation between tissue T-N level in the early growth stage (March) and total fruit yield in organic strawberry systems with multiple cultivars (Muramoto, unpublished data). Further studies with different N rates and timings are required to determine the early N demand and appropriate rate and type of pre-plant N application in organic strawberry systems.

ACKNOWLEDGEMENTS

This project was partially funded by the North American Strawberry Growers Association research grant, the Center for Agroecology and Sustainable Food Systems at UC Santa Cruz, the Halliday Foundation, and the Alfred Heller Chair in Agroecology at UC Santa Cruz.

REFERENCES

- ABAConsultants, California State Coastal Conservancy, and Monterey County (Calif.). Planning Dept. 1989. Elkhorn Slough Wetland Management Plan ABA Consultants, Capitola, Calif
- Appropriate Technology Transfer for Rural Areas (ATTRA). 2003. High-input organic agriculture [Online]. Available by <http://attra.ncat.org/attra-pub/organiccrop/tools6.html> <http://attra.ncat.org/attra-pub/organiccrop/tools6.html> (verified 6/1/2004).
- Bolda, M., L. Tourte, K. Klonsky, and J.E. Bervejillo. 2003. Sample costs to produce organic strawberries: Central coast, Santa Cruz and Monterey Counties University of California Cooperative Extension
- Campbell, C.A., R.J.K. Myers, and D. Curtin. 1995. Managing nitrogen for sustainable crop production. *Fertilizer Research* 42:277-296.
- Gaskell, M., and H. Klauer. 2001. Soil nitrate nitrogen and bell pepper yield following application of varying rates of compost or feather meal with and without prior green manure. *Hortscience* 36:473.
- Gaskell, M., B. Fouche, S.T. Koike, W.T. Lanini, J. Mitchell, and R. Smith. 2000. Organic vegetable production in California - Science and practice. *HortTechnology* 10:699-713.
- Gliessman, S.R., M.R. Werner, S.L. Swezey, E. Casswell, J. Cochran, and F. Rosado-May. 1996. Conversion to organic strawberry management: Changes in ecological processes. *California Agriculture* 50:24-31.
- Hunt, R. 1982. Plant growth curves; the Functional approach to plant growth analysis Edward Arnold Limited, London, Great Britain.
- Klonsky, K. 2003. Growth of the organic strawberry industry organic strawberry production: A short course 2003. University of California Sustainable Agriculture Research and Education Program, Davis, CA.
- Koike, S., and K.V. Subbarao. 2000. Broccoli Residues Can Control Verticillium Wilt of Cauliflower. *California Agriculture* 54:30-33.
- May, G., and M. Pritts. 1990. Strawberry nutrition. *Advances in Strawberry Production* 9:10-24.
- Monterey County Agricultural Commissioner. 2003. Monterey County Crop Report 2002.
- Morgan, M.G., and M. Henrion. 1990. Uncertainty: a Guide to dealing with uncertainty in quantitative risk and policy analysis Cambridge University Press, New York, New York, USA.
- Muramoto, J., E.C. Ellis, Z. Li, R.M. Machado, and S.R. Gliessman. 2001. Field-scale nutrient cycling and sustainability: Comparing natural and agricultural ecosystems, p. 121-134, *In* S. R. Gliessman, ed. *Agroecosystem Sustainability: Developing Practical Strategies*. CRC Press, Boca Raton, FL.
- Muramoto, J., S.R. Gliessman, S.T. Koike, D. Schmida, and R. Stephens. 2003. Maintaining agroecosystem health in the organic management of a strawberry/vegetable rotation system. Annual Meeting Abstracts, Agronomy Society of America, Crop Science Society of America, and Soil Science Society of America, Denver, CO.
- Pritts, M., and D. Handley, (eds.) 1998. Strawberry production guide for the Northeast, Midwest, and Eastern Canada, pp. 1-162. Natural Resource, Agriculture, and Engineering Service (NRAES), Ithaca, New York, 14853-5701, USA.
- University of California Integrated Pest Management Program. 1994. Integrated pest management for strawberries University of California Division of Agriculture and Natural Resources, Oakland, Calif.
- Woomer, P.L., and M.J. Swift, (eds.) 1994. The biological management of tropical soil fertility, pp. 1-243. John Wiley & Sons, West Sussex, UK.