

## Ratings Key and Abbreviations

---

**S = Susceptible; MS = Moderately Susceptible; MR = Moderately Resistant; R = Resistant. Rating is based on values in the 90th percentile of all plot-level observations for a given variety.**

**Rating scale for diseases, lodging, and shatter in site-specific summary data. Disease ratings are for area of upper leaves affected, except BYDV, which is area of plot affected. Lodging and shatter are plants affected per plot. Ratings reported in the site-specific summaries are the maximum value observed among replications of a particular variety for a particular trait.**

1 = 0-3%

2 = 4-14%,

3 = 15-29%

4 = 30-49%

5 = 50-69%

6 = 70-84%

7 = 85-95%

8 = 96-100%

**BYDV=Barley Yellow Dwarf Virus**

**Days to Heading & Maturity are from January 1<sup>st</sup> in Davis, CA.**

### **Regions**

**SacV** = Sacramento Valley

**NCSJV** = North-Central San Joaquin Valley

**SoSJV** = South San Joaquin Valley

**ImpV** = Imperial Valley

### **Breeding Programs/Companies**

**AP** = AgriPro

**APB** = Arizona Plant Breeders

**AS** = Allstar Seed

**BAG** = Baglieto Seed

**LCS** = Limagrain Cereal Systems

**NS** = Northern Seed

**OSU** = Oregon State University

**PRL** = Plant Research Limited

**SFS** = Stanislaus Farm Supply

**SY** = Syngenta

**UC** = University of California

**WB** = Westbred-Monsanto

**WWW** = World Wide Wheat

### **Crop Classifications**

#### **Wheat**

**HRS** = Hard Red Spring

**HWS** = Hard White Spring

**HRW** = Hard Red Winter

**HWW** = Hard White Winter

**SRS** = Soft Red Spring

**SWS** = Soft White Spring

**SWW** = Soft White Winter

#### **Barley**

**2RSF** = 2-ROW SPRING FEED

**6RSF** = 6-ROW SPRING FEED

**2RSF(H)** = 6-ROW SPRING FEED HOODED

**6RSF(H)** = 6-ROW SPRING FEED HOODED

**2RS(N)**: NAKED 2-ROW SPRING

**6RS(N)**: NAKED 6-ROW SPRING

**2RSM** = 2-ROW SPRING MALT

**6RSM** = 6-ROW SPRING MALT

**6RWF** = 6-ROW WINTER FEED

**6RW(H)** = 6-ROW WINTER HOODED