



Impacts of different N management regimes on nitrifier and denitrifier communities and N cycling in soil microenvironments

Angela Y.Y. Kong^{a,*}, Krassimira Hristova^b, Kate M. Scow^b, Johan Six^a

^a Department of Plant Sciences, University of California, Davis, One Shields Avenue, Davis, CA 95616, USA

^b Land, Air, and Water Resources Department, University of California, Davis, One Shields Avenue, Davis, CA 95616, USA

ARTICLE INFO

Article history:

Received 16 November 2009

Received in revised form

15 May 2010

Accepted 19 May 2010

Available online 12 June 2010

Keywords:

amoA

nosZ

Soil microenvironments

N transformation rates

Cropping system

ABSTRACT

Real-time quantitative PCR assays, targeting part of the ammonia monooxygenase (*amoA*), nitrous oxide reductase (*nosZ*), and 16S rRNA genes were coupled with ¹⁵N pool dilution techniques to investigate the effects of long-term agricultural management practices on potential gross N mineralization and nitrification rates, as well as ammonia-oxidizing bacteria (AOB), denitrifier, and total bacterial community sizes within different soil microenvironments. Three soil microenvironments [coarse particulate organic matter (cPOM; >250 μm), microaggregate (53–250 μm), and silt-and-clay fraction (<53 μm)] were physically isolated from soil samples collected across the cropping season from conventional, low-input, and organic maize–tomato systems (*Zea mays* L.–*Lycopersicon esculentum* L.). We hypothesized that (i) the higher N inputs and soil N content of the organic system foster larger AOB and denitrifier communities than in the conventional and low-input systems, (ii) differences in potential gross N mineralization and nitrification rates across the systems correspond with AOB and denitrifier abundances, and (iii) *amoA*, *nosZ*, and 16S rRNA gene abundances are higher in the microaggregates than in the cPOM and silt-and-clay microenvironments. Despite 13 years of different soil management and greater soil C and N content in the organic compared to the conventional and low-input systems, total bacterial communities within the whole soil were similar in size across the three systems (~5.15 × 10⁸ copies g⁻¹ soil). However, *amoA* gene densities were ~2 times higher in the organic (1.75 × 10⁸ copies g⁻¹ soil) than the other systems at the start of the season and *nosZ* gene abundances were ~2 times greater in the conventional (7.65 × 10⁷ copies g⁻¹ soil) than in the other systems by the end of the season. Because organic management did not consistently lead to larger AOB and denitrifier communities than the other two systems, our first hypothesis was not corroborated. Our second hypothesis was also not corroborated because canonical correspondence analyses revealed that AOB and denitrifier abundances were decoupled from potential gross N mineralization and nitrification rates and from inorganic N concentrations. Our third hypothesis was supported by the overall larger nitrifier, denitrifier, and total bacterial communities measured in the soil microaggregates compared to the cPOM and silt-and-clay. These results suggest that the microaggregates are microenvironments that preferentially stabilize C, and concomitantly promote the growth of nitrifier and denitrifier communities, thereby serving as potential hotspots for N₂O losses.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

Managed ecosystems have received much attention for their role in nitrogen (N) cycling as the increased rate of mineral N fertilizer application to crop production systems significantly contributes to the global increase of nitrous oxide (N₂O) emissions (Mosier et al., 1998) and N leaching (Foster et al., 1986). A better mechanistic understanding of nutrient cycling within cropping systems – both conventional and alternative – is necessary to promote long-term

crop management practices that enhance agroecosystem services, such as, carbon (C) storage, N retention, and mitigation of greenhouse gas emissions.

Agricultural management practices (e.g., tillage, organic amendment addition, cropping rotation, and irrigation) can strongly influence the size, composition, and activity of the microbial community in soils (Beare et al., 1992; Bossio et al., 1998; Frey et al., 1999). For example, agricultural practices, such as mineral N fertilizer application, significantly alter ammonia-oxidizing bacteria (AOB) populations, which subsequently impact nitrification and production rates of nitrous oxide (N₂O), a potent greenhouse gas, in crop production systems (e.g., Kowalchuk and Stephen, 2001). Cropping

* Corresponding author. Tel.: +1 530 752 6309; fax: +1 530 754 4361.

E-mail address: aykong@ucdavis.edu (A.Y.Y. Kong).

systems that receive high organic matter inputs have been characterized by greater microbial activity as well as greater N mineralization compared to systems receiving only mineral fertilizer-N (Gunapala and Scow, 1998; Kramer et al., 2002). However, if organic matter-C additions occur prior to the peak in N demand by the crops, then this can lead to increased microbial immobilization of NO_3^- (Recous et al., 1990; Sakala et al., 2000), which can reduce N losses from the system via leaching and/or denitrification (Robertson, 1997; Scow, 1997). Consequently, the interactions between nutrient management practices and microbial-mediated N dynamics need to be better understood in order to optimize trade-offs between greater soil organic matter levels and potential N losses in agroecosystems.

Characterizing microbial communities associated with physico-chemically and spatially heterogeneous soil microenvironments may reveal the microbial communities responsible for important ecosystem processes and elucidate how microbial processes scale up from the soil to the ecosystem level. In a comparison between aggregated and unaggregated soils, Sexstone et al. (1988) suggested that the spatial separation of available C, NO_3^- , and denitrifying bacteria was the major factor limiting denitrification in macroaggregated soil. Additionally, Seech and Beauchamp (1988) have shown that denitrifier population sizes and activity increase as the size of aggregates decrease. Contrastingly, others have shown that denitrification (i.e., N_2O production) is positively correlated with intact aggregate size (Drury et al., 2004; Miller et al., 2009). Molecular analyses have shown that the microaggregate environment selects for a microbial community dominated by *Actinobacteria* and higher bacterial abundance than the communities found on the outside of the microaggregate and those associated with macroaggregates (Ranjard et al., 2000; Mummey et al., 2006).

Recent advances in environmental molecular biology techniques have enabled researchers to investigate the relationship between soil management, microbial communities, and ecosystem processes, especially pertaining to the N cycle. For example, by comparing the relative contributions of nitrification and denitrification to total N_2O emissions measured from a ^{15}N incubation experiment with assessments of nitrifier and denitrifier community composition and abundance, Ma et al. (2008) found nitrification was the primary source of N_2O emissions in wetland soils and that cultivation increased nitrifier but not denitrifier abundance. Avrahami et al. (2002) combined N_2O flux measurements with community assays of the *amoA* and *nirK* genes and found, while greater ammonium additions increased nitrification and subsequent N_2O release rates, that the community structure of the ammonia oxidizers was not changed. Relative abundances of terminal-restriction fragments (TRFs) of amplified *nirK* fragments increased, however, with increasing ammonium additions.

Agricultural soils receiving different C and N input, in terms of quality and quantity, are expected to differ in rates of microbial-mediated N transformations and C cycling. The aims of this study were to (i) examine the responses of AOB, denitrifier, and total bacterial abundances to differences in long-term N management in

conventional (annual mineral fertilizer applications), low-input (mineral fertilizer and cover crop applied in alternating years), and organic (annual manure and cover crop additions) cropping systems across one growing season; and (ii) determine whether AOB and denitrifier densities are related to short-term N mineralization and nitrification rates across different soil microenvironments. We addressed these objectives in a long-term agricultural site where Kong et al. (2005) found significantly higher soil C stocks and greater soil aggregation in the organic compared to the conventional and low-input maize–tomato systems (*Zea mays* L.–*Lycopersicon esculentum* L.). We hypothesized that (i) the higher N inputs and soil N content of the organic system foster larger AOB and denitrifier communities than in the conventional and low-input systems, (ii) differences in potential gross N mineralization and nitrification rates across the systems correspond with AOB and denitrifier abundances, and (iii) *amoA*, *nosZ*, and 16S rRNA gene abundances are higher in the microaggregates than in the cPOM and silt-and-clay microenvironments. Real-time quantitative polymerase chain reactions (qPCR) were combined with ^{15}N isotopic pool dilution assays to test our hypotheses.

2. Materials and methods

2.1. Study site, experimental design, and field operations

The field study took place on the Long Term Research in Agricultural Sustainability (LTRAS) plots at the Russell Ranch experimental site (Davis, CA, USA; $38^\circ 32' 24''$ N $121^\circ 52' 12''$ W), which is located in a region characterized by wet winters and hot, dry summers. Two soil types dominate the site: i) Yolo silt loam (fine-silty, mixed, nonacid, thermic Typic Xerorthent) and ii) Rincon silty clay loam (fine, montmorillonitic, thermic Mollic Haploxeralf). The field study was conducted during the 2006 maize growing season in three maize–tomato (*Z. mays* L.–*L. esculentum* L.) cropping systems ($n = 3$): conventional (annual synthetic fertilizer applications), low-input (synthetic N fertilizer applied in alternate years with cover crop-N incorporated the years without synthetic N fertilization) and organic (annual composted manure and cover crop additions) (see description in Table 1). Since 1993, these maize–tomato cropping systems have been arranged in a completely randomized design with three 0.4 ha replicates under furrow irrigation and conventional tillage.

During the winter cover crop (hairy vetch: *Vicia dasycarpa*) growing season, microplots (1.0 × 1.0 m) were established in each cropping system replicate (3 treatments × 3 replicates = 9 microplots). In the final week of April 2006, the winter cover crop biomass in the microplots was mowed and roto-tilled into the microplots to a depth of 15 cm. Maize was direct-seeded into the conventional and then into the low-input and organic plots in the second and final weeks of May, respectively. At the start of May, composted manure was incorporated into the organic cropping system at a rate of 373 kg N ha⁻¹, while in the first week of June, the conventional system received 51 kg N ha⁻¹ as N–P–K starter and 170 kg N ha⁻¹ as

Table 1

Distribution of soil microenvironments and total soil nitrogen (N) levels after cover crop incorporation (T_1 sampling event) and descriptions of N inputs to the conventional, low-input, and organic maize–tomato systems. Lower-case letters indicate significant differences associated with a soil microenvironment effect on the proportions of soil microenvironments across the cropping systems ($p < 0.05$). Total soil N assigned with different upper-case letters are significantly different at the $p < 0.05$ level.

Cropping System	Distribution of Soil Microenvironments			Total Soil N (Mg ha ⁻¹)	N inputs	
	cPOM (>250 μm)	Microaggregate (53–250 μm) (%)	Silt-and-clay (<53 μm)		Even Years of Cropping	Odd Years of Cropping
Conventional	1.3 ^c	40.7 ^b	58.0 ^a	1.41 ^B	NH ₄ –NO ₃ fertilizer	NH ₄ –NO ₃ fertilizer
Low-input	2.0 ^c	33.0 ^b	64.7 ^a	1.33 ^B	winter legume cover crop	NH ₄ –NO ₃ fertilizer
Organic	2.3 ^c	36.0 ^b	61.7 ^a	1.85 ^A	winter legume cover crop and composted manure	winter legume cover crop and composted manure

ammonium nitrate side-dressing. Subsequent simulations of field operations within the microplots (cultivation, weeding, etc.) were done by hand.

2.2. Soil sample collection and aggregate separation

On October 27, 2005 (T_0 sampling event; before seeding of the winter cover crop), two soil cores (4 cm diameter; 0–15 cm) were collected from each microplot, composited, and sub-sampled for baseline soil measurements. During the 2006 maize growing season, soil core samples were collected from the microplots after cover crop incorporation (May 31st; T_i) and at harvest (September 13th; T_h). Soil bulk density was determined on an individual soil core basis. Approximately 10 g of field-moist soil was sub-sampled for moisture content and the remainder was gently broken apart and large debris was removed. Of the latter, 10 g was air-dried, and the remainder was stored at $-20\text{ }^\circ\text{C}$ until thawed at $4\text{ }^\circ\text{C}$ 20 min prior to fractionation or 2 mm sieving for ^{15}N pool dilution assays. Air-dried soil samples were ground and analyzed for elemental N concentrations, using a PDZ Europa ANCA-GSL isotope ratio mass spectrometer (Sercon, Crewe, UK). To extrapolate elemental N values from g N kg^{-1} soil to a hectare-basis, we used the minimum equivalent soil mass to correct for potential temporal changes in soil bulk density (Ellert and Bettany, 1995; Lee et al., 2009).

For each thawed soil sample, 30 g subsamples were fractionated into three fractions: coarse particulate organic matter (cPOM: $>250\text{ }\mu\text{m}$), microaggregate ($53\text{--}250\text{ }\mu\text{m}$), and silt-and-clay ($<53\text{ }\mu\text{m}$), using a microaggregate isolator according to the methodology outlined in Six et al. (2000). Thawed soil samples were submerged in deionized water at room temperature for 5 min to slake the soil. The slaked soil was immersed in deionized water on top of a $250\text{ }\mu\text{m}$ mesh screen and shaken with 50 stainless steel beads (4 mm diameter) until only cPOM and sand were retained on the $250\text{ }\mu\text{m}$ mesh screen. During shaking, a continuous and steady stream of water flowed through the device to ensure that microaggregates were immediately flushed onto a $53\text{ }\mu\text{m}$ sieve and were not exposed to any further disruption by the beads. The material on the $53\text{ }\mu\text{m}$ sieve was manually sieved by moving the sieve in an up-and-down motion 50 times over a 2 min period (Elliott, 1986), to separate water-stable microaggregates from silt-and-clay particles. All fractions were collected as separate soil suspensions, which were centrifuged at 5000 rpm for 15 min at $4\text{ }^\circ\text{C}$ (Sorvall RC-5C Plus Superspeed centrifuge, Thermo Scientific). The supernatant was discarded and the remaining soil was lyophilized and stored at $-20\text{ }^\circ\text{C}$ until further analysis. The different fractions served as the soil microenvironments of interest.

2.3. ^{15}N isotopic pool dilution assays

The remainder of the thawed soil samples was 2 mm-sieved and air-dried for extractable mineral N and N transformation rate analyses. For extractable N measurements, 10 g subsamples of air-dried soil were extracted with 50 ml of 2 M KCl solution by shaking on a reciprocal shaker for 1 h and filtering through a Whatman No. 42 filter paper (ashless). The $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$ concentrations of the soil extracts were analyzed colorimetrically by the Berthelot reaction for NH_4^+ (Forster, 1995), and by vanadium (III) chloride reduction for NO_3^- (Doane and Horwath, 2003).

Potential gross mineralization and nitrification rates were measured using ^{15}N pool dilution techniques (Barraclough, 1991). Four 10 g replicates of air-dried soil were brought to 20% water holding capacity (WHC) in 120 ml specimen cups. After equilibrating for 12 h to avoid the period of stimulated microbial activity upon re-wetting air-dried soil (Birch, 1958), solutions of ($^{15}\text{NH}_4$) $_2\text{SO}_4$ (99 atom%) for potential gross mineralization incubations and

K^{15}NO_3 (99 atom%) for potential gross nitrification incubations were added to two replicate samples, respectively. The addition of the ^{15}N solutions resulted in an approximate soil moisture level of 60% WHC. The quantity of ^{15}N added to each replicate was adjusted to achieve approximately 10 atom% ^{15}N enrichment of the NH_4^+ and NO_3^- pool sizes estimated from the extractable N measurements described above. All replicates were incubated at room temperature ($25\text{ }^\circ\text{C}$). For the potential gross mineralization assay, two replicates were destructively sampled after 3 h (t_0) and 1 d (t_1); whilst for the potential gross nitrification assay, the two remaining replicates were destructively sampled after 3 h (t_0) and 3 d (t_3). The different destructive sampling times for the two assays reflect differences in optimal incubation times that avoided re-mineralization of added N, as determined by preliminary experiments. After destructive sampling, samples were extracted with 2 M KCl, and the quantities of $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$ were measured as described above. The ^{15}N isotopic signatures of $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$ were determined by diffusion onto acidified disks (Stark and Hart, 1996) and analyzed with a PDZ Europa 20–20 isotope ratio mass spectrometer (Crewe, United Kingdom; Stable Isotope Facility of the University of California, Davis).

Rates for potential gross mineralization were calculated using the following equation (Barraclough, 1991):

$$m = \theta \times \left[\frac{\log(A_0^*/A_t^*)}{\log[(1 + \theta t)/C_0]} \right] \quad (1)$$

where m = potential gross mineralization rate, θ = the rate of change in the size of the NH_4^+ pool, A_0^* = the ^{15}N atom% excess of the NH_4^+ pool at t_0 , A_t^* = the ^{15}N atom% excess of the NH_4^+ pool at t_t , and C_0 = the size of the NH_4^+ pool at t_0 . Potential gross nitrification rates were also calculated using Equation (1), but with ^{15}N enrichments, incubation times, and pool size changes associated with the NO_3^- pool.

2.4. Soil DNA extraction procedures

Genomic DNA was extracted from soil microenvironment samples using the Bio 101 Fast DNA SPIN kit for soil (BIO 101, Vista, CA). First, humic acid, a PCR inhibitor, was removed from the soil microenvironment sample as follows: 1 g of soil was incubated in 2 ml of 0.1% $\text{Na}_4\text{P}_2\text{O}_7$ in 10 mM Tris–HCl buffer (pH 8.0)–1 mM ethylenediaminetetraacetic acid (EDTA), at room temperature for 30 min, centrifuged at $8500\times g$ for 10 min at room temperature, and the supernatant was discarded. Total DNA was extracted from 0.5 g humic acid-washed sample according to the manufacturer's instructions (BIO 101, Vista, CA). Extracted DNA was visualized by agarose (1.5% [wt/vol]) gel electrophoresis. DNA yield was quantified spectrofluorometrically using the Quant-iT fluorescent dye (Invitrogen, Carlsbad, CA).

2.5. Determination of 16S rRNA, *amoA*, and *nosZ* gene abundances

Abundances of total bacteria were determined using the real-time quantitative TaqMan PCR method as described in Suzuki et al. (2001). Real-time qPCR was performed in a 20 μl reaction mixture that consisted of 4 μl of template DNA (SOM fractions: $\sim 2.5\text{ ng }\mu\text{l}^{-1}$; Whole soil: $\sim 1.1\text{ ng }\mu\text{l}^{-1}$), 10 μl of TaqMan Universal PCR Master Mix (Applied Biosystems, NJ, USA), 4 μl of H_2O , 0.8 μl of each primer, and 0.4 μl of the probe. The primers BACT1369F (5'-CGG TGA ATA CGT TCY CGG-3'), PROK1492R (5'-GGW TAC CTT GTT ACG ACTT-3') and probe TM1389 (5'-CTT GTA CAC ACC GCC CGTC-3') (Suzuki et al., 2001) were used at concentrations of 800 nM, 800 nM, and 200 nM, respectively. The PCR protocol for bacterial 16S rRNA quantification was as follows: 2 min at $50\text{ }^\circ\text{C}$, 10 min at $95\text{ }^\circ\text{C}$, and then 40 cycles consisting of 15 s at $95\text{ }^\circ\text{C}$, and 1 min at $56\text{ }^\circ\text{C}$. We acknowledge that

novel bacterial sequences that were not targeted by the TaqMan probe that was used in this study to determine 16S rRNA gene abundances might have resulted in an underestimation of the total bacterial community size in our soil, which might subsequently have led to the calculation of higher relative abundances of the functional genes (e.g., *amoA* and *nosZ*).

Ammonia-oxidizing and denitrifying communities were investigated by quantifying the abundances of the functional genes, *amoA*, which encodes for the active subunit of ammonia monooxygenase (Rotthauwe et al., 1997), and *nosZ*, which encodes for nitrous oxide reductase (Henry et al., 2006). Abundances of AOB and denitrifiers were both determined using the SYBR green real-time qPCR method. As described by Cavagnaro et al. (2007) for the *amoA* qPCR reaction, 1.2 μl of the A189 forward primer (5'-GNG ACT GGG ACT TCT GG-3'; 0.3 μM ; Holmes et al., 1995) and 3.6 μl of the *amoA*-2R' reverse primer (5'-CCC CTC KGS AAA GCC TTC TTC-3'; 0.9 μM ; Rotthauwe et al., 1997) were used to amplify the *amoA* gene in a 20 μl reaction mixture with 5 μl of template DNA (SOM fractions: $\sim 1.6 \text{ ng } \mu\text{l}^{-1}$; Whole soil: $\sim 1.0 \text{ ng } \mu\text{l}^{-1}$). The PCR conditions were as follows: 15 s at 95 $^{\circ}\text{C}$, and then 40 cycles consisting of 15 s at 95 $^{\circ}\text{C}$, 30 s at 55 $^{\circ}\text{C}$, and 31 s at 72 $^{\circ}\text{C}$, followed by a dissociation stage of 15 s at 95 $^{\circ}\text{C}$, 30 s at 60 $^{\circ}\text{C}$, and 15 s at 95 $^{\circ}\text{C}$. The real-time qPCR *nosZ* assay was carried out according to modifications by Henry et al. (2006) in a reaction volume of 20 μl , and the assay mixture contained 10 μl of 2 \times ABI Power SYBR I Green PCR Master Mix, 0.8 μl of each *nosZ* primer (*nosZ*2F: 5'-CGC RAC GGC AAS AAG GTS MSS GT-3', 0.3 μM and *nosZ*2R: 5'-CAK RTG CAK SGC RTG GCA GAA-3', 0.3 μM), and 5 μl of template DNA (SOM fractions: $\sim 2.1 \text{ ng } \mu\text{l}^{-1}$; Whole soil: $\sim 1.0 \text{ ng } \mu\text{l}^{-1}$). Thermal cycling conditions for the *nosZ* primers were as follows: an initial cycle of 95 $^{\circ}\text{C}$ for 10 min, then six cycles of 95 $^{\circ}\text{C}$ for 15 s, 65 $^{\circ}\text{C}$ for 30 s, 72 $^{\circ}\text{C}$ for 30 s, followed by 40 cycles of 95 $^{\circ}\text{C}$ for 15 s, 60 $^{\circ}\text{C}$ for 15 s, 72 $^{\circ}\text{C}$ for 30 s, and 83 $^{\circ}\text{C}$ for 30 s (acquisition data step), and, finally, one cycle at 95 $^{\circ}\text{C}$ for 15 s and 60 $^{\circ}\text{C}$ for 30 s, to 95 $^{\circ}\text{C}$ for 15 s.

Fluorescence signals were used to calculate C_T (threshold cycle) values for the *amoA*, *nosZ*, and 16S rRNA gene copies using the internal software of the thermocycler (7300 System SDS software, V1.2.2, Applied Biosystems, CA, USA). The parameter, C_T (threshold cycle), is the cycle number at which the fluorescence emission crosses a threshold within the logarithmic increase phase.

We acknowledge that the high clay and high humic acid content associated with the silt-and-clay and the cPOM, respectively, may have affected DNA extraction efficiency and may have inhibited the qPCR reactions. We also acknowledge that an increase in a specific subpopulation of nitrifiers or denitrifiers in the whole soil or SOM fractions may not have been detected against a larger *amoA*- or *nosZ*-containing population that remained inactive. However, because 16S rRNA gene abundances were detected in all SOM fractions and in the whole soil, we are confident that our methods did not bias the gene abundance results.

2.6. Standard curve and quantification of *amoA*, *nosZ*, and 16S rRNA genes

External standard curves illustrating the relationship between *amoA*, *nosZ*, and 16S rRNA copy numbers and respective C_T values were generated with 10-fold serial dilutions of known copy numbers of the *amoA*, *nosZ*, and 16S rRNA genes, respectively. DNA from three plasmids containing *amoA*, *nosZ*, and 16S rRNA gene fragments amplified from *Nitrosomonas europaea* ATCC 19718, *Bradyrhizobium japonicum* USDA 110, and *Escherichia coli* K-12, respectively, with the forward and reverse primers for *amoA*, *nosZ*, and 16S rRNA (listed above) were extracted with a Plasmid Mini Kit (Qiagen). Plasmid concentrations ($\text{ng } \mu\text{l}^{-1}$) were measured spectrofluorometrically using the Quant-iT fluorescent dye method. Copy numbers of *amoA*,

nosZ, and 16S rRNA were directly calculated from the respective concentrations of extracted plasmid DNA. The following equations were used to convert C_T values to *amoA*, *nosZ*, and 16S rRNA copy numbers, respectively:

$$C_T = -3.37 \times \text{Log}_{10}(\text{amoA copies reaction}^{-1}) + 41.47 \text{ and } R^2 = 0.98 \quad (2)$$

$$C_T = -3.35 \times \text{Log}_{10}(\text{nosZ copies reaction}^{-1}) + 38.67 \text{ and } R^2 = 0.99 \quad (3)$$

$$C_T = -3.54 \times \text{Log}_{10}(\text{16S copies reaction}^{-1}) + 41.93 \text{ and } R^2 = 0.99 \quad (4)$$

Standard curves were linear over six orders of magnitude and the detection limit was approximately 4.0×10^5 and 2.0×10^5 copies g dry soil $^{-1}$ for the *amoA* and *nosZ* real-time qPCRs, respectively, and $\sim 1.0 \times 10^5$ copies g dry soil $^{-1}$ for the 16S rRNA real-time qPCR.

2.7. Data analyses

Whole soil N measurements and the distribution of aggregate size classes within T_i were analyzed using analyses of variance (ANOVA) by the PROC MIXED procedure of the Statistical Analysis System (SAS; SAS Institute, 2002). To compare differences in *amoA*, *nosZ*, and 16S rRNA gene abundances among the three sampling events, repeated measures analyses were performed using the PROC MIXED procedure. The data were analyzed as a completely randomized design, with cropping system as the main plot and plot as a random factor. A standard variance components covariance structure was specified with the TYPE = VC option in the model statement. Differences between means were calculated based on least significant difference tests, with the PDIF option of the LSMEANS statement. Letters for mean separation in PROC MIXED were assigned using the macro PDMIX 800 (Saxton, 1998). All differences discussed were significant at the $p < 0.05$ probability level, unless otherwise stated.

To assess the potential effects of the fractionation/wet-sieving procedure on recoveries of microbial populations in the SOM fractions, the gene copy numbers for *amoA*, *nosZ*, and 16S rRNA in the three soil microenvironments corresponding to the same soil sample, respectively, were summed together (i.e., copies g $^{-1}$ soil for cPOM + microaggregates + silt-and-clay = summed copies for soil sample) and then compared to their corresponding gene abundances in the whole soil (for all soil samples). We found that recoveries in the SOM fractions were not drastically different from abundances in the whole soil. Although we did not expect the sum of the gene abundances in the SOM microenvironments to be identical to the abundances in the whole soil, we are confident that the wet-sieving procedure did not bias the gene abundances in the SOM fractions.

Gene abundances for *amoA*, *nosZ*, and 16S rRNA genes and correlations with environmental variables (NH_4^+ and NO_3^- concentrations and potential gross mineralization and nitrification rates) were analyzed with canonical correspondence analysis (CCA) (Lepš and Šmilauer, 2003), using the CANOCO software (Microcomputer Power, Ithaca, NY, USA). The Monte Carlo permutation test was used to test the significance of the correlations between microbial community gene abundances and measured environmental variables.

Initially, we intended to measure denitrification rates to compare with *nosZ* gene abundances. However, in the process of re-analyzing soils due to unforeseen set-backs, the amount of each soil sample dwindled and, as a result, we did not have enough soil to accurately conduct the denitrification rate measurements following

aggregate separation, baseline soil measurements, three gene assays, and mineralization plus nitrification rate assays for each soil sample. Nevertheless, we feel that the molecular assays and N transformation rate experiments that were conducted encompass a wide breadth of N cycling processes.

3. Results

3.1. Physical and chemical soil properties

Across the three sampling events, total soil N levels were highest in the organic (1.85 Mg N ha⁻¹) and similar between the conventional and low-input systems (1.41 and 1.33 Mg N ha⁻¹, respectively). Total soil N levels were greater at *T*₀ than at *T*_i and *T*_h; however, because trends in N levels were similar across the sampling events within each of the soil microenvironments and N level distribution among the soil microenvironments at *T*_h were similar to *T*_i, only total soil N concentrations at *T*_i are shown in Table 1. The distributions of soil microenvironments were similar between the cropping systems and did not change from *T*₀ to *T*_h. At *T*_i, the silt-and-clay fraction (<53 μm) was the greatest by weight proportion across the systems, representing 58.0%–64.7% of the soil mass, while the microaggregate (53–250 μm) and cPOM (>250 μm) fractions comprised 36.6% and 1.9%, respectively, across the systems (Table 1).

Ammonium concentrations were significantly affected by cropping system and a cropping system × sampling event interaction, while only sampling event influenced NO₃⁻ concentrations (Fig. 1). At *T*₀, NH₄⁺ concentrations were higher in the organic (5.0 μg N g⁻¹ dry soil) than in the conventional and low-input systems (2.9 and 2.2 μg N g⁻¹ dry soil). After cover crop incorporation into each cropping system and immediately before the application of either

mineral- or composted manure-N, NH₄⁺ levels at *T*_i were greater than at *T*₀ (5.2 versus 3.4 μg N g⁻¹ dry soil). Concentrations of NH₄⁺ at *T*_i decreased in the following order: conventional ≥ organic ≥ low-input (6.2, 5.0, and 4.4 μg N g⁻¹ dry soil, respectively). At *T*_h, NH₄⁺ concentrations were the highest of the season and were greatest in the organic (13.3 μg N g⁻¹ dry soil), intermediate in the conventional (10.0 μg N g⁻¹ dry soil), and lowest in the low-input (5.6 μg N g⁻¹ dry soil). Nitrate pools were at least twice as large as ammonium pools in the three cropping systems. Unlike NH₄⁺ levels, NO₃⁻ concentrations peaked at *T*_i (41.2 μg N g⁻¹ dry soil) and were lower at *T*₀ and *T*_h (23.1 and 20.1 μg N g⁻¹ dry soil, respectively).

3.2. Potential gross N transformation rates

Interactions between cropping systems and sampling events were observed for potential gross N transformation rates in the whole soil. Potential gross mineralization and nitrification rates were similar in magnitude despite the larger pool size of NO₃⁻ than NH₄⁺ (Fig. 1). At *T*₀, gross mineralization rates were not different among the conventional, low-input, and organic cropping systems (~3.42 μg N g⁻¹ dry soil day⁻¹). At *T*_i, shortly after incorporation of the winter cover crop biomass, gross mineralization rates were fastest in the conventional system (6.05 μg N g⁻¹ dry soil day⁻¹) and slowest in the organic system (1.53 μg N g⁻¹ dry soil day⁻¹). By the end of the season, the gross mineralization rates were 9.57, 5.27, and 3.89 μg N g⁻¹ dry soil day⁻¹, in the organic, low-input, and conventional systems, respectively. Potential gross nitrification rates were similar between the low-input and conventional systems (~2.56 μg N g⁻¹ dry soil day⁻¹) at *T*₀, *T*_i, and *T*_h, while rates for the organic systems (~4.34 μg N g⁻¹ dry soil day⁻¹) were generally the highest (but only significantly higher at *T*₀; Fig. 1).




Fig. 1. Ammonium (A) and nitrate concentrations (B) (μg N g⁻¹ dry soil) and potential gross mineralization (C) and nitrification rates (D) (μg N g⁻¹ dry soil day⁻¹) for the conventional, low-input, and organic cropping systems, before the start of the cover crop growing season (*T*₀), after cover crop incorporation (*T*_i), and during maize harvest (*T*_h). Bars represent means (*n* = 3). Different letters within the respective figures for ammonium concentrations, gross mineralization rates, and gross nitrification rates indicate differences associated with a significant sampling event × cropping system effect (*p* < 0.05). For nitrate concentrations, different letters symbolize differences among sampling events (*p* < 0.05).

Table 2

Ratios of potential gross nitrification rates to gross mineralization rates across the conventional, low-input, and organic cropping systems, before the start of the cover crop growing season (T_0), after cover crop incorporation (T_i), and during maize harvest (T_h) ($n = 3$). Values with different letters are significantly different at the $p < 0.05$ level.

Sampling Event	Cropping System		
	Conventional	Low-input	Organic
T_0	1.0 (bc)	0.75 (c)	1.78 (b)
T_i	0.46 (c)	0.69 (c)	4.0 (a)
T_h	0.72 (c)	0.48 (c)	0.29 (c)

The ratio of gross nitrification rates over gross mineralization rates (Nit:Min) were calculated to assess competition between heterotrophic microorganisms and autotrophic nitrifiers for NH_4^+ (Table 2). In these soils, Nit:Min ratios were generally low, ranging from 0.29 to 4.0, with the Nit:Min ratio for the organic system rising from T_0 (1.78) to T_i (4.0) and falling from T_i to T_h (0.29). Cropping systems did not clearly influence the ratio of Nit:Min across the different sampling events.

3.3. Total bacterial, AOB, and denitrifier community abundances

Total copy numbers for 16S rRNA genes averaged across the cropping systems increased from 1.06×10^8 copies g^{-1} soil at T_0 to 3.34×10^8 and 7.70×10^8 copies g^{-1} soil at T_i and T_h , respectively. The size of ammonia-oxidizing and denitrifying bacterial communities changed significantly over the course of the season, but not congruently. Total *amoA* gene abundances among cropping systems at T_0 and T_i were ~ 6 and ~ 2 times higher in the organic (1.63×10^8 and 1.75×10^8 copies g^{-1} soil, respectively) than the conventional and low-input systems, respectively; however at T_h , densities of *amoA* genes did not differ among the cropping systems (Table 3). Total *nosZ* copy numbers were similar among all cropping systems at T_i , but were also the lowest across the season (4.39×10^6 copies g^{-1} soil). Copies of *nosZ* were greater in the conventional and low-input systems than in the organic at T_h . Copy numbers of *nosZ* at the end of the season in the conventional and low-input systems (T_h) were also higher than at T_0 , and T_i , whereas the denitrifier abundances in the organic system were not significantly different throughout the season (Table 3).

A significant interaction between cropping system, soil micro-environment, and sampling event effects was found for the *amoA*, *nosZ*, and 16S rRNA gene copy numbers of the soil microenvironments. Across the season, a cropping system effect was only observed for the copy numbers of *amoA*, *nosZ*, and 16S rRNA genes in the microaggregate fraction (Figs. 2 and 3, and 4). Also, copy numbers associated with the microaggregate fraction were higher than gene abundances in the cPOM and silt-and-clay. Total bacterial abundance in the silt-and-clay was on average 2.79×10^7 copies g^{-1} soil, while copy numbers of *amoA* and *nosZ* in silt-and-clay ranged from 0 or below detection limit to 3.88×10^6 copies g^{-1} soil and 0 or below

detection limit to 4.38×10^7 copies g^{-1} soil, respectively. In the cPOM, copy numbers of *amoA*, *nosZ*, and 16S rRNA were on average 2.77×10^5 , 1.09×10^6 , and 5.32×10^7 copies g^{-1} soil, respectively.

3.4. Canonical correspondence analysis

The first two canonical axes of the CCA of *amoA*, *nosZ*, and 16S rRNA gene abundances, NH_4^+ and NO_3^- concentrations, and potential gross mineralization and nitrification rates at T_0 , T_i , and T_h explained 35.6% of the variance in the gene abundance data and 100% of variance in the total gene abundance-environment parameter relationship (Fig. 5). The first canonical axis, which explained 85.5% of the total gene abundance-environmental parameter variation ($p = 0.006$), was dominated mainly by NH_4^+ concentrations (0.48) and, to a smaller degree, by NO_3^- concentrations (0.20). The second canonical axis explained an additional 14.5% of the total gene abundance-environmental parameter variation ($p = 0.006$) and was also dominated by the effects of NO_3^- (0.27) and NH_4^+ concentrations (0.17), indicating that NO_3^- and NH_4^+ concentrations are the most influential environmental gradients, of those measured, on the distribution of nitrifier and denitrifier genes among samples. However, when sampling event was included as a covariable in the CCA, canonical axes 1 (x axis) and 2 (y axis) explained 98.2% and 1.8% of the gene abundance-environmental variable variance in the data, respectively, but the inclusion of seasonality in the CCA rendered the axes insignificant (data not shown).

4. Discussion

4.1. Long-term management effects on N transformation rates and AOB, denitrifier, and total bacterial community dynamics

For the past 13 years, cover crop- and composted manure-N have been incorporated into the organic system at the Russell Ranch at rates nearly two and five times greater than the rates of mineral N and mineral plus cover crop-N input to the conventional and low-input systems, respectively. It was expected that inorganic N availability and the results from the isotope dilution experiments would mirror these input differences between the cropping systems. However, short-term NH_4^+ and NO_3^- concentrations and potential gross N transformation rates did not demonstrate a consistently greater soil N supplying capacity for the organic compared to the conventional and low-input systems. At T_0 , which took place after the 2005 growing season, NH_4^+ levels were at their lowest, likely due to the removal of plant residues and the lack of irrigation at that time of the year. In addition, NH_4^+ levels were lower in the low-input and conventional systems than in the organic system, which has been suggested to release NH_4^+ more gradually than conventional systems (Burger and Jackson, 2003). Several months after mineral and/or composted manure N incorporation into the systems, NH_4^+ concentrations were the greatest for the season and reflected the relative rates of N input to the

Table 3

Total copy numbers of 16S rRNA, *amoA*, and *nosZ* genes measured from whole soil, before the start of the cover crop growing season (T_0), after cover crop incorporation (T_i), and during maize harvest (T_h) ($n = 3$). Values for *amoA* and *nosZ* with different upper-case and lower-case letters are significantly different within respective gene abundances ($p < 0.05$). For 16S rRNA gene copy numbers, values followed by different lower-case letters are significantly greater than values for other sampling events, within a cropping system ($p < 0.05$).

Cropping System Sampling Event	16S rRNA			<i>amoA</i>			<i>nosZ</i>		
	Conventional	Low-input	Organic	Conventional	Low-input	Organic	Conventional	Low-input	Organic
	(No. gene copies g^{-1} soil)								
T_0	1.22×10^8 (xy)	2.48×10^8 (xy)	2.25×10^8 (xy)	2.48×10^7 (C)	2.72×10^7 (C)	1.63×10^8 (A)	3.93×10^7 (bc)	4.94×10^6 (d)	7.71×10^6 (d)
T_i	8.54×10^8 (y)	6.01×10^7 (y)	8.81×10^7 (y)	9.34×10^7 (B)	6.40×10^7 (BC)	1.75×10^8 (A)	4.43×10^6 (d)	5.55×10^6 (d)	3.20×10^6 (d)
T_h	1.25×10^9 (x)	8.80×10^8 (x)	1.83×10^8 (x)	5.57×10^7 (BC)	2.98×10^7 (C)	3.44×10^7 (C)	7.65×10^7 (a)	5.58×10^7 (ab)	8.83×10^6 (d)




Fig. 2. Mean copy numbers of 16S rRNA genes within coarse particulate organic matter (cPOM: >250 μm), microaggregate (53–250 μm), and silt-and-clay (<53 μm) soil microenvironments of the conventional (A, D, G), low-input (B, E, H), and organic (C, F, I) cropping systems, before the start of the cover crop growing season (T_0), after cover crop incorporation (T_i), and during maize harvest (T_h) ($n = 3$). Bars shown with different letters are significantly different across soil microenvironments, cropping systems, and sampling events ($p < 0.05$).

systems (organic > conventional > low-input). In contrast, NO_3^- levels peaked at T_i and were not different among the systems. Under incubation conditions, Burger and Jackson (2003) found that NO_3^- immobilization by heterotrophic microbes was greater in organic ($\sim 2.5 \mu\text{g N g}^{-1} \text{d}^{-1}$) than conventionally managed soil ($\sim 0.5 \mu\text{g N g}^{-1} \text{d}^{-1}$), as greater NO_3^- immobilization would be expected for systems with high soil C content. The reasoning for the lack of differences in NO_3^- concentrations among the cropping systems, as well as the discrepancies between dynamics observed in the NH_4^+-N - and $\text{NO}_3^- -\text{N}$ levels in this study are unclear.

Potential gross N mineralization and nitrification rates in the organic system, which received more organic matter-N inputs than the other systems, were not consistently higher than N transformation rates in the conventional and low-input systems; only at T_h were potential gross mineralization rates higher in the organic system than in the low-input and conventional systems. The magnitude of the gross mineralization rates (Fig. 1), which were nearly two times faster than the $<3 \mu\text{g N g}^{-1} \text{soil day}^{-1}$ reported in Burger and Jackson (2003) for the same conventional and organic cropping systems, implied that the NH_4^+ pool turned over daily. The discrepancies between our gross mineralization rates with those from Burger and Jackson (2003) may be related to our shorter incubation times (1 day versus 5 days). On the other hand,

gross nitrification rates measured in this study were similar ($\sim 3 \mu\text{g N g}^{-1} \text{soil day}^{-1}$) to the rates reported in Burger and Jackson (2003). Gross nitrification rates for the systems reflected the overall trends in soil N content among the systems at T_0 (e.g., organic > low-input \geq conventional). However, like Burger and Jackson (2003), who found that gross nitrification rates were not statistically different between the conventional and organic soils, gross nitrification rates in our study became more similar among the three cropping systems over the course of the season. The lack of dramatic differences in potential gross nitrification rates among the systems at T_i and T_h , suggested that, despite different quantities and qualities of N input, NH_4^+ levels were adequate in all three systems. In addition, the lack of differences suggested that the nitrification activity of the AOB populations was not limited by the supply of NH_4^+ or the soil conditions created by the additions of the different N inputs.

We postulated that AOB, denitrifier, and total bacterial community sizes would vary according to differences in the magnitudes of the measured inorganic N and soil C levels of the cropping systems (i.e., organic > low-input = conventional) and would correlate to the potential gross N transformation rates. In a long-term study conducted on soil types similar to those at Russell Ranch, Gunapala and Scow (1998) found that C and manure-C inputs led to higher microbial biomass in organic compared to conventional cropping




Fig. 3. Mean copy numbers of the *amoA* gene within coarse particulate organic matter (cPOM: >250 μm), microaggregate (53–250 μm), and silt-and-clay (<53 μm) soil micro-environments of the conventional (A, D, G), low-input (B, E, H), and organic (C, F, I) cropping systems, before the start of the cover crop growing season (T_0), after cover crop incorporation (T_i), and during maize harvest (T_h) ($n = 3$). Bars shown with different letters are significantly different across soil microenvironments, cropping systems, and sampling events ($p < 0.05$).

systems. Shi and Norton (2000) showed that available NH_4^+ limits both nitrification rates and the nitrifier population size. In our study, gross nitrification rates only weakly reflected NH_4^+ concentrations, while the abundances of *amoA* gene copies in the whole soil reflected soil C trends at T_0 and T_i , but did not mirror measured NH_4^+ levels or potential gross nitrification rates. For example, copies of *amoA* genes at T_i were intermediate in the conventional system (Table 3), but NH_4^+ levels were similar to the organic and higher than the low-input system, respectively (Fig. 1). The lower AOB abundances in the conventional compared to the organic system at T_i and the similar AOB abundances between these two systems at T_h do not concur with results from Okano et al. (2004). There, they found AOB communities to be significantly larger in annually fertilized than in unfertilized soil and suggested that ammonium fertilization has a long-term effect on the population sizes of AOB. Nevertheless, *amoA* gene abundances in this study are on the same order of magnitude as their study on the same maize-tomato systems (0–15 cm). The higher ratios of Nit:Min that we calculated for the organic than conventional system suggested that autotrophic nitrifiers were more competitive for NH_4^+ than the heterotrophic microorganisms (Hart et al., 1994). This result suggested that AOB gene copies should have been higher in the organic than the conventional and low-input systems, which was indeed found to be the case at T_0 and T_i .

It was expected that the higher organic C inputs from the cover crop and composted manure additions into the organic system would lead to a greater abundance of *nosZ* gene copies, because organic fertilizers contain readily available C compounds, which can stimulate denitrification (Paul and Beauchamp, 1989; Rochette et al., 2000). Dambreville et al. (2006) reported that the denitrifying community, determined using the *narG* and *nosZ* genes as molecular markers in restriction fragment length polymorphism (RFLP) assays, was more diverse in the plot amended with organic (composted pig manure) fertilizer than the plot amended with mineral (ammonium nitrate) fertilizer. Using qPCR to quantify the *nosZ* gene, Hallin et al. (2009) found nearly one order of magnitude more *nosZ* gene copies in systems receiving organic fertilizer (solid cattle manure and sewage sludge) compared to the system receiving only ammonium sulfate. Hence, the lower abundance of *nosZ* gene copies across the season in the organic system was unexpected. Furthermore, the lower abundance of *nosZ* gene copies in the organic cropping system did not reflect NO_3^- concentrations, which were not different across the cropping systems. Interestingly, the denitrifier abundances increased in the conventional and low-input systems across the season, but remained lower and did not change in the organic system. We found that the ratios of *nosZ*:16S rRNA gene abundances within the systems did not significantly change across the season




Fig. 4. Mean copy numbers of the *nosZ* gene within coarse particulate organic matter (cPOM: >250 μm), microaggregate (53–250 μm), and silt-and-clay (<53 μm) soil microenvironments of the conventional (A, D, G), low-input (B, E, H), and organic (C, F, I) cropping systems, before the start of the cover crop growing season (T_0), after cover crop incorporation (T_i), and during maize harvest (T_h) ($n = 3$). Bars shown with different letters are significantly different across soil microenvironments, cropping systems, and sampling events ($p < 0.05$).

(data not shown), thereby implying that the size of the denitrifier communities increased along with the total bacterial community. Complex interactions between soil water content, O_2 content, availability of C, and NO_3^- -N levels are understood to be environmental factors that affect microbiological denitrification (Firestone, 1982; Tiedje, 1988). Soil moisture and NO_3^- -N levels, were similar across the three cropping systems. Hence, in our study, the increase in denitrifier abundances in the conventional and low-input systems compared to the lack of change in the organic systems, may be linked more to the effects of C availability on the denitrifier community, as C availability has been recognized as one of the most important regulatory factors controlling denitrification in soil (Burford and Bremner, 1975; Weier et al., 1993).

4.2. Soil microenvironments select for N cycling communities

Our *amoA*, *nosZ*, and 16S rRNA gene abundance data agree with other reports that suggest the location of soil microorganisms within the soil matrix can affect their metabolism and activity (Gupta and Germida, 1988; Lensi et al., 1995; Philippot et al., 1996; Ranjard et al., 2000). Ammonia oxidizing, denitrifier, and total bacterial communities were generally larger in the microaggregate (Figs. 2–4)

than in the cPOM (>250 μm) and silt-and-clay. This was not surprising as microaggregates are important in the protection and stabilization of C (Jastrow, 1996; Six et al., 1998; Gale et al., 2000) and might provide better physical protection for microorganisms than the cPOM and silt-and-clay (Ranjard and Richaume, 2001). Moreover, C availability has been shown to increase rates of denitrification (Beauchamp et al., 1989; McCarty and Bremner, 1992; Luo et al., 1999), and especially within the more anoxic environment of aggregate structures (Sexstone et al., 1985; Parkin, 1987; Drury et al., 2004).

Our results suggest that the microaggregate is a microenvironment that is more conducive to microbial proliferation and nitrifying and denitrifying activity than the cPOM and silt-and-clay. Consequently, our hypothesis that *amoA*, *nosZ*, and 16S rRNA gene abundances are higher in the microaggregate than in the cPOM and silt-and-clay fractions was corroborated. Furthermore, significant cropping system \times soil microenvironment effects found for the gene abundances indicate that crop management interacts with the aggregates in selecting for nitrifiers, denitrifiers, and total bacterial communities. Larger denitrifier populations in the microaggregates of the organic and low-input compared to the conventional system at T_i may indicate that microaggregates of the organic and low-input systems have greater C availability and NO_3^- concentrations than the




Fig. 5. Canonical correspondence analysis (CCA) of measures of the 16S rRNA, *amoA*, and *nosZ* gene abundances (plotted as squares) in relation to soil environmental variables (plotted as vectors) for samples collected before the start of the cover crop growing season (T_0 : empty circles), after cover crop incorporation (T_1 : filled triangles), and during maize harvest (T_1 : X symbols). Canonical axes 1 (x axis) and 2 (y axis) explain 85.5% and 14.5% of the gene abundance–environmental variable variance in the data, respectively ($p = 0.006$).

conventional system. In contrast, the higher abundance of AOB in the microaggregates of the conventional system suggested that this microenvironment was characterized by NH_4^+ levels and conditions relatively more favorable to AOB than the conditions associated with microaggregates of the low-input and organic systems. Additionally, the consistent growth of nitrifiers across the season, particularly in the microaggregates of the conventional system, concurrent with a decrease in denitrifiers, suggested a steady increase in availability of O_2 over the season as the soil became drier.

4.3. Relationship between microbial community abundance and N processing

The combined significance of the first and second canonical axes in the first CCA of this study ($p = 0.006$) suggested that AOB and denitrifier community abundances correlated to potential gross N mineralization and nitrification rates and to NH_4^+ and NO_3^- concentrations; however, the first and second canonical axes were no longer significant when sampling event was included as a covariable in the CCA. Therefore, our data suggested that seasonality substantially influences the relationship between gene abundances and inorganic N concentrations and transformation rates, and that this relationship is a weak one. The lack of correlation between NO_3^- concentrations and *nosZ* gene abundance is in agreement with Kandeler et al. (2006), who found that the soil nitrate content could not explain the abundance of 16S rRNA, *narG*, *nirK*, *nirS*, and *nosZ* genes in the early stage of primary succession in a glacier foreland. Moreover, other studies have also found that gene abundance and microbial community composition cannot predict corresponding ecosystem processes. For example, Ma et al. (2008) found that, although different nitrifier and denitrifier communities were specific to landforms within cultivated versus uncultivated wetlands, these differences were not related to land-use or landform differences in N_2O emissions. Similarly, Dandie et al. (2008) found that differences in denitrifier community compositions and abundances in a potato (*Solanum tuberosum* L.) cropping system were not related to denitrification rates and N_2O emissions.

The current approaches used to study microbial communities with respect to soil processes have made substantial inroads in

our understanding of the linkages between the two. Nevertheless, to determine whether a relationship exists between ecosystem N processes and soil microbial population densities, future studies must ensure that approaches to measuring these relationships are not confounded by temporal or spatial biases. Moreover, the methods used to quantify nitrifiers and denitrifiers must strive to encompass all microorganisms responsible for nitrification and denitrification. Recently, Leininger et al. (2006) found a high abundance and wide distribution of archaeal ammonia monooxygenase (*amoA*) genes in the soil environment. To comprehensively characterize the nitrifying populations in soil, the abundances of autotrophic ammonia-oxidizing bacteria of the β - and γ -subgroups of proteobacteria should be considered along with the abundances of archaea.

4.4. Conclusions

This work sought to relate the abundances of AOB, denitrifiers, and total bacterial communities within different soil microenvironments to their activity along a gradient of long-term N management regimes. We found that, although crop management influenced AOB and denitrifier community abundance, changes in AOB and denitrifier community sizes did not relate to changes in NH_4^+ or NO_3^- concentrations and that potential gross N mineralization and nitrification rates were decoupled from nitrifier and denitrifier community abundances. Consequently, our study did not show a strong link between nitrifier and denitrifier abundances and N cycling processes. This study did find that microaggregates were characterized by larger nitrifier, denitrifier, and total bacterial communities that fluctuated more than communities in the cPOM and silt-and-clay fractions, particularly in low-input and organic systems; hence, the microaggregate appears to be a microenvironment that is conducive for nitrifiers and denitrifiers (e.g., higher C accumulation and NO_3^- content), and is, consequently, a potential hotspot for N_2O loss.

Acknowledgements

Many thanks to Marisa Alcorta and the entire staff at the Russell Ranch for their assistance in the field. We also thank Engil Isadora Pujol Pereira, Alice Yan, and Theresa Yim for their invaluable work in the laboratory. Thanks to Neil Willits and Dr. John Battles for assisting us with the statistical analyses in this study and to two anonymous reviewers for their comments to improve this manuscript. Funding for this project was provided by the Kearney Foundation of Soil Science, University of California, a grant from the Western Sustainable Agriculture Research and Education Program, and by grant number 5 P42 ES04699 from the National Institute of Environmental Health Sciences, NIH. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIEHS, NIH.

References

- Avrahami, S., Conrad, R., Braker, G., 2002. Effect of soil ammonium concentration on N_2O release and on the community structure of ammonia oxidizers and denitrifiers. *Applied and Environmental Microbiology* 68, 5685–5692.
- Barraclough, D., 1991. The use of mean pool abundances to interpret ^{15}N tracer experiments. I. Theory. *Plant and Soil* 131, 89–96.
- Beare, M.H., Parmelee, R.W., Hendrix, P.F., Cheng, W., Coleman, D.D., Crossley Jr., D.A., 1992. Microbial and faunal interactions and effects on litter nitrogen and decomposition in agroecosystems. *Ecological Monographs* 62, 569–591.
- Beauchamp, E.G., Trevors, J.T., Paul, J.W., 1989. Carbon sources for bacterial denitrification. *Advances in Soil Science* 10, 113–142.
- Birch, H.F., 1958. The effect of soil drying on humus decomposition and nitrogen availability. *Plant and Soil* 10, 9–31.
- Bossio, D.A., Scow, K.M., Gunapala, N., Graham, K.J., 1998. Determinants of soil microbial communities: effects of agricultural management, season, and soil type on phospholipid fatty acid profiles. *Microbial Ecology* 36, 1–12.

- Burford, J.R., Bremner, J.M., 1975. Relationships between the denitrification capacities of soils and total, water-soluble and readily decomposable soil organic matter. *Soil Biology & Biochemistry* 7, 389–394.
- Burger, M., Jackson, L.E., 2003. Microbial immobilization of ammonium and nitrate in relation to ammonification and nitrification rates in organic and conventional cropping systems. *Soil Biology & Biochemistry* 35, 29–36.
- Cavagnaro, T.R., Jackson, L.E., Scow, K.M., Hristova, K.R., 2007. Effects of arbuscular mycorrhizas on ammonia oxidizing bacteria in an organic farm soil. *Microbial Ecology* 54, 618–626.
- Dambreville, C., Hallet, S., Nguyen, C., Morvan, T., Germon, J.-C., Philippot, L., 2006. Structure and activity of the denitrifying community in a maize-cropped field fertilized with composted pig manure or ammonium nitrate. *FEMS Microbiology Ecology* 56, 119–131.
- Dandie, C., Burton, D.L., Zebbarth, B.J., Henderson, S.L., Trevors, J.T., Goyer, C., 2008. Changes in bacteria denitrifier community abundance over time in an agricultural field and their relationship with denitrification activity. *Applied and Environmental Microbiology* 74, 5997–6005.
- Doane, T.A., Horwath, W.R., 2003. Spectrophotometric determination of nitrate with a single reagent. *Analytical Letters* 36, 2713–2722.
- Drury, C.F., Yang, X.M., Reynolds, W.D., Tan, C.S., 2004. Influence of crop rotation and aggregate size on carbon dioxide production and denitrification. *Soil and Tillage Research* 79, 87–100.
- Ellert, B.H., Bettany, J.R., 1995. Calculation of organic matter and nutrients stored in soils under contrasting management regimes. *Canadian Journal of Soil Science* 75, 529–538.
- Elliott, E.T., 1986. Aggregate structure and carbon, nitrogen, and phosphorus in native and cultivated soils. *Soil Science Society of America Journal* 50, 627–633.
- Firestone, M.K., 1982. Biological denitrification. In: Stevenson, F.J. (Ed.), *Nitrogen in Agricultural Soils*. American Society of Agronomy, Madison, pp. 289–326.
- Forster, J.C., 1995. Soil nitrogen. In: Alef, K., Nannipieri, P. (Eds.), *Methods in Applied Soil Microbiology and Biochemistry*. Academic Press, San Diego, CA, pp. 79–87.
- Foster, S.S.D., Bridge, L.R., Geake, A.K., Lawrence, A.R., Parker, J.M., 1986. The Groundwater Nitrate Problem: A Summary of Research on the Impact of Agricultural Land-use Practices on Groundwater Quality Between 1976 and 1985. *Hydrogeol. Rep. Br. Geol. Surv. No. 86/2*. Br. Geol. Surv., Oxfordshire, England.
- Frey, S.D., Elliott, E.T., Paustian, K., 1999. Bacterial and fungal abundance and biomass in conventional and no-tillage agroecosystems along two climatic gradients. *Soil Biology & Biochemistry* 31, 573–585.
- Gale, W.J., Cambardella, C.A., Bailey, T.B., 2000. Root-derived carbon and the formation and stabilization of aggregates. *Soil Science Society of America Journal* 64, 201–207.
- Gunapala, N., Scow, K.M., 1998. Dynamics of soil microbial biomass and activity in conventional and organic farming systems. *Soil Biology & Biochemistry* 30, 805–816.
- Gupta, V.V.S.R., Germida, J.J., 1988. Distribution of microbial biomass and its activity in different soil aggregate size classes as affected by cultivation. *Soil Biology & Biochemistry* 20, 777–786.
- Hallin, S., Jones, C.M., Schloter, M., Philippot, L., 2009. Relationship between N-cycling communities and ecosystem functioning in a 50-year-old fertilization experiment. *ISME Journal* 3, 597–605.
- Hart, S.C., Nason, G.E., Myrold, D.D., Perry, D.A., 1994. Dynamics of gross nitrogen transformations in an old-growth forest: the carbon connection. *Ecology* 75, 880–891.
- Henry, S., Bru, D., Stres, B., Hallet, S., Philippot, L., 2006. Quantitative detection of the *nosZ* gene, encoding nitrous oxide reductase, and comparison of the abundances of 16S rRNA, *narG*, *nirK*, and *nosZ* genes in soils. *Applied and Environmental Microbiology* 72, 5181–5189.
- Holmes, A.J., Costello, A., Lidstrom, M.E., Murrell, J.C., 1995. Evidence that particulate methane monooxygenase and ammonia monooxygenase may be evolutionarily related. *FEMS Microbiology Letters* 132, 203–208.
- Jastrow, J.D., 1996. Soil aggregate formation and the accrual of particulate and mineral-associated organic matter. *Soil Biology & Biochemistry* 28, 656–676.
- Kandeler, E., Deiglmayr, K., Tschirko, D., Bru, D., Philippot, L., 2006. Abundance of *narG*, *nirS*, *nirK*, and *nosZ* genes of denitrifying bacteria during primary successions of a glacier foreland. *Applied and Environmental Microbiology* 72, 5957–5962.
- Kong, A.Y.Y., Six, J., Bryant, D.C., Denison, R.F., van Kessel, C., 2005. The relationship between carbon input, aggregation, and soil organic carbon stabilization in sustainable cropping systems. *Soil Science Society of America Journal* 69, 1078–1085.
- Kowalchuk, G.A., Stephen, J.R., 2001. Ammonia-oxidizing bacteria: a model for molecular microbial ecology. *Annual Review of Microbiology* 55, 485–529.
- Kramer, A.W., Doane, T.A., Horwath, W.R., van Kessel, C., 2002. Combining fertilizer and organic inputs to synchronize N supply in alternative cropping systems in California. *Agriculture, Ecosystems and Environment* 34, 43–50.
- Lee, J., Hopmans, J.W., Rolston, D.E., Baer, S.G., Six, J., 2009. Determining soil carbon stock changes: simple bulk density corrections fail. *Agriculture, Ecosystems, and Environment* 134, 251–256.
- Leininger, S., Urich, T., Schloter, M., Schwark, L., Qi, J., Nicol, G.W., Prosser, J.I., Schuster, S.C., Schleper, C., 2006. Archaea predominate among ammonia-oxidizing prokaryotes in soils. *Nature* 442, 806–809.
- Lensi, R., Clays-Josserand, A., Jocteur Monrozier, L., 1995. Denitrifiers and denitrifying activity in size fractions of a mollisol under permanent pasture and continuous cultivation. *Soil Biology & Biochemistry* 27, 61–69.
- Leps, J., Smilauer, P., 2003. *Multivariate Analysis of Ecological Data Using CANOCO*. University Press, Cambridge, UK.
- Luo, J., Tillman, R.W., Ball, P.R., 1999. Factors regulating denitrification in a soil under pasture. *Soil Biology & Biochemistry* 31, 913–927.
- Ma, W.K., Bedard-Haughn, A., Siciliano, S.D., Farrell, R.E., 2008. Relationship between nitrifier and denitrifier community composition and abundance in predicting nitrous oxide emissions from ephemeral wetland soils. *Soil Biology & Biochemistry* 40, 1114–1123.
- McCarty, G.W., Bremner, J.M., 1992. Availability of organic carbon for denitrification of nitrate in subsoils. *Biology and Fertility of Soils* 14, 219–222.
- Miller, M.N., Zebbarth, B.J., Dandie, C.E., Burton, D.L., Goyer, C., Trevors, J.T., 2009. Denitrifier community dynamics in soil aggregates under permanent grassland and arable cropping systems. *Soil Science Society of America Journal* 73, 1843–1851.
- Mosier, A.R., Duxbury, J.M., Freney, J.R., Heinemeyer, O., Minami, K., 1998. Assessing and mitigating N₂O emissions from agricultural soils. *Climatic Change* 40, 7–38.
- Mummey, D.L., Holben, W.E., Six, J., Stahl, P.D., 2006. Spatial stratification of soil microbial populations in diverse soils: Rubrobacteria and Gemmatimonada are abundant in water-stable microaggregate interiors while Acidobacteria are primarily associated with macroaggregates. *Microbial Ecology* 51, 404–411.
- Okano, Y., Hristova, K.R., Leutenegger, C.M., Jackson, L.E., Denison, R.F., Gebreyesus, B., Lebauer, D., Scow, K.M., 2004. Application of real-time PCR to study effects of ammonium on population size of ammonia-oxidizing bacteria in soil. *Applied and Environmental Microbiology* 70, 1008–1016.
- Parkin, T.B., 1987. Soil microsites as a source of denitrification variability. *Soil Science Society of America Journal* 51, 1194–1199.
- Paul, J.W., Beauchamp, E.G., 1989. Effect of carbon constituents in manure on denitrification in soil. *Canadian Journal of Soil Science* 69, 49–61.
- Philippot, L., Renault, P., Sierra, J., Henault, C., Clays-Josserand, A., Chenu, C., Chaussod, R., Lensi, R., 1996. Dissimilatory nitrite-reductase provides a competitive advantage to *Pseudomonas* spp. RTC01 to colonize the centre of soil aggregates. *FEMS Microbiology Ecology* 21, 175–185.
- Ranjard, L., Richaume, A.S., 2001. Quantitative and qualitative microscale distribution of bacteria in soil. *Research in Microbiology* 152, 707–716.
- Ranjard, L., Poly, F., Combrisson, J., Richaume, A., Gourbière, F., Thioulouse, J., Nazaret, S., 2000. Heterogeneous cell density and genetic structure of bacterial pools associated with various soil microenvironments as determined by enumeration and DNA fingerprinting approach (RISA). *Microbial Ecology* 39, 263–272.
- Recous, S., Mary, B., Faurie, G., 1990. Microbial immobilization of ammonium and nitrate in cultivated soils. *Soil Biology & Biochemistry* 22, 913–922.
- Robertson, G.P., 1997. Nitrogen use efficiency in row-crop agriculture: crop nitrogen use and soil nitrogen loss. In: Jackson, L.E. (Ed.), *Ecology in Agriculture*. Academic Press, San Diego, pp. 347–363.
- Rochette, P., Bochove, E.V., Prévost, D., Angers, D.A., Côté, D., Bertrand, N., 2000. Soil carbon and nitrogen dynamics following application of pig slurry for the 19th consecutive years: II. Nitrous oxide fluxes and mineral nitrogen. *Soil Science Society of America Journal* 64, 1396–1403.
- Rothauwe, J.-H., Witzel, K.-P., Liesack, W., 1997. The ammonia monooxygenase structural gene *amoA* as a functional marker: molecular fine-scale analysis of natural ammonia-oxidizing populations. *Applied and Environmental Microbiology* 63, 4704–4712.
- Sakala, W.D., Cadisch, G., Giller, K.E., 2000. Interactions between residues of maize and pigeonpea and mineral N fertilizers during decomposition and N mineralization. *Soil Biology & Biochemistry* 32, 679–688.
- SAS Institute, 2002. *Statistical Analysis System, Version 9.2*. SAS Institute, Cary, NC.
- Saxton, A.M., 1998. A macro for converting mean separation output to letter groupings in Proc Mixed. In: *Proceedings of 23rd SAS Users Group Intl. SAS Institute, Cary, NC*, pp. 1243–1246.
- Scow, K.M., 1997. Soil microbial communities and carbon flow in agroecosystems. In: Jackson, L.E. (Ed.), *Ecology in Agriculture*. Academic Press, San Diego, pp. 367–413.
- Seech, A.G., Beauchamp, E., 1988. Denitrification in soil aggregates of different sizes. *Soil Science Society of America Journal* 52, 1616–1621.
- Sexstone, A.J., Revsbech, N.P., Parkin, T.B., Tiedje, J.M., 1985. Direct measurement of oxygen profiles and denitrification rates in soil aggregates. *Soil Science Society of America Journal* 49, 645–651.
- Sexstone, A.J., Parkin, T.B., Tiedje, J.M., 1988. Denitrification response to soil wetting in aggregate and unaggregated soil. *Soil Biology & Biochemistry* 20, 767–769.
- Shi, W., Norton, J.M., 2000. Microbial control of nitrate concentrations in an agricultural soil treated with dairy waste compost or ammonium fertilizer. *Soil Biology & Biochemistry* 32, 1453–1457.
- Six, J., Elliott, E.T., Paustian, K., Doran, J., 1998. Aggregation and soil organic matter accumulation in cultivated and native grassland soils. *Soil Science Society of America Journal* 62, 1367–1377.
- Six, J., Elliott, E.T., Paustian, K., 2000. Soil macroaggregate turnover and microaggregate formation: a mechanism for C sequestration under no-tillage agriculture. *Soil Biology & Biochemistry* 32, 2099–2103.
- Stark, J.M., Hart, S.C., 1996. Diffusion technique for preparing salt solutions, Kjeldahl digests, and persulfate digests for nitrogen-15 analysis. *Soil Science Society of America Journal* 60, 1846–1855.
- Suzuki, M.T., Taylor, L.T., Delong, E.F., 2001. Phylogenetic analysis of ribosomal RNA operons from uncultivated coastal marine bacterioplankton. *Environmental Microbiology* 3, 323–331.
- Tiedje, J.M., 1988. Ecology of denitrification and dissimilatory nitrate reduction to ammonium. In: Zehnder, A.J.B. (Ed.), *Biology of Anaerobic Microorganisms*. John Wiley, New York, pp. 179–244.
- Weier, K.L., Doran, J.W., Power, J.F., Walters, D.T., 1993. Denitrification and the dinitrogen/nitrous oxide ratio as affected by soil water, available carbon, and nitrate. *Soil Science Society of America Journal* 57, 66–72.