



## EDITION #8: Fresh Tomato & Bell Pepper

Volume 2, Issue 2

February 2006

### IN THIS ISSUE

**New Farm Advisor - San Joaquin County**  
**Greenhouse Tomatoes Emerge as a Major Market Force**

**Statewide Fresh Tomato Variety Results**  
**Bell Pepper Variety Results**

**Water Requirements of Bell Peppers**

**Pepper Virus Diseases: A Review**

**Tomato Spotted Wilt Virus**

**Pepper Stip**

**Psyllids in Peppers & Tomatoes**

**Stink Bug Field Research**

Conservation Biological Control

Cultural Control

Chemical Control

**Nematode Resistant Varieties NOT always Resistant**

**Preemergence Herbicides in Transplanted Bell Peppers**

**Postharvest Handling of Grape Tomatoes**

### SOURCES OF INFORMATION

**Farm Advisors – Vegetable Crops**  
**San Joaquin Valley**

#### FRESNO

Shannon Mueller (559) 456-7261

Kurt Hembree (559) 456-7556

#### KERN

Joe Nuñez (661) 868-6222

#### MERCED & MADERA

Scott Stoddard (209) 385-7403

#### SAN JOAQUIN

Brenna Aegerter (209) 468-2085

#### STANISLAUS

Jan Mickler (209) 525-6800

#### TULARE & KINGS

Michelle Le Strange

(559) 685-3309, Ext. 220

#### Newsletter Editor

Michelle Le Strange

4437-B S. Laspina Street

Tulare, CA 93274

E-mail: mlestrange@ucdavis.edu

### Meet the New Farm Advisor...

I would like to introduce myself- I am the new vegetable crops Farm Advisor for San Joaquin County. I have recently relocated to the San Joaquin Valley from that “other” valley to the north and I am very excited to be here and working with the vegetable crop industry!



At UC Davis, I worked in the Department of Plant Pathology with Mike Davis and Tom Gordon. I conducted field and laboratory research and participated in the diagnosis of problems in field and vegetable crops from samples submitted from across the state. Among the types of problems we saw were viral, bacterial, and fungal diseases, insect injury, nutrient problems and other abiotic disorders. Although I am a plant pathologist by training, I am anticipating the challenge of tackling whatever crop production problems are faced in this area.

This season I will be participating in the statewide variety evaluation projects for both processing and fresh market tomatoes and I’ll be launching some pest management projects, including the evaluation of a powdery mildew model that might help growers predict disease in the San Joaquin Valley. And there will be another bell pepper variety trial in San Joaquin County this year. I look forward to meeting you and working with you!

--- Brenna Aegerter

### Announcement

## 38th California Nematology Workshop

**Tuesday, March 28, 2003, 8 AM - 4:30 PM**

**University of California Extension Center**

**1200 University Ave, Riverside, CA**

This annual workshop offers pest management professionals and growers the latest information on problems caused by plant-parasitic nematodes and on their potential solutions. Target audience for this program includes pest control advisors and operators, growers, pesticide and biocontrol industry representatives, landscapers, municipal and state employees, parks and recreation personnel, educators and consultants. A superb lineup of speakers and workshop presenters will share their expertise concerning nematode-related issues. Posters will inform about the latest Nematology research activities at the University of California, CDFA, USDA and industry. Breakout session will give the audience an opportunity to sharpen their skill in nematode and identification, disease diagnostics, and sampling procedures. For information and registration: go to [www.nematology.ucr.edu](http://www.nematology.ucr.edu) or contact antoon.ploeg@ucr.edu, 951-827-3192.

# North American Greenhouse Tomatoes Emerge as a Major Market Force

*Linda Calvin, Economic Research Service and Roberta Cook, UCCE Marketing Economist, UC Davis*

The rapidly growing greenhouse tomato industry has become an important part of the North American fresh tomato industry. Greenhouse tomatoes now represent an estimated 17 percent of U.S. fresh tomato supply. Even though greenhouse tomatoes still constitute a minority share of the U.S. fresh tomato market, their influence is concentrated and growing in retail channels, which represent about half of U.S. tomato consumption. *Around 37% of all fresh tomatoes sold in U.S. retail stores are now greenhouse, compared with negligible amounts in the early 1990s.*

Greenhouse tomatoes are just one more development in a trend toward more differentiated fresh tomato offerings, including more variety in fieldgrown tomatoes. New types of tomatoes, improved varieties and handling, and positive health benefits associated with eating tomatoes have all contributed to a 30% rise in U.S. consumption of fresh tomatoes since 1985, with estimated 2003 annual per capita consumption levels around 19.4 pounds.

***Growth in the greenhouse industry has challenged growers of fresh field tomatoes.*** With rising consumption of all tomatoes, field tomato sales in the U.S. retail market increased through 2001, in part due to new fresh field products, such as grape tomatoes. But in 2002, the combined retail sales volume of all field tomato types began to slip. Field tomatoes still dominate the growing foodservice market (restaurants, schools, hospitals, etc.) where greenhouse tomatoes are scarce. *Foodservice sales are increasingly essential to the health of the field tomato industry.*

While greenhouse tomatoes have higher per unit costs of production and generally higher retail prices than field tomatoes, several other characteristics have contributed to the growth in this sector. Since they are protected from weather and other conditions affecting open field production, greenhouse tomatoes generally have a much more uniform appearance than field

tomatoes. They are also less prone to swings in production volumes. These factors lead to greater consistency in quality, volumes, and pricing—issues of particular concern to the retail and foodservice industries.

The United States, Canada, and Mexico have all developed major greenhouse industries. The U.S. is the largest North American market for greenhouse tomatoes, and U.S. imports from Canada and Mexico are larger than domestic production. In recent years, the growth in U.S. imports has exceeded the growth in U.S. production. In 2003, Canada accounted for an estimated 46% of U.S. imports of greenhouse tomatoes. Mexico's share was 45%. *Mexico is the primary foreign winter supplier to the U.S. market and Canada is the primary summer foreign supplier.* As the greenhouse tomato industry has transitioned from niche to mainstream status, it has become part of a more integrated North American market, following the pattern established by the field tomato industry.

***The greenhouse industry is facing growing pains.*** With rapid growth in Canada and the United States during the 1990s, greenhouse tomato prices declined, causing financial problems for some growers. More recently, as the industry has expanded in Mexico, heterogeneity in production methods has increased. Growers in the United States and Canada, and some Mexican growers, have high-technology and high-cost greenhouses. Many of these growers view the growth of lower technology greenhouses and shade houses in Mexico with some alarm. Higher expected year-round production volumes in Mexico portend greater competition in all seasons, and continued downward pressure on price.

## **Seasonality Drives Market**

Much of the U.S. greenhouse tomato industry began in the northeast in the early 1990s, with production in the same months as Canadian producers. Eventually, several producers moved west and south, lured by the prospect of producing tomatoes year-round and capturing a slice of the high-priced winter market. *The four largest*

*greenhouse tomato firms in the U.S. are located in Arizona, Texas, Colorado and coastal southern California, and represent 67% of domestic production.*

Expanding winter production in Mexico will likely reduce greenhouse tomato prices and increase competitive pressure on year-round U.S. growers. Mexico's greenhouse tomato industry is the fastest growing in North America and the most varied. In Mexico, large field tomato grower-exporters in Sinaloa on the northwest coast and Baja California peninsula are experimenting with protected culture, either shade houses or greenhouses, near their field operations. *In contrast, U.S. field tomato growers usually have no connections to the greenhouse industry.*

Several clusters of greenhouses are also emerging in temperate, higher altitude areas in central and north central Mexico, and in Imuris in northern Sonora, near the U.S. border. *As greenhouse production in these areas expands, Mexico will become more of a competitive force in all seasons.*

## **Greenhouse Tomato Prices Fall**

Despite rising demand for greenhouse tomatoes, the industry is facing downward price pressures, as demand growth has sometimes been outpaced by expanding supply. Production of the leading greenhouse tomato products—beefsteak and cluster—has now grown to the point where they are becoming mainstream commodities. As the industry matures, greenhouse tomato growers strive for continual product innovation as a strategy for adding value, stimulating consumer interest, and maintaining margins and profitability. The expanding product line currently consists of smaller cluster tomatoes (cocktail tomatoes, including Campari), roma and mini roma cluster tomatoes, heirloom, and different-colored tomatoes. *Greenhouse tomato producers tend to be closer to the pulse of consumers because they market a retail- and consumer-ready product. Also, they increasingly market directly to retailers, and not through intermediaries, such as repackers and wholesalers, as most field tomato shippers do.*

## **Impacts on Field Tomatoes**

Competition from greenhouse tomatoes has brought major changes in the quantity and composition of field tomato sales. While

## Canada leads North American greenhouse tomato production in 2003

Item	United States	Canada	Mexico	North America
Greenhouse tomato production (1,000 metric tons)	160	220	148	528
Greenhouse tomato area (hectares)	330	446	950	1,726
Average greenhouse tomato yield (metric tons/hectare)	484	494	156	378
Fresh field tomato production, excluding processing (1,000 metric tons)	1,594	27	1,804	3,425
Average fresh field tomato yield (metric tons/hectare)	32	15	28	25
Greenhouse share of total fresh production, by country (percent)	9	89	8	13
Estimated greenhouse exports to U.S. (1,000 metric tons) <sup>1</sup>	NA	130	126	256

<sup>1</sup>Official imports of greenhouse tomatoes are thought to be underreported for Mexico due to tariff code misclassification; 58,357 metric tons of greenhouse tomato imports from Mexico were reported by the U.S. Department of Commerce in 2003. The figure shown here includes estimated additional miscoded imports, based on information from industry sources obtained by Cook and Calvin. This figure may include some production from shade houses.

NA=Not applicable.

Sources: Statistics Canada, Ontario Greenhouse Vegetable Producers' Marketing Board, British Columbia Vegetable Marketing Commission, U.S. Department of Commerce, interviews by Cook and Calvin, USDA's National Agricultural Statistics Service, USDA's Foreign Agricultural Service.

total retail quantity sold of all fresh tomatoes increased from 1999 to 2003, the volume of field tomatoes declined after 2001, with the share falling from 69 to 63%. *Over the same years, the share of all round tomatoes (mature green and vine ripe) declined from 43 to 31%. The roma share fell from 23 to 19%, but the grape and cherry category grew from 3 to 13%. Most grape and cherry tomatoes are field grown, mitigating the impact of greenhouse tomatoes on the field-grown category. Within the declining round category, the share of mature green tomatoes fell from 78 to 39%, with vine ripe tomatoes benefiting.*

While mature green tomatoes are being forced out of the retail market by competition from both greenhouse and other field tomato types, they still dominate the expanding foodservice market, which represents about half of U.S. tomato consumption. With declining retail sales, the mature green industry is increasingly dependent on the foodservice market, where greenhouse tomatoes have not yet made significant inroads. However, this could change. *Some greenhouse firms have begun to experiment with developing acceptable products for foodservice use.* If foodservice demand falters, mature green tomato growers would need to consider other alternatives, with serious industry structural adjustments likely. Growers could continue to attempt to reposition field tomatoes through new varieties,

products, and packaging with more commercial appeal. Alternatively, the industry could diversify into the greenhouse industry, either through alliances with existing producers or through direct investment. However, greenhouse tomato production is very capital- and technology-intensive, creating barriers to entry. In addition, the rapid greenhouse expansion in the United States was accompanied by mixed profitability results; thus, most field tomato growers did not consider the greenhouse industry an attractive alternative. But recent profitability in the California field industry caused by weather-induced high prices may provide the financial where-withal for some field growers to explore greenhouse production. If they were to invest, they would be new entrants in a maturing industry.

### Greenhouse and Field Tomato Market Interactions Increase

In the early days of the evolution of greenhouse tomatoes, the greenhouse and field tomato sectors operated on a relatively independent basis. Now that they are a major market force, greenhouse tomatoes are increasingly influenced by supply and demand trends in the fresh field tomato industry, and vice versa. In fall 2004, a weather-induced period of short supplies of fresh field tomatoes enabled greenhouse producers to benefit from a brief period of extraordinarily high prices as buyers substituted greenhouse for field tomatoes, where possible. In contrast,

earlier in summer 2004, a record-high supply of greenhouse tomatoes caused greenhouse prices to decline, making them even more attractive to retail buyers, and placing a damper on demand for fresh field tomatoes. With greater supply has come an increased willingness on the part of consumers, retailers, and foodservice users to experiment with tomato types.

### Mexico Will Shape the Future

Notwithstanding brief periods of abnormally high prices, average grower prices for greenhouse tomatoes have been trending downward. If this trend continues, some parts of the North American greenhouse tomato industry may become less viable. Growers will continue to seek the lowest cost production regions and form marketing alliances to build year round supply. Greater competition means that new entrants have less room for error; the learning curve is shorter than in the 1990s, when the industry was in its infancy and average prices were higher. *The greatest source of uncertainty for the future of the North American greenhouse tomato industry will be the changing structure of the Mexican industry, which is still seeking out the best locations, technology packages, and management practices.* U.S. and Canadian growers will be following developments in Mexico closely when making their future investment and marketing decisions.

### This article is drawn from . . .

Greenhouse Tomatoes Change the Dynamics of the North American Fresh Tomato Industry, by Roberta Cook & Linda Calvin, ERR-2, USDA/ERS, Apr. 2005, available at: [www.ers.usda.gov/publications/err2/](http://www.ers.usda.gov/publications/err2/)

# Statewide Fresh Market Tomato Variety Field Evaluations for 2005

Scott Stoddard, Michelle Le Strange, Bob Mullen (Emeritus) and Jan Mickler, UCCE  
Farm Advisors, Merced & Madera, Tulare & Kings, San Joaquin, and Stanislaus Counties

## Introduction

UCCE conducts fresh market tomato variety trials in three areas in the San Joaquin Valley to evaluate the performance of new varieties and breeding lines from commercial plant breeders for the mature green market. These variety trials evaluate and compare fruit quality characteristics and yield in commercial production fields with different types of soil, management, and growing conditions. This market includes both round and “roma” type tomatoes.

## Procedure

Trials are laid out as randomized complete block designs with 4 replications (observation lines are not replicated but are planted adjacent to the replicated plots). Plots are transplanted and managed concurrently as the commercial field in which they are located. Harvest is done by hand at the same time as the rest of the field, picking from a 10 foot section from the center of the plot. At harvest, fruit are sorted by culls, color, and size. Small fruit (2 – 2.25”) are picked but are not included in the total market yield.

## Results

**Replicated Lines:** Results for marketable yield and fruit size for Fresno, Merced, and San Joaquin Counties are shown in Figure 1. Shady Lady and Quali T-21 are the standards to which the other varieties are compared. In Fresno, BHN 580 was the clear standout with regard to yield, with a mean yield over 2400 boxes/A. This was largely a result of an over-production of jumbo sized fruit.

Merced also had a clear winner with AT-37, at over 2500 boxes per acre. Overall, the production of XL fruit was much lower in Merced compared to the other locations.


There was no variety in San Joaquin County that was so markedly higher yielding than the rest. AT-37, Q-21, Catalyst, and RFT 500-311 all yielded similar to each other at around 2000 boxes per acre.

The LSD’s for Fresno, Merced, and San Joaquin Counties were 211, 424, and 360 boxes per acre, respectively. Additional information about this trial can be found in the full report posted on the Merced County website at <http://cemerced.ucdavis.edu>.

**Observed Lines:** The combined market yields for each county are shown in Figure 2. Because there is no replication in the observed lines, statistical analysis could be performed only on the combined data set. SRT 6784 did particularly well in Fresno, while BHN 525 and PX 2942 yielded well in Merced and San Joaquin locations. Combining locations, no significant differences among varieties were found for yield or size, mainly because of the large amount of variability in the data

**Romas:** A replicated roma trial was conducted in San Joaquin County. At that location, Miroma performed better than the other lines. Contact Jan Mickler or Bob Mullen for more information. Six roma varieties were observed in Fresno (visit [http://cetulare.ucdavis.edu/Vegetable\\_Crops/](http://cetulare.ucdavis.edu/Vegetable_Crops/))

**Figure 1.** Yield by size class for all three locations. Stacked bars show medium fruit on top, large in middle, and extra-large on bottom. Error bars are standard error of the mean for each variety. Total height of the bar is the total market yield.



**Figure 2.** Total market yield results. Error bars represent one standard error of the mean. Variety yields are not significantly different.

## 2005 Bell Pepper Variety Evaluation Trial

*Benny Fouche and Bob Mullen (Emeritus), Farm Advisors, UCCE San Joaquin County*

Now that there are a number of new pepper cultivars available to producers, information on yield and fruit quality, as well as disease resistance or tolerance is desirable for the local industry. This year's trial at Biglieri Farms on the Borden Ranch near Dry Creek, east of Galt, California was transplanted on June 10<sup>th</sup>. The soil type at the trial site was a Wyman silt loam and the trial field was alternate-row furrow-irrigated throughout the season. The resulting crop stand was excellent with vigorous early plant growth. A very hot July and August caused some plant stress, loss of fruit set and a subsequent delay in fruit maturity. Hand harvest of the trial was on September 10th.

The trial included fifteen replicated varieties arranged in a randomized complete block design. In addition to marketable red and green yield figures, data on crop maturity and fruit size were taken (Table 1). Best quality fruit, including blocky shape and good fruit color and size was led by Double Up, Encore, Mercado, Red Bell, RPP 9650, Affinity, RPP 16900, and Baron. Fruit size for most of the lines evaluated was predominately jumbo and extra-large. Other than some fruit sunburn and blossom end rot and some cat-faced fruit, there were no other fruit defect problems. There was virtually no worm damage in the trial and none of the fruit had Pepper Spot (STIP).

*The same varieties were evaluated in Morgan Hill by Aziz Baameur. His report is available at UCCE Santa Clara Co. website (<http://cesantaclara.ucdavis.edu/>).*

**Table 1. Yield, maturity, and fruit size percent for 15 bell pepper varieties – Galt, CA 2005**

Variety	Marketable yield/acre (red + green)		Crop maturity at harvest (%)			Fruit size (%) <sup>2</sup>					Total yield/acre
	Tons <sup>1</sup>	Boxes	Red	Green	Culls	Jumbo	Extra-large	Large	Med	Small	Tons
Red Bell	16.0	1,281	1.5	58.5	40.0	34.2	25.9	23.6	9.9	6.4	26.8
Double Up	15.9	1,272	9.8	56.0	34.2	30.3	20.1	28.5	10.5	10.6	24.3
Encore	146	1,167	9.1	55.2	35.7	55.8	18.8	10.6	7.4	7.4	22.6
RPP16900	13.8	1,103	14.4	49.1	36.5	3.3	19.6	28.9	22.7	25.5	21.7
Mercado	13.5	1,080	4.3	57.2	38.5	66.3	17.3	9.7	2.9	3.8	22.1
RPP9650	13.4	1,074	8.4	47.1	44.5	64.5	21.8	10.9	1.1	1.7	23.7
RPP9661	13.4	1,071	5.3	57.3	37.4	53.1	14.4	8.9	9.7	13.9	21.3
Baron	13.4	1,068	12.1	48.7	39.2	4.1	27.7	32.7	21.7	13.8	22.1
Wizard	13.1	1,049	4.7	54.7	40.6	47.0	23.3	10.3	6.1	13.3	21.9
Affinity	12.7	1,016	11.3	46.0	42.7	51.0	21.9	14.8	3.6	8.7	22.9
Stiletto	12.1	967	3.4	50.2	46.4	9.5	31.7	27.2	19.1	12.5	22.1
Crusader	10.7	857	2.7	52.3	45.0	34.4	25.8	12.8	13.6	13.4	19.2
Excel	10.1	804	3.0	47.7	49.3	24.9	19.9	32.4	7.0	15.8	19.8
Jupiter	9.7	778	1.6	42.1	56.3	51.1	23.3	14.9	3.7	7.0	22.1
Escarlata	7.4	590	6.4	29.3	64.3	20.0	10.2	20.0	24.5	25.3	21.4
<b>Average</b>	<b>12.7</b>	<b>1,012</b>									
LSD <sup>3</sup>	4.2	336									
C.V.	23.3%	23.3%									

<sup>1</sup>Values represent the average of four replications

<sup>2</sup>Pepper fruit sizing data: Jumbo: >8.5 oz; Extra-large: 7 – 8.5 oz; Large: 6 – 7 oz; Medium: 5.3 – 6 oz; Small: <5.3 oz

<sup>3</sup>Least significant difference at 5% significance level

**Many thanks to the cooperators and to the participating seed companies for their support of this work!**

# Water Requirements of Irrigated Bell Peppers

Tom Trout and James Ayars, USDA/ARS, Water Management Research Laboratory, Parlier

## Introduction

There has been a shift in cropping from long season high water requirement crops (tomato, cotton) to short season vegetable crops (lettuce, pepper, broccoli, onion) on the west side of the San Joaquin Valley (SJV). There has also been a shift in irrigation systems from surface irrigation to pressurized systems i.e. sprinkler and microirrigation. There is very little information describing the crop water requirements for vegetable crops grown in this region using sprinkler and microirrigation. This is a report of the results of a field study that determined the crop water requirements for a bell pepper crop grown on the westside of the SJV using drip and furrow irrigation.

## Materials and Methods

Three different irrigation systems were installed at the West Side Research and Extension Center to evaluate and compare irrigation methods commonly used to grow vegetable crops on the west side. These include:

- 1) a furrow irrigation system,
- 2) a surface drip irrigation system, and
- 3) a subsurface drip irrigation system with drip laterals installed 12 inches deep.

Water was applied with each system at four different irrigation levels in order to determine the application amount needed to obtain maximum yield. Amounts of applied water were equal to 50, 75, 100 or 125% of the crop evapotranspiration rate determined from water use in a well watered crop lysimeter. (*Lysimeter is a device for measuring the percolation of water through soils.*)

The 12 irrigation treatments were arranged in a split-plot experimental design with four replicate plots per treatment. Each plot was 300 feet long and consisted of four crop beds spaced 40 inches from center to center; outside beds served as borders between treatments.

An irrigation control system applied all drip irrigations automatically in response to crop lysimeter water use. The lysimeter (which has drip tubing installed 12 inches deep) and all drip irrigation treatments in the field were watered after 0.08 inch of crop evapotranspiration was measured by the lysimeter. This resulted in several applications each day to match peak water use. Furrow irrigated plots were watered weekly based on the accumulated water use over the previous 7 days.

Bell peppers (var Baron) were planted on April 25, 2005 as transplants with a planting density of 17,000 plants/ac (10-inch in row spacing by 40-inch row spacing). Harvest was in July and early August. Plants were grown following

normal cultural practices, which included pre-plant and irrigation applied nutrients. Sprinkler irrigation was used to establish seedlings.

Water applied to each treatment was recorded automatically using electronic flow meters installed in the irrigation manifold. Crop evapotranspiration was measured with a lysimeter and with a Bowen Ratio system installed in the pepper field. A second Bowen Ratio system was installed in the grass field next to the pepper field. Crop ET measured by the Bowen Ratio system in the peppers was divided by the grass ET measured by the Bowen Ratio system in the grass field to calculate daily crop coefficient K<sub>c</sub> values.

Peppers were harvested 3 times from a 30 foot section of the center 2 rows of each treatment. The peppers were sorted into green and red market peppers and culls.

## Results and Discussion

The applied water for each of the treatments is summarized in **Table 1**. The ET measured by the crop lysimeter was 20-inches and the data show that the target ET levels were met for the drip systems and approximately 5% higher in the furrow systems.

The daily evapotranspiration for the crop and the grass reference are plotted in **Figure 1**. The data show that there was approximately 0.3-0.4 inch of water lost per day in the grass and pepper crop during July and August with the pepper crop ET being higher than the grass. These data were used to calculate the bell pepper crop coefficient shown in **Figure 2**. The K<sub>c</sub> in July and August was between 1 and 1.2 for the pepper crop and was an average across all the treatments.

The yield results for the 2005 experiment are summarized in **Table 2**. The data show that the two drip treatments had the similar yields at three of the irrigation levels. Furrow irrigation yields were less than either of the drip treatments at both 100% and 125% irrigation levels. At the 50% level there was no significant difference across the system type. Comparing the mean values of the water treatments, the data show that the yields for the 50% treatments are less and the mean yield for the 125% water treatment was statistically greater than the intermediate treatments.

The water use efficiency data (**Table 3**) were generally lowest for the furrow treatments and highest for the subsurface drip treatments with the exception being the 125% treatment.

**Table 1. Applied irrigation water (inches) on bell pepper irrigation trial at WSREC in 2005.**

Irrigation levels				
Irrigation Methods	50% ET	75% ET	100% ET	125% ET
Furrow	10.7	15.7	20.8	26.0
Surface Drip	10.0	14.9	19.9	24.7
Sub Surface Drip	9.9	14.9	19.7	24.7


**Table 2. Pepper market yield (Tons/acre) at WSREC experimental sited during 2005 growing season.**

Irrigation levels					
Irrigation Methods	50% ET	75% ET	100% ET	125% ET	Mean
Furrow	9.5 ef	11.7df	10.1 def	11.9 d	10.8 b
Surface Drip	8.5 f	11.9 d	15.4 c	19.9 a	13.9 ab
Sub Surface Drip	10.1 def	15.5 c	17.5 bc	18.0 ab	15.3 a
Mean	9.4 c	13.1 b	14.3 b	16.6 a	13.6


LSD(0.05) for irrigation methods = 3.4 T/A LSD (0.05) for irrigation levels = 2.2 T/A LSD (0.05) for interaction (M x L) = 2.3 T/A

**Table 3. Water use efficiency in Tons/Acre/inch of applied water**

Irrigation levels				
Irrigation Methods	50% ET	75% ET	100% ET	125% ET
Furrow	0.89	0.75	0.49	0.46
Surface Drip	0.85	0.80	0.77	0.81
Sub Surface Drip	0.95	1.04	0.89	0.73



**Figure 1. Daily evapotranspiration of grass ( $ET_o$ ) and pepper ( $ET_c$ ) at the West Side Research and Extension Center in July 1 to August 17, 2005 measured by the Bowen Ratio technique.**



**Figure 2. Pepper crop coefficient calculated using Bowen Ratio data from July 1 to August 17, 2005.**

# Pepper Virus Diseases: A Review

*Steven Koike, Richard Smith, and Aziz Baameur, UCCE Farm Advisors, Monterey and Santa Clara Counties*

**Introduction and Significance:** Pepper is susceptible to a large number of virus pathogens. Worldwide, over 70 such agents have been documented to some degree, and other virus-like diseases have yet to be fully characterized. Some of these virus diseases are economically important throughout the world, while others are significant in only specific, limited areas. For pepper growers in California, perhaps ten viruses are of regular or periodic concern (see Table 1). The particular set of viruses that might be of economic importance can change over time. In the early to mid-1990s, cucumber mosaic, pepper mottle, and tobacco etch viruses were perhaps the most commonly encountered pepper virus problems. Now in the 2000s, that situation may have changed significantly.

**Symptoms and Diagnostic Features:** For any particular pepper virus, the incidence and expression of disease symptoms will vary greatly depending on the strain and virulence of the virus, pepper species and cultivar, age of pepper plant when infected, means of inoculation (e.g. whether the virus entered the plant by mechanical abrasion or was initially in the pepper seed), vector type and strain, population of the vector, and environmental conditions. With few exceptions, symptoms caused by different viruses often resemble each other, thereby making field diagnosis difficult and ill advised. Virus disease diagnosis is further complicated when more than one viral agent infects the pepper plant. Clinical tests are required to positively identify viral agents in plants.

**Disease Cycle:** Disease cycles of the various pepper viruses are similar. Primary inoculum mostly comes from infected weed hosts, volunteer Solanaceous plants, or existing pepper plantings. However, for the few pepper seedborne viruses, the germinating seedling or infected transplant will be the inoculum source. Insect or nematode vectors then move the viruses from infected plants to healthy plants. Vector movement generally dictates pathogen distribution for most of these diseases. Some of these pepper viruses can be readily transmitted by mechanical means such as handling by workers and pruning tools.

**Control:** Virus diseases tend to be difficult to control. The use of resistant cultivars would be the best option for growers; however, for the California pepper industry we do not yet have suitably resistant peppers that have the necessary horticultural features. Remove reservoirs of virus pathogens by controlling weeds, volunteer peppers, and other Solanaceous volunteers near fields. Plow under old pepper fields soon after harvest is completed. Carefully inspect and remove any transplants that show virus symptoms and vector activity. Crop rotation is a good practice in general, though such rotation usually does not assist in virus disease management unless the vectors are soilborne nematodes. For seedborne pepper viruses, use seed that has been tested and found to not have detectable levels of the pathogen, or that has a pathogen level below significant thresholds. Controlling the vector does not prevent virus

infections from taking place. However, management of insect vectors is important and should be attempted by applying insecticides and other insect control materials, planting crops on reflective mulches to repel vectors, or planting crops under netting, fabric, or plastic tunnels to exclude vectors. Soilborne nematode vectors can be managed by using soil-applied fumigants, rotating crops, and cultivating regularly to reduce the growth of host weeds and volunteers.

**Pepper Virus Survey:** Because of severe crop losses in 2004 due to pepper viruses, we conducted field surveys in coastal California pepper growing regions to identify pepper virus incidence. Symptomatic pepper plants were randomly collected from fields in the Gilroy, Hollister, and King City areas. Various pepper types (Anaheim, ancho, bell, jalapeno) were collected and tested using serological assays. The survey was conducted in 2004 and 2005. Results were similar for both years. The great majority of samples were infected by either cucumber mosaic virus (CMV) or tomato spotted wilt virus (TSWV). Many samples were co-infected with CMV and TSWV. Other pepper viruses were found but were very low in incidence and clearly were not important factors. Such incidental finds included the following: alfalfa mosaic virus, potato virus Y, tobacco etch virus, tobacco mosaic virus.

**Summary:** Pepper viruses will continue to be a long-term concern for pepper growers. In some seasons, such as 2004, virus diseases will cause significant crop losses. In other seasons the viruses will be less important and disease incidence will be low. Continued research efforts will be warranted to further understand and define the causes of such virus problems and to develop suitable resistant cultivars. With the worldwide movement of plant materials, it will be certain that sometime in the future new virus pathogens will make their way into California. Growers, field personnel, and extension researchers should therefore monitor pepper virus situations and be aware of such new developments. Contact the Farm Advisor in your region, if you see outbreaks of virus symptoms on peppers.

**Table 1. Summary of some pepper viruses found in CA**

Pathogen	Acronym	Virus group	Primary transmission
alfalfa mosaic virus	AMV	alfamovirus	Aphid
cucumber mosaic virus	CMV	cucumovirus	Aphid
pepper mottle virus	PepMoV	potyvirus	Aphid
potato virus Y	PVY	potyvirus	Aphid
tobacco etch virus	TEV	potyvirus	Aphid
pepper mild mottle virus	PMMoV	tobamovirus	seed, mechanical
tobacco mosaic virus	TMV	tobamovirus	seed, mechanical
tomato mosaic virus	ToMV	tobamovirus	seed, mechanical
tomato spotted wilt virus	TSWV	tospovirus	Thrips
beet curly top virus	BCTV	geminivirus	Leafhopper

## Tomato Spotted Wilt Virus (TSWV)

*Bryce Falk and Mike Davis, Plant Pathology, UC Davis, Michelle Le Strange and Scott Stoddard, Farm Advisors*

TSWV is a relatively recent and non-uniform problem in CA tomatoes and peppers. It is transmitted from plant to plant by at least 10 specific thrips vectors, including the western flower thrips (*Frankliniella occidentalis*), which is the most widespread and important vector for TSWV worldwide.

**The TSWV:Thrips Vector Transmission relationship** is different than most insect-transmitted plant viruses, and must be considered for any disease control strategy. Only the 1<sup>st</sup> or 2<sup>nd</sup> instar larvae (first stages after egg hatch) can acquire the virus from infected plants and then as adults transmit the virus to plant hosts. Adult thrips cannot acquire the virus; they can only spread it along IF they acquired it when young. Another fact is that the virus is not passed along from the adult to the egg. Thus, the eggs giving rise to the young nymphs must be on plants that are already TSWV infected. Inoculum sources must be hosts for both the thrips vector and the TSWV.

Knowing these facts we presume that important sources of TSWV inoculum are host plants that also support thrips populations. Unfortunately TSWV has one of the widest host ranges of any plant virus, infecting at least 168 plant species in 29 families. Economic hosts include tomatoes, peppers, celery, legumes, lettuce and many ornamentals; whereas weed hosts include nightshade, tree tobacco and jimson weed. Research is underway to determine the TSWV/thrips inoculum sources in some areas of the SJV. Please call a farm advisor, if you see significant incidence of this disease this season.

**Symptoms of the disease** vary, but young leaves tend to turn bronze, develop necrotic spots and streaks, and eventually, young shoots dieback and entire parts of the plant collapse and seem to wilt. One of the most diagnostic characteristics is the development of chlorotic or yellow ringspots on fruit; these rings are most obvious on red fruit, but also occur on green.

## Pepper Stip

*Joe Nuñez, UCCE Farm Advisor, Kern County*

### What is Stip?

Pepper stip, or color spotting, has become a serious problem for many bell pepper growers the past few seasons. There has been a lot of confusion as to what stip is and what it looks like. Pepper stip causes greenish-brown spots that are about 1/4 inch in diameter, slightly sunken below the surface on the fruit. They are most commonly seen on the mature red fruit but occasionally occur on green fruit as well. The spots appear just as the fruit begins to turn red.

Here in the southern San Joaquin Valley, stip has been prevalent on hybrid elongated red bell types or the so called "Maccabi" types. It also seemed to occur in early summer after a warm spring. This is contradictory to the report of stip in other pepper growing regions of the state. Along the coast and in Northern California, pepper stip is described as occurring after a cool period during the short days of fall on blocky open pollinated green bell types. Stip is apparently a mysterious disease of which very little is known.

Some things can be said about pepper stip. It is a physiological disorder that seems to be dependent on the environment for it to occur. Here the warm spring weather may have triggered it; in other parts of the state short, cool days are required before stip appears. It also appears to be a calcium imbalance, similar to blossom end rot. However, some reports say it is due to lack of calcium in the fruit while others report too much calcium in the fruit. There is also a difference in varietal susceptibility. In the southern San Joaquin Valley and in the southern deserts we know that the elongated Maccabi types are very susceptible to stip while blocky types are resistant. Other parts of the state report that it is on the blocky types that stip is found.

### Earlier trials

Stip research trials conducted in San Benito and San Joaquin Counties by Farm Advisors Richard Smith and Bob Mullen, in 1998 showed some interesting results. They learned two important things that may be beneficial to pepper growers here. First, that there are differences in susceptibility to stip between varieties and secondly that the incidence of stip could be reduced in the most susceptible varieties with calcium applications.

A variety trial evaluating open pollinated varieties for tolerance to pepper stip was conducted in San Benito County. The cultivar Gusto was nearly completely resistant and it was followed by varieties that were intermediated in susceptibility to pepper stip: Taurus and Cal Wonder 300. The remainder of the varieties tested (Yolo Wonder A and B, Jupiter, Keystone Resistant Giant, Grande Rio 66, Mercury, Pimlico, Loribelle, Capistrano, Merced, Emerald Giant, D-93, and Pip) were susceptible to stip. Foliar calcium applications beginning at first flower and continuing weekly until fruit began to turn red did not affect the incidence of pepper stip in San Benito County. However foliar applications did reduce the incidence of pepper stip in San Joaquin County. Five applications of foliar calcium at 0.5 gallon per acre of Cal Max reduced the incidence of pepper stip on Grande Rio and Yolo A by 85 and 60%, respectively.

Because so little is known about stip it is difficult to make management recommendations. Right now the only recommendations can be made are to plant varieties that are less susceptible to the disorder and maintain adequate but not excessive amounts of calcium.

## **Beware of Psyllids: Tomatoes, Peppers, Eggplant, & Potatoes at Risk**

*John Trumble, Entomologist, UC Riverside and Eric T. Natwick, UCCE Farm Advisor, Imperial County*

The tomato psyllid, *Paratrioza cockerelli*, also known as the potato psyllid has been showing up in pepper fields throughout southern California. PCAs and growers should be aware that nymphs have been found infesting peppers and other crops in the family: Solanaceae, such as tomato, potato and eggplant are at risk of becoming infested.

Tomato psyllids resemble tiny cicadas with clear wings that rest roof-like over the back and are about 2 mm long. Adults are mostly dark brown or black with white markings and they jump when disturbed. Psyllids are related to whiteflies, aphids and leafhoppers. Eggs are deposited on the underside of the leaf along the edge and in the upper plant canopy. They are football-shaped and very small, a 10X hand lens is required to see them, and on a short stalk. Psyllid nymphs are flat, shades of light green to greenish-yellow, and fringed with short spines around the edge. They resemble immature soft scale, but unlike insects, they move when disturbed. The nymphs develop through five instars in as little as two weeks.

**Damage is caused only by psyllid nymphs.** They inject a salivary toxin that causes a plant response known as psyllid yellows. Symptoms include yellowing and an upward curling of leaflets nearest the stem on the top part of the plant. However, yellowing is the most common symptom, initially found on the leaf edges. Other symptoms include an overall yellowing with enlarged nodes, shortened internodes, and development of clusters of small leaves in the axillary buds that appear rosetted. If the nymphs are removed from the plant, the progression of the disease will stop.

There are no specific treatment thresholds established for tomato psyllids on tomatoes or peppers. Insecticides used for aphid control, such as pyrethroids, and Provado, also control psyllids.

## **Stink Bug Research in Tomatoes**

*Frank Zalom and Corin Peas, Dept of Entomology, UC Davis*

### **CONSERVATION BIOLOGICAL CONTROL**

A study was conducted to assess feasibility of enhancing biological control of stink bugs by introduction and management of a nectar resource for parasitoids. Three sites were identified in March, 2003, on fresh market tomato farms that had a history of stink bug problems. Each site was planted with mixed heirloom tomato varieties and managed organically. Buckwheat, the nectar source we had intended to use, was broadcast seeded during the week of April 18-25 at the row ends of treatment plots, but two sites on silty soil failed to establish. Because of the time needed to re-establish buckwheat at other sites or to re-seed buckwheat at the two sites which failed to establish, alyssum was substituted. Alyssum had been studied previously with respect to its ability to provide a nectar source for biological control agents in other crops. One alyssum and one control plot were randomly placed along each selected border. Alyssum was transplanted at all three sites on May 20 and 21. Five alyssum plants spaced at 6 inches were planted on the row end of 15 rows in each treatment plot.

Sentinel egg masses were used to evaluate stink bug egg parasitism in treatment and control plots. The egg masses were from colonies established from stink bug adults captured in tomato fields earlier in the season. Paper towels within each colony served as an oviposition substrate. On July 3 and July 31, 8 sentinel egg masses were placed at 0, 20 and 50 feet from the alyssum and control borders of each study field. On Sept 5, 10 egg masses were deployed at each distance. The egg masses were retrieved a week after they were placed in the

field, and placed individually into small perforated zip lock bags. The bags were kept at room temperature and observed for parasitism. Once signs of parasitism (black eggs) were observed parasitized egg masses were transferred to individual glass scintillation vials with a ventilated cap and held for parasitoid emergence.

Parasitism of consperse stink bug egg masses was observed in all treatment and control replicates (Table 1). Average parasitism rates ranged from approximately 4% to 50%. Significant differences in parasitism were detected in 2 of three sample periods. Sentinel egg mass parasitism was significantly higher ( $p < 0.05$ ) 1ft from alyssum borders than 1ft from the control in the second and third sample periods (Jul 31-Aug 6, Sept 5-12). Parasitism was also significantly higher ( $p < 0.05$ ) 20 ft from alyssum borders than in the control. No significant differences in parasitism were detected further than 20 ft from the border and during the first sample period. However, all dates and distances with the exception of two showed higher mean parasitism adjacent to alyssum than in the control, although not significant ( $p > 0.05$ ).

This study shows promise for the use of in-field nectar sources for the enhancement of stink bug egg parasitoids. We suspect that actual parasitism could be even higher than we measured due to the artificial nature of our egg placement and other factors, and the affect of the nectar source in enhancing other parasitoids and generalist predators. However, the effect of field scale application of this technique on actual parasitism, fruit quality and other insects is not known.

**Table 1. Stink bug egg parasitism measured by sentinel masses placed in fresh market tomatoes at 3 distances from borders during 3 sampling periods, 2003.**

\*Significantly different from control by t-test

Treatment	Dist. Ft	PERCENT (%) of eggs parasitized					
		July 3-9		July 31 - Aug 6		Sept 5-12	
		%	SE	%	SE	%	SE
Alyssum	1	11.4	± 5.7	53.3	± 2.0 *	43.3	± 3.5 *
Control	1	7.6	± 7.6	45.6	± 2.9	21.7	± 7.0
Alyssum	20	9.9	± 1.6	39.1	± 6.6	49.8	± 8.2 *
Control	20	4.1	± 2.1	46.1	± 4.4	17.7	± 1.8
Alyssum	50	4.3	± 3.7	49.2	± 2.6	27.5	± 7.5
Control	50	4.8	± 4.8	28.8	± 11.9	22.5	± 9.5
Alyssum	All	8.5	± 1.9	47.2	± 3.2	40.2	± 5.6
Control	All	5.5	± 2.9	40.2	± 4.5	20.7	± 6.0

## CULTURAL CONTROL RESEARCH

A study was conducted to assess the role of springtime weeds which host stink bugs on the stink bug densities later observed in adjacent tomato fields. Eight processing tomato fields that were scouted during March and April, 2003, were selected for this study. Each field had one adjacent border area that was comprised of greater than 50% weedy hosts (cheeseweed, wild radish and mustard) of stink bugs and an opposing border where the weeds had been controlled early in the spring.

Each border replicate was sampled using tray shake samples to determine stink bug densities in July and again in August or early September. Fruit damage estimates were made following the second sampling date. Five tray shake samples were taken at 8 monitoring sites along both treatment borders of each field. These in-field monitoring sites were within 60 feet (12 rows) of both opposing field margins, and nearer to the center of the border to avoid edge effects from other borders. At harvest, 200 fruit were sampled from each border area, and individual fruit scored for fruit damage by stink bugs. The 200 fruit were a composite of 25 fruit collected at each in-field monitoring site where the stink bugs were also sampled. Treatments were analyzed by one-way ANOV of treatments using each field as a replicate.

Stink bugs were detected in all 8 fields and fruit damage in the border areas sampled ranged from 9.5% to 67% (Table 2). The majority (81%) of the stink bugs found at all sites were the consperse stink bug. Most of the others captured were the red shouldered stink bug, *Thyanta pallidovirens*, however the data presented here is solely for the consperse stink bug. Four of the fields were treated with insecticides for stink bugs between the first and second sampling periods and these are noted in Table 2. Stink bugs were significantly higher ( $p < 0.05$ ) in tomatoes adjacent to weedy host borders than in tomatoes adjacent to a non-host borders during the first sample period (Table 3). Average number of bugs per tray shake during the second sampling period and percent damage were

higher in the host border treatment than in the non-host border treatment, but the differences were not significant ( $p > 0.05$ ).

The results of this study demonstrate the importance of spring weed hosts nearby tomato fields to the incidence of stink bugs in tomatoes. Stink bugs were not precluded from colonizing tomatoes near borders without weed hosts, yet the tomatoes adjacent to these borders had lower incidence early season. It seems clear that control of weeds nearby future tomato fields and on a greater landscape scale can be expected to reduce damaging populations in tomatoes. These weed hosts must be removed prior to tomato seedling establishment or transplanting of the fields. The results of this study suggests that management of weed hosts that may function as a trap crop for stink bugs could be beneficial for growers. However, there are certainly confounding factors that could play a role in the success of such an approach. Location of additional overwintering habitat outside of the immediate field borders that is beyond the grower's control, such as riparian areas and orchard floor vegetation, may also play a role.

**Table 2. Number of stink bugs per tray shake and percent stink bug damage at field borders in relation to availability of an adjacent host.**

Field	Treat.	# of bugs/ shake		% damage
		July	Aug/Sept	
7	Host	0.075	0.225	38.0
7	No host	0	0.100	25.5
11A	Host	0.075	1.950	58.5
11A	No host	0.025	0.875	51.5
1A	Host	0.150	0.100*	61.0
1A	No host	0.025	0.050*	40.5
1B	Host	0.050	0 *	32.5
1B	No host	0	0 *	23.5
B14	Host	0.200	0.025*	36.0
B14	No host	0.025	0 *	42.6
B2	Host	0.025	0.025*	45.5
B2	No host	0	0.125*	67.0
LT	Host	0.500	0.900	62.0
LT	No host	0.075	1.075	56.0
R1	Host	0.350	0.600	38.0
R1	No host	0.025	0.025	9.5

\* fields treated with insecticide for stink bugs

**Table 3. ANOV statistics for stink bugs per tray shake and percent stink bug damage at field borders in relation to availability of an adjacent host.**

Treatment	df	F=	p=
Host vs Non Host (1 <sup>st</sup> Sample Period)	1,14	6.84	0.02
Host vs Non Host (2 <sup>nd</sup> Sample Period)	1,14	0.48	0.50
Host vs Non Host (% Fruit Damage)*	1,14	0.67	0.43

## CHEMICAL CONTROL STUDIES

For the past several years, we have been evaluating alternative pesticides which could serve as a replacement for the organophosphates methamidophos (Monitor) and dimethoate for control of stink bugs. Insecticides tested include pyrethroids, neonicotinoids, insect growth regulators, and various tank mixes of products.

### FIELD PLOTS

Stink bug populations were established in the tomato plots by artificial infestation using methods we developed for several years. The source of the stink bugs was a colony we established in spring from adults collected at several sites. For our field infestation, egg masses from the colony were collected during a 4 day period and taken to the field where the paper towel strips containing the egg masses were attached to the undersides of leaves within the plant canopy. One egg mass was placed for each 10 foot length of tomato row in the experimental planting between July 22 and July 25 which approximated the predicted harvest date for stink bugs in the Davis area using the degree-day phenology model developed by Cullen and Zalom. The model can be found at UCIPM (<http://www.ipm.ucdavis.edu>).

The insecticides tested were applied on July 12 (Fulfill and Platinum, only) or August 1. All treatments but Platinum were applied using an Echo Duster/Mister air assist sprayer at a volume equivalent to 50 gal/acre. Platinum was applied by spraying the product on the soil at the base of each plant in 4.25 oz. of water per plant using a Hudson pump sprayer. The plots were furrow irrigated the day after the soil application was made. Treatments were assigned to plots in a completely randomized design, with three replicates per treatment and six replicates of the untreated control. Stink bug treatment efficacy was evaluated by shake sampling and determining

damage at harvest. Five shake samples were taken from each replicate on August 11, ten days following the application of most treatments. Fruit damage was evaluated on September 10, by scoring 100 red fruit that were randomly sampled from each replicate for stink bug feeding. One of the 6 untreated plots could not be evaluated because damage and secondary fungal infections were so high that only a few of the fruit maintained sufficient integrity to allow scoring to proceed.

### RESULTS

Differences between treatments were confirmed by 1-way ANOV for both number of stink bugs per tray shake sample and damage at harvest (**Table 4**). Significant treatment differences in number of stink bugs per tray shake relative to the untreated control were found for Dibrom, Warrior + Actara, Warrior + Platinum, F1785 (flonicamid, an unregistered product from FMC), MustangMax + F1785, Warrior + Assail, Lannate + Danitol, and Knack + Danitol. Significant treatment differences in damage at harvest relative to the untreated control were found for Warrior + Actara, MustangMax, MustangMax + F1785, Warrior + Assail, and Lannate + Danitol. That the combination of Lannate and Danitol was significant is interesting given that in field tests in 1986 and 1987 we showed that the combination of Lannate and Asana (a pyrethroid as is Danitol) also provided significant control of stink bugs. In those trials, neither Lannate nor Asana provided significant control of stink bugs when applied alone.

### SUMMARY

These results confirm for the most part our previous observations that combinations of certain pyrethroid insecticides such as Warrior with a neonicotinoid pesticide afford better control of stink bug than do other alternatives.

**Table 4. Number of stink bugs per 5 tray shake samples on August 11, percent damage at harvest (September 10) UC Davis, 2003.**



Treatments applied on August 1, except as indicated.

1-way ANOV results:

$F=1.719$ ,  $df=21,38$ ,  $p=0.0327$  for tray shake samples;

$F=2.150$ ,  $df=18,40$ ,  $p=0.0220$  for damage at harvest.

<sup>1</sup>Application date for this product: July 12.

<sup>2</sup>Not registered for use on tomatoes.

\*Formulated rate/acre

\*\*Means are significantly different ( $p<0.05$ )

from untreated by t-test following  $\log(x+1)$  transformation

Treatment	Rate (a.i./ac)	No. of bugs 5 tray shakes 8/11	% Damage at harvest 9/10
Untreated	NA	2.3 ± 0.6	73 ± 9
Dibrom 8	1 pt*	0.9 ± 0.2 **	57 ± 9
Fulfill <sup>1</sup>	2.75 oz*	1.4 ± 0.4	70 ± 16
Activol	3.00%	2.1 ± 0.5	65 ± 26
Warrior	3.84 oz*	1.7 ± 0.6	52 ± 5
Actara	4.0 oz*	1.3 ± 0.4	66 ± 18
Warrior + Actara	3.84 oz* + 4.0 oz*	1.0 ± 0.4 **	30 ± 18**
Platinum <sup>1</sup>	8.0 oz*	1.2 ± 0.4	69 ± 13
Warrior + Platinum <sup>1</sup>	3.84 oz* + 8.0 oz*	0.5 ± 0.2 **	49 ± 23
MustangMAX (L)	0.018 lb	1.6 ± 0.9	51 ± 22
MustangMAX (H)	0.025 lb	1.5 ± 0.5	46 ± 19**
F1785 (L) <sup>2</sup>	0.054 lb	0.8 ± 0.3 **	62 ± 21
F1785 (H) <sup>2</sup>	0.071 lb	1.2 ± 0.3	58 ± 18
MustangMAX (H) + F1785 (H) <sup>2</sup>	0.025 lb + 0.071 lb	0.6 ± 0.3 **	35 ± 06**
Assail 70WP	2.39 oz*	1.8 ± 0.5	65 ± 23
Warrior + Assail	3.84 oz* + 2.39 oz*	0.7 ± 0.4 **	31 ± 09**
Dimethoate 4EC	1.5 pts*	1.8 ± 0.4	76 ± 11
Lannate + Danitol	0.9 lb + 0.2 lb	0.0 ± 0.0 **	36 ± 04**
Knack 0.86 EC + Danitol	0.054 lb + 0.2 lb	0.6 ± 0.2 **	48 ± 28

# Root-knot Nematode Resistant Tomatoes not always Resistant.

*Antoon Ploeg, Nematology Specialist, UC Riverside*

Root-knot nematodes (*Meloidogyne* species) are the most important nematode parasites of tomato, and although a large number of root-knot nematode species exist, only a few species are important. Warm-loving species such as *M. incognita* and *M. javanica* are usually found in greenhouse production in Northern California and in the field in Central and Southern California. Another common species, *M. hapla*, prefers cooler conditions and is more widely distributed in the Northern part of the state.

Nematodes multiply rapidly on their hosts when temperatures are favorable, and populations can increase from hardly detectable to very high levels (>10,000 per pint of soil) within one growing season. The nematodes are closely associated with the host roots, which they modify to produce typical galling. The developing nematodes initially occur within these galls, but later deposit numerous egg masses, containing up to 400 eggs, that are “glued” to the outside of the roots.

Nematodes take nutrients away from the plant, and the damaged roots become an easier target for infection by fungal and bacterial pathogens. In other crops (e.g. melon) it has been shown that nematode infection in the early stages of plant growth is particularly damaging, whereas established healthy plants can tolerate a certain level of nematode infection without plant growth becoming affected. However, even plants that initially escape nematode infection and show no obvious above-ground symptoms may still exhibit severe galling and harbor large nematode numbers at harvest time.

The use of fumigant nematicides has long been an important tool in nematode management, but fewer products are available, and costs may be prohibitive. Alternatively, nematode-resistant cultivars can be grown and fortunately, tomato is one of a few crops where nematode-resistant varieties are available.

**Nematode resistance:** In all nematode-resistant tomato varieties the resistance is based on one gene (*Mi*-gene). As a management strategy the use of nematode-resistant varieties has several advantages:

- it requires no major changes in farming practices,
- it reduces the need for chemical control,
- it is reliable, and
- it has a positive effect on a following nematode-susceptible crop because it lowers the nematode populations.

There are however a few drawbacks:

- 1) Although nematode-resistant tomato varieties are resistant to three very important warm-climate root-

knot nematode species (*M. javanica*, *M. incognita*, *M. arenaria*), they are not resistant to *M. hapla*.


- 2) Resistance is specifically against root-knot nematodes, and not against other types of nematodes.
- 3) Resistance breaks down when soil temperatures reach 82°F, and thus may not be useful in hot desert-type conditions.
- 4) There are a limited number of resistant varieties, and thus for certain desired tomato types resistant varieties may not be available.

**Grafting tomatoes?** To circumvent this last limitation, susceptible tomato with the desired fruit type can be grafted onto nematode-resistant rootstocks. This practice is very common in high value greenhouse tomato production in Asian and European countries. Recently we studied the effect of grafting on fruit production and nematode population levels using a California root-knot nematode population (*M. incognita*), greenhouse tomato variety Blitz, and nematode-resistant rootstock variety Beaufort. As expected, on the non-grafted susceptible controls, fruit yields decreased, galling increased, and nematode populations were higher at harvest as nematode inoculum levels increased. The grafted tomatoes did not exhibit yield loss, even at high inoculum levels, but surprisingly, root galling and nematode levels were only slightly lower than on the susceptible controls. Soil temperature had remained below the critical level of 82F, the *Mi*-gene was present in the rootstocks, and the root-knot nematodes had not previously been exposed to resistant tomato. Therefore, high-temperature resistance breaking, a seed mix-up (the rootstock did not really contain the resistance gene), or the use of a highly virulent resistance-breaking population as reasons for the unexpected results were excluded.

In a second experiment, another nematode-resistant tomato variety (Hypeel45) was included as a rootstock, as well as another root-knot nematode population. Results from this experiment showed that both resistant rootstocks prevented yield loss at increasing inoculum densities, but that rootstock Beaufort again showed severe galling and allowed nematode multiplication. Rootstock Hypeel45 however only had very minor galling and only few nematodes were recovered from this rootstock, even at very high nematode inoculum densities.

**Summary:** We conclude that one of the “resistant” rootstocks (Beaufort) was tolerant rather than resistant, leaving a high nematode population behind at harvest. This may have important consequences for the performance of a following susceptible crop. At this moment we do not have an explanation for the difference in galling and nematode levels between the two “resistant” rootstocks both containing the nematode resistance *Mi*-gene.

## U.S. Per Capita Utilization of Fresh Bell Peppers and Tomatoes, 1985-2003 (with and without U.S. Greenhouse tomato production added to field grown as of 1998)



Source USDA/ERS July 2004, Vegetable Yearbook  
Roberta Cook and Linda Calvin estimated '98-03 tomato consumption to reflect unaccounted for Greenhouse tomato production.

## Trials with Preemergence Herbicides in Transplanted Bell Peppers

Richard Smith and Michelle Le Strange, UCCE Farm Advisors, Monterey and Tulare & Kings Counties

### INTRODUCTION

Peppers are long-season vegetables that have several weed control challenges: They compete weakly with weeds for the first 40 to 60 days following transplanting. They are a long-season crop in many production districts that can be subject to flushes of both winter and summer weeds over the course of their growing cycle.

The preemergence herbicides registered for peppers have gaps in the spectrum of weeds that they control. As a result, growers may spend from \$200 to \$350/acre on weed management. Field selection, field sanitation, cultivation and the use of plastic mulches are cultural practices that reduce weed pressure in production fields. Fumigation provides substantial weed control and is frequently used in conjunction with plastic mulches which improves the level of weed control provided by both techniques.

Goal Tender was registered in California in 2004 for use with plastic mulch and provides control of Little Mallow (*Malva parviflora*) which is only partially controlled by fumigants and other preemergence herbicides registered for use on peppers. However, many acres of peppers are not grown with plastic mulch, and weed control is a challenge. Devrinol, Prefar and Treflan are registered preemergence herbicides in peppers. Dual Magnum is registered under a 24C and provides good control of hairy nightshade (*Solanum sarrachoides*) and yellow nutsedge (*Cyperus esculentus*) which are not controlled

by the other preemergence materials. Late season weed control is also an important issue in this crop.

The objective of these studies was to examine at transplant and layby herbicide combinations for peppers that can provide long-term and economical weed control for peppers grown without plastic mulch. The herbicides tested included: Dual Magnum 7.62 (s-metolachlor), Goal Tender 4F (oxyfluorfen), Outlook 6.0 (dimethenamid), flumioxazin (Chateau) impregnated on fertilizer, and Dacthal 75W (DCPA).

### METHODS

Field trials were conducted on the Central Coast (Monterey and Santa Clara Counties) and Fresno County in 2005 to provide an evaluation of the test herbicides over a wider range of growing conditions and weed spectra.

**Central Coast Trial:** The trial was conducted with a cooperating grower in Gilroy. Goal Tender treatments were applied onto shaped beds two weeks prior to transplanting the peppers on April 28. The field was transplanted on May 13. The at-planting treatments were applied over-the-top of the plants immediately following transplanting. Sprinkler irrigation was started 5 hours following transplanting applying 0.38 inch of water. Layby applications were made on June 16 and the material was incorporated with the last sprinkler irrigation before the field was switched to drip irrigation. The

plots were hand weeded on June 3 and the July 1 weed evaluations reflect newly sprouted weeds following the layby application. The plots were not cultivated prior to the July 1 weed evaluation. Each plot was one 40-inch bed wide by 25 feet long and replicated four times in a randomized complete block design (RCBD). All sprayed treatments were applied to the entire bed in 74 gallons of water per acre with two passes of a one nozzle wand with an 8008E teejet nozzle at 30 psi. Flumioxazin on fertilizer granules was spread by hand on the bed top immediately following transplanting. Soil type was Pacheco silt loam and the variety was Baron.

**Fresno County Trial:** The field trial was conducted on a Panoche clay loam soil at the UC West Side Research and Extension Center (WSREC) near Five Points. On June 7 the bell pepper variety “Jupiter” was transplanted in single rows into 40” beds. Within row plant spacing was 10”. Plot size was two 40-inch beds x 70 feet of row length and replicated 4 times in a RCBD.

Preplant applications of Goal Tender were made onto shaped beds on May 10, 28 days prior to transplanting the peppers and incorporated with 0.50 inches of rainfall. The at-planting treatments of Dual Magnum, Outlook and flumioxazin were applied over the top of the plants and the field was sprinkler irrigated applying 0.50 inches of water immediately following transplanting. Sprinkler irrigation continued as needed for a few weeks and then switched to furrow irrigation. On July 25 the field was machine cultivated before layby applications of the herbicides were made as a directed spray to the base of the plants. These applications were incorporated by sprinkler irrigation.

All sprayed treatments were applied to the entire plot in 30 gallons of water per acre using a CO<sub>2</sub> backpack sprayer @30 psi and a 2 nozzle boom with 8003evs tips. Flumioxazin on fertilizer granules was hand broadcast over the top of the peppers. There were two untreated checks: one was handweeded twice in addition to layby cultivation and the other was allowed to grow weedy all season.

Plots were evaluated for phytotoxicity to the peppers and weed control on July 1, July 22, and August 12. Pepper stand counts were made on July 13. A portion of each plot (25’ row) was hand harvested on August 23 (west bed) and on September 8 (east bed) and the yields were combined for total yield. Tables 2 & 3 list treatments and evaluations.

## RESULTS AND DISCUSSION

**Central Coast:** Hairy Nightshade was the dominant weed at this site. The best weed control was provided by Outlook, then by Dual Magnum and Goal Tender on the 21 days after treatment (DAT) (Table 1). Flumioxazin impregnated on fertilizer provided good weed control in two treatments, but not on one. This may be due to problems with obtaining an even distribution of this dry granular material on the top of the bed. Devrinol was at a distinct disadvantage at this site because Hairy Nightshade was the main weed at this site and it did not control it. Outlook caused stunting of the plants 21 DAT, and while the stunting was reduced 28 DAT, it was still quite noticeable. There was no difference in the stand among

treatments, but there were some instances of burned pepper plants in the flumioxazin treatment, presumably where a prill of the material lodged against the stem of a plant. All herbicides except Devrinol reduce time to weed the plots, but Goal and flumioxazin on fertilizer tended to take more time than Dual Magnum and Outlook. There were no differences in weed control among the layby applications (data not shown) and this test did not provide a good opportunity to evaluate the long-term weed control system for peppers. There were no significant differences in yield among the treatments (data not shown) which indicates that the initial phytotoxicity observed on the Outlook treatments did not translate to reduced yield.



Fresno: June 1, 2005 - Goal Tender provided very good weed control prior to transplanting peppers.

**Fresno County:** Because of excessive rainfall many weed seeds germinated in the untreated area of the field after the preplant treatments of Goal Tender but before the peppers were transplanted. Goal Tender was extremely effective in controlling all of the weeds (see photo). However, prior to transplanting it was necessary to cultivate and reshape the beds, thus destroying the herbicide layer and the effectiveness of Goal Tender. Weed control ratings on July 1 and July 22 (a few days before layby) showed how Goal Tender was no longer effective (Table 3).

Weeds were vigorous and abundant throughout the season and included several broadleaf species and virtually no grasses except for occasional jungle rice (*Echinochloa colonum*). The major broadleaf weeds were prostrate, tumble, and redroot pigweeds (*Amaranthus blitoides*, *A. albus*, and *A. retroflexus*); primarily black nightshade (*Solanum nigrum*), but also some hairy nightshade (*S. sarrachoides*) and lanceleaf groundcherry (*Physalis lanceifolia*); common lambsquarters (*Chenopodium album*); and purslane (*Portulaca oleracea*). Mustards, shepherds-purse (*Capsella bursa-pastoris*) and London rocket (*Sisymbrium irio*), were initially present prior to layby, but were taken out with the layby cultivation and were not serious competitors. Puncturevine (*Tribulus terrestris*) was also scattered throughout the experimental site but was not included in the weed counts because its populations were too random.

**At planting applications:** Although weed control was initially excellent, Outlook really hurt the peppers with an over the top application and many plants remained stunted for the entire season. Pepper yields were extremely reduced. As

the season progressed weeds germinated and the small pepper plants offered little competition. Flumioxazin provided good weed control and only slight pepper phytotoxicity was observed using the dry granular formulation, although some care was given to try to keep the prills off of the pepper plants during the broadcast application. Weed control is probably compromised by this method of application due to the difficulty of obtaining uniform coverage. Dual Magnum provided the best weed control. A little damage was seen on the peppers, but yields were not affected.

**Layby applications:** After layby there was not a lot of new weeds that germinated however, weeds that were missed by cultivations continued to grow. Dual, Outlook, and Dacthal all provided good to excellent weed control when applied at layby. All of the Goal Tender preplant plots and the flumioxazin at planting plots were improved with the layby applications. Dual, Outlook, and flumioxazin were effective on nightshades, and reduced pigweeds, purslane, and lambsquarters populations to varying degrees, although none of these products provided complete control of these weeds in this experiment. Still a hand weeding crew would have been able to clean up the field in a relatively short time, if the pepper field had been treated with almost any of these combinations.

## CONCLUSION

The Central Coast trial provided evidence that Goal Tender applied to shaped beds prior to transplanting (and subsequently not worked prior to transplanting) provided acceptable safety to the peppers and good weed control. This use pattern could provide an alternative “at planting” treatment and can provide weed control for the first 30 days following transplanting. Outlook was applied over-the-top in both trials, but was more damaging to the peppers in the Fresno trial. This material did not reduce yields in the Central Coast trial and should be further examined as a pretransplant application. Both trials showed that flumioxazin impregnated on fertilizer has promise as a post transplant application on peppers. The Fresno Trial showed that Dual Magnum, Outlook and Dacthal all provided good layby weed control. Dacthal is already registered for this use, but the Dual Magnum label would need to be adjusted to allow this use.

In summary, these trials showed promise for developing a weed control system to provide early and late season weed control for peppers grown without plastic.

**Table 1. Central Coast Trial. Post transplant evaluations**

Code	Applications	Lbs a.i./A	Material/A	Night-shade 21 DAT	Total Weeds 21 DAT	Phyto 21 DAT	Phyto 28 DAT	Plants per plot 21 DAT	Time to weed (hrs/A) 21 DAT
1	Dual Magnum 7.62 <i>Fb</i> * Dual Magnum 7.62	1.43 1.43	1.50 pts 1.50 pts	0.5	0.5	0.3	0.0	36.8	1.6
2	Dual Magnum 7.62 <i>Fb</i> Outlook 6.0	1.43 0.60	1.50 pts 0.80 pt	0.8	0.8	0.8	0.3	35.5	1.3
3	Dual Magnum 7.62 <i>Fb</i> Dacthal 75W	1.43 7.00	1.50 pts 9.3 lbs	1.3	1.3	0.3	0.1	36.0	1.3
4	Goal Tender 4F <sup>1</sup> <i>Fb</i> Dual Magnum 7.62	0.50 1.43	1.00 pt 1.50 pts	2.3	3.3	0.8	0.4	35.3	3.6
5	Goal Tender 4F <sup>1</sup> <i>Fb</i> Outlook 6.0	0.50 0.60	1.00 pt 0.80 pt	2.5	2.8	0.3	0.0	35.0	3.3
6	Goal Tender 4F <sup>1</sup> <i>Fb</i> Dacthal 75W	0.50 7.00	1.00 pt 9.3 lbs	2.3	2.3	0.5	0.0	35.8	2.8
7	Outlook 6.0 <i>Fb</i> Dual Magnum 7.62	0.60 1.43	0.80 pt 1.50 pts	0.3	0.3	1.0	1.3	34.8	1.2
8	Outlook 6.0 <i>Fb</i> Outlook 6.0	0.60 0.60	0.80 pt 0.80 pt	0.3	0.3	1.5	0.8	35.3	1.2
9	Outlook 6.0 <i>Fb</i> Dacthal 75W	0.60 7.00	0.80 pt 9.3 lbs	0.5	0.5	1.3	0.8	36.5	1.1
10	Flumioxazin impregnated on fertilizer <i>Fb</i> Dual Magnum 7.62	0.094 1.43	188 lbs 1.50 pts	1.0	1.3	1.3	1.0	35.0	2.2
11	Flumioxazin impregnated on fertilizer <i>Fb</i> Outlook 6.0	0.094 0.60	188 lbs 0.80 pt	1.0	1.0	1.5	1.0	35.5	2.2
12	Flumioxazin impregnated on fertilizer <i>Fb</i> Dacthal 75W	0.094 7.00	188 lbs 9.3 lbs	4.0	4.0	0.8	0.5	36.5	3.4
13	Devrinol <i>Fb</i> Dacthal 75W	1.50 7.00	3.0 lbs 9.3 lbs	11.8	12.0	0.0	0.0	35.8	8.4
14	Untreated	---	---	11.8	13.3	0.0	0.0	36.3	7.7
	LSD (0.05)			3.8	3.7	1.4	1.0	NS	2.5

<sup>1</sup> – applied 16 days prior to transplanting. \* *Fb* = Followed by

**Table 2. Fresno Trial. Weed control ratings and Weed counts**

	Applications			Weed CONTROL Ratings *			Weed Counts per plot				TOTAL Brdlvs
	Preemergence Herbicides	Lbs a.i. per Acre	Material per Acre	---- all broadleaf weeds ----			August 12, 2005* (67 DAT)				
				24 DAT	45 DAT	67 DAT	Pig	Night	Purs	Lamb	
1	Dual Magnum 7.62 <i>Fb Dual Magnum</i>	1.43 1.43	1.5 pt 1.5 pt	9.8 a	9.2 a	9.6 ab	10.7	0.0	6.7	2.0	19.3 ab
2	Dual Magnum <i>Fb Outlook</i>	1.43 0.60	1.5 pt .75 pt	10.0 a	9.5 a	8.5 bc	15.0	0.3	3.7	3.0	22.0 ab
3	Dual Magnum <i>Fb Dacthal</i>	1.43 7.00	1.5 pt 9.5 lb	10.0 a	9.5 a	9.7 a	13.0	0.3	3.7	0.0	17.0 a
4	Goal Tender 4F <sup>1</sup> <i>Fb Dual Magnum</i>	0.50 1.43	1 pt 1.5 pt	1.3 c	4.0 c	8.7 abc	13.7	7.0	2.3	0.7	23.7 ab
5	Goal Tender <sup>1</sup> <i>Fb Outlook</i>	0.50 0.60	1 pt .75 pt	1.5 c	6.2 b	9.3 abc	6.0	2.3	2.7	1.7	12.7 a
6	Goal Tender <sup>1</sup> <i>Fb Dacthal</i>	0.50 7.00	1 pt 9.5 lb	1.0 c	4.0 c	8.8 abc	13.7	2.7	2.0	0.3	18.7 ab
7	Outlook 6.0 <i>Fb Dual Magnum</i>	0.60 1.43	.75 pt 1.5 pt	10.0 a	9.0 a	8.3 c	32.3	0.0	5.3	4.0	41.7 bc
8	Outlook <i>Fb Outlook</i>	0.60 0.60	.75 pt .75 pt	10.0 a	9.6 a	8.5 bc	21.7	0.0	3.7	5.7	31.0 ab
9	Outlook <i>Fb Dacthal</i>	0.60 7.00	.75 pt 9.5 lb	9.7 a	9.5 a	7.0 d	37.0	1.0	9.7	8.3	56.0 c
10	Flumioxazin impregnated on fertilizer <i>Fb Dual Magnum</i>	0.094 1.43	150 lbs 1.5 pt	7.7 b	8.3 a	9.0 abc	15.0	0.0	3.0	4.7	22.7 ab
11	Flumioxazin <i>Fb Outlook</i>	0.094 0.60	150 lbs .75 pt	9.7 a	8.7 a	8.8 abc	20.7	0.0	3.3	5.0	29.0 ab
12	Flumioxazin <i>Fb Dacthal</i>	0.094 7.00	150 lbs 9.5 lb	8.2 b	8.3 a	8.8 abc	18.3	1.3	4.7	2.0	26.3 ab
13	Untreated - weeded			0.7 c	8.3 a	6.7 d	32.3	19.7	3.3	1.7	57.0 c
14	Untreated - weedy			1.0 c	1.7 d	0.7 e	64.3	24.0	24.7	17.0	130.0 d
	LSD (0.05)			1.0	1.5	1.2	20.5	4.0	5.3	6.5	24.4

1- Applied 28 days before transplanting. \* July 1=24 DAT; July 22=45 DAT; Aug 12=67 DAT. Aug 12 =18 days post layby application.

**Table 3. Fresno Trial. Pepper yield, Stand counts, and Phytotoxicity ratings\***

	Applications	Lbs a.i. per Acre	Material per Acre	Pepper Yield lbs/plot				Peppers/plot 36 DAT	Phytotoxicity	
				Good	Small	Sun- burn	Total		24 DAT	67 DAT*
1	Dual Magnum 7.62 <i>Fb Dual Magnum</i>	1.43 1.43	1.5 pt 1.5 pt	77.6 ab	18.7	15.9	112.2 ab	186.0 bcd	1.3 b	0.0 d
2	Dual Magnum <i>Fb Outlook</i>	1.43 0.60	1.5 pt .75 pt	71.3 ab	18.4	13.9	103.5 ab	184.5 cd	1.5 b	3.5 bcd
3	Dual Magnum <i>Fb Dacthal</i>	1.43 7.00	1.5 pt 9.5 lb	61.0 ab	16.9	11.7	89.5 ab	194.5 abcd	1.0 b	0.0 cd
4	Goal Tender 4F <sup>1</sup> <i>Fb Dual Magnum</i>	0.50 1.43	1 pt 1.5 pt	62.6 ab	20.7	14.3	97.6 ab	186.5 bcd	0.2 b	0.3 d
5	Goal Tender <sup>1</sup> <i>Fb Outlook</i>	0.50 0.60	1 pt .75 pt	91.6 a	12.6	11.8	116.0 a	196.5 abc	0.5 b	0.0 d
6	Goal Tender <sup>1</sup> <i>Fb Dacthal</i>	0.50 7.00	1 pt 9.5 lb	66.0 ab	12.9	10.2	89.1 ab	200.0 ab	0.2 b	0.0 d
7	Outlook 6.0 <i>Fb Dual Magnum</i>	0.60 1.43	.75 pt 1.5 pt	47.2 bc	8.9	9.5	65.5 bc	190.5 abcd	3.7 a	6.0 ab
8	Outlook <i>Fb Outlook</i>	0.60 0.60	.75 pt .75 pt	41.8 bc	10.7	13.7	66.2 bc	187.0 bcd	3.7 a	2.0 abc
9	Outlook <i>Fb Dacthal</i>	0.60 7.00	.75 pt 9.5 lb	21.0 c	7.1	3.7	31.9 c	188.0 bcd	5.2 a	6.5 a
10	Flumioxazin impregnated on fertilizer <i>Fb Dual Magnum</i>	0.094 1.43	150 lbs 1.5 pt	57.5 abc	15.5	11.3	84.4 ab	182.0 cd	1.0 b	1.0 cd
11	Flumioxazin <i>Fb Outlook</i>	0.094 0.60	150 lbs .75 pt	79.4 ab	15.4	17.6	112.3 ab	194.0 abcd	0.5 b	0.5 d
12	Flumioxazin <i>Fb Dacthal</i>	0.094 7.00	150 lbs 9.5 lb	65.3 ab	16.3	7.3	89.0 ab	203.5 a	1.5 b	2.0 bcd
13	Untreated - weeded			78.4 ab	19.2	10.0	107.5 ab	181.5 d	0.3 b	0.3 d
14	Untreated - weedy			20.9 c	6.2	4.5	31.5 c	182.5 cd	0.0 b	0.0 d
	LSD (0.05)			39.8	8.8	9.1	47.4	14.6	1.9	2.8

1- Applied 28 days before transplanting. \* July 1=24 DAT; July 13=36 DAT; July 22=45 DAT; Aug 12=67 DAT. Aug 12 = 18 days post layby.

# Postharvest Handling Considerations for Grape Tomatoes

Marita Cantwell, UCCE Postharvest Vegetable Specialist, Mann Laboratory, UC Davis

During 2003 and 2004 postharvest research was conducted on grape tomatoes. These small grape-size tomatoes have high flavor potential and are now popular among consumers. As in other tomatoes, stage of maturity at harvest is critical to maximize eating quality. **Table 1** shows the difference in composition when the fruit were harvested at different maturity stages (3, 4, and 5) and ripened to full red color (table-ripe stage). Although the % soluble solids (°Brix) differed considerably among the 3 maturity stages, the actual sugar concentrations were much less different. As expected, sugars are correlated to % soluble solids. The fruit harvested at higher maturity stage had higher % titratable acidity in Test 1 but not Test 2. No significant differences were noted in Ascorbic acid (Vit. C) concentration among fruit ripened from different stages of maturity. There were no significant differences at the table-ripe stage in fruit weight, red color or firmness in relation to the 3 stages of harvest (data not shown).


Storage temperature will be a critical factor in maintaining the postharvest quality of fruit. Changes in visual appearance, color (objective color value, the lower the hue value, the redder the fruit), firmness (the lower the value, the softer the fruit) and % weight loss of stage 4 fruit packaged in clamshells in relation to storage at 4 temperatures is shown in **Figure 1**. As with all tomatoes, a storage temperature of 41°F is too low as evidenced by the lack of color development. Storing at 50°F retarded color development. Firmness changes were similar among the fruit stored at 41, 50 and 59°F. In other tests, we stored fruit at 55°F and consider this the best storage temperature (slow softening and color change but not prevent eventual normal color development). Shriveling and firmness loss is closely correlated with % weight loss (data not shown).

A complete report is available at the Postharvest Website) <http://postharvest.ucdavis.edu>

**Table 1.** Composition of table-ripe grape tomatoes (cv Amsterdam) harvested at different stages of maturity and ripened at 69°F.

Initial color stage	% Soluble solids	Sugars mg/ml juice	pH	% Titratable acidity	SS:TA ratio	Vitamin C mg/100ml
<b>Test 1</b>						
Stage 3	5.5	30.1	4.37	0.51	10.8	92.6
Stage 4	6.0	30.7	4.36	0.53	11.3	90.4
Stage 5	7.5	33.6	4.49	0.57	13.2	85.8
LSD.05	0.8	2.8	0.12	0.05	--	ns
<b>Test 2</b>						
Stage 4	7.0	30.9	4.28	0.65	10.8	57.4
Stage 5	7.6	34.3	4.48	0.53	14.3	63.4
LSD.05	0.5	3.4	0.07	0.06	--	ns

**Figure 1:** Visual quality, weight loss, firmness and color of grape tomatoes (cv. Amsterdam) sorted in vented clamshells at 4 Temps.



# SOURCES OF INFORMATION – TOMATOES & PEPPERS

## PUBLICATIONS FROM UC

Many items are available at no cost from local UCCE offices or the World Wide Web.

**UC Vegetable Research & Information Center**  
(UC VRIC) [www.vric.ucdavis.edu](http://www.vric.ucdavis.edu)

**UC IPM (homepage)**  
[www.ipm.ucdavis.edu](http://www.ipm.ucdavis.edu)

**UC Weed Research & Information Center:**  
(UC WRIC) [www.wric.ucdavis.edu](http://www.wric.ucdavis.edu)


**UC Postharvest Technology:**  
<http://postharvest.ucdavis.edu>  
(be sure to browse the Produce Facts)

**UC Ag Economics: Cost of Production Guidelines**  
<http://coststudies.ucdavis.edu> or (530) 752-1515

**UC Ag & Natural Resources Catalogue**  
<http://anrcatalog.ucdavis.edu>

**IPM Tomato Manual, #3274**  
**IPM Tomato Pest Management Guidelines #3470**  
*Identification & Management of Complex Tomato Diseases (available through UC VRIC)*

**Fresh Market Tomato Publication in CA, #8017**  
**Processing Tomato Production in CA, #7228**  
**Bell Pepper Production in CA, #7217**  
**IPM Pepper Pest Management Guidelines #3460**  
*Scheduling Irrigation: When & How Much, #3396*



**Mark Your Calendars**  
**May 21-23, 2006**

Renaissance Esmeralda Resort  
Palm Springs, CA

[www.internationalpepper.com](http://www.internationalpepper.com)

## INDUSTRY ORGANIZATIONS

### California Tomato Commission

[www.tomato.org](http://www.tomato.org)  
Fresh Market Tomato Industry  
1625 E. Shaw Avenue, Suite 106  
Fresno, CA 93710  
(559) 230-0116

### California Pepper Commission

531-D North Alta Avenue  
Dinuba, CA 93618  
(559) 591-3925

## WEATHER & IRRIGATION

CIMIS - CA Irrigation Management & Info System  
CA Dept Water Resources - [www.cimis.water.ca.gov](http://www.cimis.water.ca.gov)

## GOVERNMENT

CDFA - [www.cdfa.ca.gov](http://www.cdfa.ca.gov)  
CDPR - [www.cdpr.ca.gov](http://www.cdpr.ca.gov)

CA AG Statistics Services - <http://www.nass.usda.gov/ca>  
Curly Top Virus Control Program - (559) 445-5472

## PESTICIDE LABELS

CDMS – Ag Chemical Information Services  
<http://www.cdms.net/pfa/LUpdate.Msg.asp>  
GREENBOOK – <http://www.greenbook.net/>

## MARKET NEWS

<http://www.produceforsale.com/producemarkets.htm>

## The Vegetable Notes Newsletter is available ONLINE.

To download this or previous editions go to  
*UCCE Tulare County website:*  
[http://cetulare.ucdavis.edu/Vegetable\\_Crops/](http://cetulare.ucdavis.edu/Vegetable_Crops/)  
You can also sign up to receive this newsletter online.  
We welcome your comments. Send to newsletter editor:  
[mlestrange@ucdavis.edu](mailto:mlestrange@ucdavis.edu)

### Other UCCE county websites in the SJV:

*Fresno County:* <http://cefresno.ucdavis.edu>  
*Kern County:* <http://cekern.ucdavis.edu>  
*Kings County:* <http://cekings.ucdavis.edu>  
*Merced County:* <http://cemerced.ucdavis.edu>  
*San Joaquin County:* <http://cesanjoaquin.ucdavis.edu>  
*Stanislaus County:* <http://cestanislaus.ucdavis.edu>



# Vegetable Notes

*UCCE Tulare & Kings and Fresno Counties*  
*Michelle Le Strange and Shannon Mueller, Farm Advisors*

**Newsletter Volume 2, Issue #2:**

*Fresh Tomato & Bell Pepper  
Research Progress Reports*

The University of California prohibits discrimination or harassment of any person on the basis of race, color, national origin, religion, sex, gender identity, pregnancy (including childbirth, and medical conditions related to pregnancy or childbirth), physical or mental disability, medical condition (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or status as a covered veteran (covered veterans are special disabled veterans, recently separated veterans, Vietnam era veterans, or any other veterans who served on active duty during a war or in a campaign or expedition for which a campaign badge has been authorized) in any of its programs or activities.

University policy is intended to be consistent with the provisions of applicable State and Federal laws.

Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Staff Personnel Services Director, University of California, Agriculture and Natural Resources, 300 Lakeside Drive, 6<sup>th</sup> Floor, Oakland, CA 94612-3550, (510) 987-0096.

Cooperative Extension  
University of California  
4437-B S. Laspina Str.  
Tulare, CA 93274

Nonprofit Organization  
US Postage Paid  
Visalia, CA 93277  
Permit No. 240