University of California

Agriculture and Natural Resources

Making a Difference for California



(Estimated Crop Evapotranspiration or ET_C)

04/06/04 41

Crops (Leafout Date)	#	#148 Merce	d		#39 Parlier		#258 Lemon Cove				
	04/26 - 05/02	Accum'd	05/03 - 05/09	04/26 - 05/02	Accum'd	05/03 - 05/09	04/26 - 05/02	Accum'd	05/03 - 05/09		
	Water	Seasonal	Estimated	Water	Seasonal	Estimated	Water	Seasonal	Estimated		
	Use	Water Use	ETc	Use	Water Use	ETc	Use	Water Use	ETc		
Almonds (3/1) *	1.18	6.55	1.31	1.25	6.78	1.33	1.13	6.45	1.41		
Pistachio (4/20) * **	0.30	0.42	0.51	0.32	0.43	0.53	0.28	0.39	0.55		
Citrus (2/1)	0.97	7.81	1.05	1.04	8.16	1.05	0.94	7.82	1.11		
Raisin Grapes (3/11) (11 ft. row spacing)	0.42	1.18	0.51	0.48	1.24	0.52	0.43	1.17	0.55		
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	0.44	1.56	0.51	0.48	1.61	0.52	0.43	1.53	0.55		
Walnuts (4/20)	0.15	0.27	0.80	0.19	0.33	0.82	0.18	0.33	0.84		
Stone Fruit (3/11)	0.75	2.72	0.80	0.80	2.84	0.82	0.73	2.74	0.84		
Past 7 days precipitation (inches)		0.00			0.00			0.06			
Accumulated precipitation (inches) (1/1/2024)		14.98			8.64			9.12			

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of atrip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹													
Crops		#148 Merce	d			#39 Parlier							
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/1)	1.8	1.6	1.4	1.2	1.9	1.7	1.5	1.3	1.7	1.5	1.3	1.2	
Pistachio (4/20)	0.5	0.4	0.4	0.3	0.5	0.4	0.4	0.3	0.4	0.4	0.3	0.3	
Citrus (2/1)	1.5	1.3	1.1	1.0	1.6	1.4	1.2	1.1	1.4	1.3	1.1	1.0	
Raisin Grapes (3/11) (11 ft. row spacing)***	As	ssume all gra	pe	0.4	Assume all grape 0.5			0.5	As	ipe	0.4		
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis) ***	irrig	ation type is	drip	0.5	irrigation type is drip		drip	0.5	irrigation type is		drip	0.5	
Walnuts (4/20)	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.2	0.2	
Stone Fruit (3/11)	1.2	1.0	0.9	0.8	1.2	1.1	0.9	0.8	1.1	1.0	0.9	0.8	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

	PAST	WEEKLY	APPLIED W	ATER IN G	GALLON PER	R TREE OR	VINE					
Crops		#148 Merce	ed			#39 Parlier						
Almonds 115 Trees/A	425	378	331	283	449	401	354	307	401	354	307	283
Pistachio 106 Trees/A	125	100	100	75	125	100	100	75	100	100	75	75
Citrus 110 Trees/A	370	321	272	247	395	346	296	272	346	321	272	247
Raisin Grapes 566 Vines/A	Assume all grape			19	Assume all grape 24			24	A	19		
Winegrapes 622 Vines/A	irrig	ation type is	drip	22	irrigation type is drip			22	irrigation type is drip			22
Walnuts 76 Trees/A	71	71	71	71	107	107	71	71	107	71	71	71
Stonefruit 172 Trees/A	189	158	142	126	189	174	142	126	174	158	142	126
For further information concerning all counties receiving this report, contact	et the Fresno C	o. Farm Advi	sor's office at	(559) 241-7	526.							

HEALTHY FOOD SYSTEMS . HEALTHY ENVIRONMENTS . HEALTHY COMMUNITIES . HEALTHY CALIFORNIANS UCCE/DWR Weekly Crop Water Use Report

HEALTHY FOOD SYSTEMS . HEALTHY ENVIRONMENTS . HEALTHY COMMUNITIES . HEALTHY CALIFORNIANS

University of California

Agriculture and Natural Resources

Making a Difference for California

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_{C})

Crops (Leafout Date)	#	#124 Panoche				2 Five Point	ts	#15 Stratford		
	04/26- 05/02	Accum'd	05/03- 05/09		04/26- 05/02	Accum'd	05/03-05/09	04/26- 05/02	Accum'd	05/03- 05/09
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc
Almonds $(3/1) *$	1.44	6.63	1.52		1.51	6.77	1.54	1.27	6.36	1.54
Pistachio (4/20) * **	0.37	0.48	0.60		0.39	0.52	0.62	0.33	0.44	0.62
Citrus (2/1)	1.22	8.34	1.24		1.29	8.67	1.26	1.07	8.24	1.26
Raisin Grapes (3/11) (11 ft. row spacing)	0.51	1.28	0.58		0.54	1.34	0.60	0.45	1.22	0.60
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	0.53	1.68	0.58		0.55	1.69	0.60	0.48	1.59	0.60
Walnuts (4/20)	0.21	0.36	0.91		0.22	0.38	0.91	0.19	0.35	0.91
Stone Fruit (3/11)	0.92	3.03	0.91		0.98	3.02	0.91	 0.83	2.79	0.91
Past 7 days precipitation (inches)		0.00				0.00			0.00	
Accumulated precipitation (inches) (1/1/2024)		6.41				6.69			5.29	

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹													
Crops	#124 Panoche					#2 Five Poi	nts						
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/1)	2.2	1.9	1.7	1.5	2.3	2.0	1.8	1.6	2.0	1.7	1.5	1.3	
Pistachio (4/20)	0.6	0.5	0.4	0.4	0.6	0.5	0.5	0.4	0.5	0.4	0.4	0.3	
Citrus (2/1)	1.9	1.6	1.4	1.3	2.0	1.7	1.5	1.4	1.6	1.4	1.3	1.1	
Raisin Grapes (3/11) (11 ft. row spacing)	As	ssume all gra	ipe	0.5	Assume all grape 0.6			0.6	As	ipe	0.5		
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	irrig	ation type is	drip	0.6	irrigation type is drip		drip	0.6	irrigation type is		drip	0.5	
Walnuts (4/20)	0.3	0.3	0.2	0.2	0.3	0.3	0.3	0.2	0.3	0.3	0.2	0.2	
Stone Fruit (3/11)	1.4	1.2	1.1	1.0	1.5	1.3	1.2	1.0	1.3	1.1	1.0	0.9	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

	PAST	WEEKLY .	APPLIED W	ATER IN G	GALLON PER	R TREE OR	VINE					
Crops		#124 Panoc	he			#2 Five Point	nts					
Almonds 115 Trees/A	519 449 401			354	543	472	425	378	472	401	354	307
Pistachio 106 Trees/A	149	125	100	100	149	125	125	100	125	100	100	75
Citrus 110 Trees/A	469	395	346	321	494	420	370	346	395	346	321	272
Raisin Grapes 566 Vines/A	Assume all grape			24	Assume all grape 29			29	As	24		
Winegrapes 622 Vines/A	irrig	ation type is	drip	26	irrigation type is drip		26	irrigation type is drip		drip	22	
Walnuts 76 Trees/A	107	107	71	71	107	107	107	71	107	107	71	71
Stonefruit 172 Trees/A	221	189	174	158	237	205	189	158	205	174	158	142
For further information concerning all counties receiving this report, conta	ct the Fresno C	Co. Farm Adv	isor's office a	t (559) 241-7	7526.							

UCCE/DWR Weekly Crop Water Use Report