

WEEKLY SOIL MOISTURE LOSS IN INCHES
(Estimated Crop Evapotranspiration or ET_c)
07/19/24 through 07/25/24

Crops (Leafout Date)	#148 Merced			#39 Parlier			#258 Lemon Cove		
	07/19 - 07/25 Water Use	Accum'd Seasonal Water Use	07/26 - 08/01 Estimated ET _c	07/19 - 07/25 Water Use	Accum'd Seasonal Water Use	07/26 - 08/01 Estimated ET _c	07/19 - 07/25 Water Use	Accum'd Seasonal Water Use	07/26 - 08/01 Estimated ET _c
Almonds (3/1) *	2.03	28.34	1.99	2.14	29.77	1.91	2.00	28.89	1.90
Pistachio (4/20) * **	2.10	20.69	2.06	2.21	21.80	1.98	2.06	21.18	1.97
Citrus (2/1)	1.26	22.12	1.22	1.31	23.34	1.14	1.21	22.62	1.13
Raisin Grapes (3/11) (11 ft. row spacing)	1.37	15.43	1.33	1.43	16.22	1.27	1.34	15.78	1.26
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	1.55	16.65	1.51	1.64	17.57	1.45	1.52	17.07	1.44
Walnuts (4/20)	2.17	18.29	2.13	2.28	19.39	2.05	2.13	18.84	2.04
Stone Fruit (3/11)	1.96	20.62	2.06	2.08	21.89	1.98	1.93	21.22	1.97
Past 7 days precipitation (inches)	0.00			0.00			0.00		
Accumulated precipitation (inches) (1/1/2024)	14.98			8.98			9.71		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop
 * Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.
 ** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹

Crops	#148 Merced				#39 Parlier				#258 Lemon Cove			
	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
System Efficiency >>												
Almonds (3/1)	3.1	2.7	2.4	2.1	3.3	2.9	2.5	2.3	3.1	2.7	2.4	2.1
Pistachio (4/20)	3.2	2.8	2.5	2.2	3.4	2.9	2.6	2.3	3.2	2.7	2.4	2.2
Citrus (2/1)	1.9	1.7	1.5	1.3	2.0	1.7	1.5	1.4	1.9	1.6	1.4	1.3
Raisin Grapes (3/11) (11 ft. row spacing)	Assume all grape			1.4	Assume all grape			1.5	Assume all grape			1.4
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	irrigation type is drip			1.6	irrigation type is drip			1.7	irrigation type is drip			1.6
Walnuts (4/20)	3.3	2.9	2.6	2.3	3.5	3.0	2.7	2.4	3.3	2.8	2.5	2.2
Stone Fruit (3/11)	3.0	2.6	2.3	2.1	3.2	2.8	2.4	2.2	3.0	2.6	2.3	2.0

¹ The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

Crops	#148 Merced				#39 Parlier				#258 Lemon Cove			
Almonds 115 Trees/A	732	638	567	496	779	685	590	543	732	638	567	496
Pistachio 106 Trees/A	797	698	623	548	847	722	648	573	797	673	598	548
Citrus 110 Trees/A	469	420	370	321	494	420	370	346	469	395	346	321
Raisin Grapes 566 Vines/A	Assume all grape			67	Assume all grape			72	Assume all grape			67
Winegrapes 622 Vines/A	irrigation type is drip			70	irrigation type is drip			74	irrigation type is drip			70
Walnuts 76 Trees/A	1179	1036	929	822	1251	1072	965	857	1179	1000	893	786
Stonefruit 172 Trees/A	474	410	363	332	505	442	379	347	474	410	363	316

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.



WEEKLY SOIL MOISTURE LOSS IN INCHES
 (Estimated Crop Evapotranspiration or ET_c)
 07/19/24 through 07/25/24

Crops (Leafout Date)	#124 Panoche				#2 Five Points				#15 Stratford		
	07/19- 07/25 Water Use	Accum'd Seasonal Water Use	07/26- 08/01 Estimated ETc		07/19- 07/25 Water Use	Accum'd Seasonal Water Use	07/26- 08/01 Estimated ETc		07/19- 07/25 Water Use	Accum'd Seasonal Water Use	07/26- 08/01 Estimated ETc
Almonds (3/1) *	2.36	30.56	1.99		2.28	31.45	2.12		2.36	28.50	2.12
Pistachio (4/20) * **	2.41	22.82	2.06		2.33	23.49	2.19		2.40	21.08	2.19
Citrus (2/1)	1.42	24.14	1.22		1.39	24.99	1.28		1.42	22.80	1.28
Raisin Grapes (3/11) (11 ft. row spacing)	1.57	16.87	1.33		1.51	17.41	1.43		1.56	15.61	1.43
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	1.81	18.22	1.51		1.74	18.73	1.62		1.80	16.83	1.62
Walnuts (4/20)	2.50	20.29	2.13		2.41	20.90	2.26		2.49	18.84	2.26
Stone Fruit (3/11)	2.33	23.00	2.06		2.24	23.58	2.19		2.32	21.25	2.19
Past 7 days precipitation (inches)	0.00				0.00				0.00		
Accumulated precipitation (inches) (1/1/2024)	6.60				6.86				5.43		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop
 * Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.
 ** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹

Crops	#124 Panoche					#2 Five Points					#15 Stratford			
	65%	75%	85%	95%		65%	75%	85%	95%		65%	75%	85%	95%
System Efficiency >>	65%	75%	85%	95%		65%	75%	85%	95%		65%	75%	85%	95%
Almonds (3/1)	3.6	3.1	2.8	2.5		3.5	3.0	2.7	2.4		3.6	3.1	2.8	2.5
Pistachio (4/20)	3.7	3.2	2.8	2.5		3.6	3.1	2.7	2.5		3.7	3.2	2.8	2.5
Citrus (2/1)	2.2	1.9	1.7	1.5		2.1	1.9	1.6	1.5		2.2	1.9	1.7	1.5
Raisin Grapes (3/11) (11 ft. row spacing)	Assume all grape			1.7		Assume all grape			1.6		Assume all grape			1.6
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	irrigation type is drip			1.9		irrigation type is drip			1.8		irrigation type is drip			1.9
Walnuts (4/20)	3.8	3.3	2.9	2.6		3.7	3.2	2.8	2.5		3.8	3.3	2.9	2.6
Stone Fruit (3/11)	3.6	3.1	2.7	2.5		3.4	3.0	2.6	2.4		3.6	3.1	2.7	2.4

¹ The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

Crops	#124 Panoche					#2 Five Points					#15 Stratford			
	850	732	661	590		826	708	638	567		850	732	661	590
Almonds 115 Trees/A	850	732	661	590		826	708	638	567		850	732	661	590
Pistachio 106 Trees/A	922	797	698	623		897	772	673	623		922	797	698	623
Citrus 110 Trees/A	543	469	420	370		518	469	395	370		543	469	420	370
Raisin Grapes 566 Vines/A	Assume all grape			82		Assume all grape			77		Assume all grape			77
Winegrapes 622 Vines/A	irrigation type is drip			83		irrigation type is drip			79		irrigation type is drip			83
Walnuts 76 Trees/A	1358	1179	1036	929		1322	1143	1000	893		1358	1179	1036	929
Stonefruit 172 Trees/A	568	489	426	395		537	474	410	379		568	489	426	379

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.