#### University of California

**Agriculture and Natural Resources** 



HEALTHY FOOD SYSTEMS + HEALTHY ENVIRONMENTS + HEALTHY COMMUNITIES + HEALTHY CALIFORNIANS

## UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

#### WEEKLY SOIL MOISTURE LOSS IN INCHES (Estimated Crop Evapotranspiration or ET<sub>C</sub>) 07/26/24 through 08/01/24 Crops (Leafout Date) #148 Merced #39 Parlier #258 Lemon Cove 07/26 - 08/01 Accum'd 08/02 - 08/08 07/26 - 08/01 Accum'd 08/02 - 08/08 07/26 - 08/01 Accum'd 08/02 - 08/08 Water Seasonal Estimated Water Seasonal Estimated Water Seasonal Estimated Water Use ETc Water Use ETc Water Use ETc Use Use Use Almonds (3/1) \* 1.99 30.33 1.94 2.03 31.80 1.85 1.98 30.87 1.82 Pistachio (4/20) \* \*\* 2.06 22.75 2.01 2.10 23.90 1.88 2.05 23.23 1.82 Citrus (2/1) 1.22 23.34 1.17 1.24 24.58 1.12 23.82 1.12 1.20 Raisin Grapes (3/11) (11 ft. row spacing) 1.33 1.36 17.58 1.22 1.19 16.76 1.29 1.32 17.10 Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis) 1.51 18.17 1.47 1.55 19.12 1.39 1.50 18.57 1.37 Walnuts (4/20) 2.13 20.42 2.08 2.17 21.56 1.95 2.12 20.96 1.89 Stone Fruit (3/11) 2.06 22.68 2.01 2.10 23.99 2.05 23.27 1.82 1.88 0.00 0.00 0.00 Past 7 days precipitation (inches) 14.98 8.98 9.71 Accumulated precipitation (inches) (1/1/2024)

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

\* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

\*\* Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY <sup>1</sup>													
Crops		#148 Merce	d			#39 Parlier			#258 Lemon Cove				
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/1)	3.1	2.7	2.3	2.1	3.1	2.7	2.4	2.1	3.0	2.6	2.3	2.1	
Pistachio (4/20)	3.2	2.7	2.4	2.2	3.2	2.8	2.5	2.2	3.2	2.7	2.4	2.2	
Citrus (2/1)	1.9	1.6	1.4	1.3	1.9	1.7	1.5	1.3	1.8	1.6	1.4	1.3	
Raisin Grapes (3/11) (11 ft. row spacing)	Assume all grape			1.4	Assume all grape			1.4	Assume all grape			1.4	
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	irrigation type is drip			1.6	irrigation type is drip			1.6	irrigation type is drip			1.6	
Walnuts (4/20)	3.3	2.8	2.5	2.2	3.3	2.9	2.6	2.3	3.3	2.8	2.5	2.2	
Stone Fruit (3/11)	3.2	2.7	2.4	2.2	3.2	2.8	2.5	2.2	3.2	2.7	2.4	2.2	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE													
Crops	#148 Merced				#39 Parlier					#258 Lemon Cove			
Almonds 115 Trees/A	732	638	543	496	732	638	567	496	708	614	543	496	
Pistachio 106 Trees/A	797	673	598	548	797	698	623	548	797	673	598	548	
Citrus 110 Trees/A	469	395	346	321	469	420	370	321	444	395	346	321	
Raisin Grapes 566 Vines/A	Assume all grape			67	Assume all grape 67			67	A	67			
Winegrapes 622 Vines/A	irrig	ation type is	drip	70	irrigation type is drip			70	irrigation type is drip			70	
Walnuts 76 Trees/A	1179	1000	893	786	1179	1036	929	822	1179	1000	893	786	
Stonefruit 172 Trees/A	505	426	379	347	505	442	395	347	505	426	379	347	
For further information concerning all counties receiving this report, contact	the Fresno Co	. Farm Adviso	or's office at	(559) 241-752	26.								

## University of California **Agriculture and Natural Resources**

Making a Difference for California

#### WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Cron Evanatranspiration or FT)

	(Est		p Evapotransp 24 through 08		51 <sub>C</sub> )						
Crops (Leafout Date)	#	#124 Panoche				2 Five Point	ts	#15 Stratford			
	07/26- 08/01	Accum'd	08/02-08/08		07/26- 08/01	Accum'd	08/02- 08/08	07/26- 08/01	Accum'd	08/02-08/08	1
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc	
Almonds (3/1) *	2.10	32.66	1.93		2.11	33.56	2.08	2.17	30.67	2.08	
Pistachio (4/20) * **	2.17	24.99	2.00		2.18	25.67	2.15	2.24	23.32	2.15	
Citrus (2/1)	1.28	25.42	1.16		1.28	26.27	1.26	1.33	24.13	1.26	
Raisin Grapes (3/11) (11 ft. row spacing)	1.41	18.28	1.28		1.41	18.82	1.41	1.45	17.07	1.41	
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	1.61	19.82	1.46		1.62	20.34	1.59	1.67	18.50	1.59	
Walnuts (4/20)	2.24	22.53	2.07		2.25	23.15	2.22	2.31	21.15	2.22	
Stone Fruit (3/11)	2.17	25.17	2.00		2.18	25.76	2.15	 2.24	23.49	2.11	
Past 7 days precipitation (inches)		0.00				0.00			0.00		
Accumulated precipitation (inches) (1/1/2024)		6.60				6.86			5.43		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

\* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

\*\* Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

Crops		#124 Panoc	che			#2 Five Poin	nts					
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	3.2	2.8	2.5	2.2	3.2	2.8	2.5	2.2	3.3	2.9	2.6	2.3
Pistachio (4/20)	3.3	2.9	2.6	2.3	3.4	2.9	2.6	2.3	3.4	3.0	2.6	2.4
Citrus (2/1)	2.0	1.7	1.5	1.3	2.0	1.7	1.5	1.3	2.0	1.8	1.6	1.4
Raisin Grapes (3/11) (11 ft. row spacing)	As	ssume all gra	ape	1.5	A	ssume all gra	ipe	1.5	As	1.5		
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	irrig	ation type is	drip	1.7	irrigation type is drip			1.7	irrigation type is drip			1.8
Walnuts (4/20)	3.4	3.0	2.6	2.4	3.5	3.0	2.6	2.4	3.6	3.1	2.7	2.4
Stone Fruit (3/11)	3.3	2.9	2.6	2.3	3.4	2.9	2.6	2.3	3.4	3.0	2.6	2.4

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE												
Crops	#124 Panoche					#2 Five Poi	nts					
Almonds 115 Trees/A	756	661	590	519	756	661	590	519	779	685	614	543
Pistachio 106 Trees/A	822	722	648	573	847	722	648	573	847	747	648	598
Citrus 110 Trees/A	494	420	370	321	494	420	370	321	494	444	395	346
Raisin Grapes 566 Vines/A	Assume all grape			72	Assume all grape 72			72	As	72		
Winegrapes 622 Vines/A	irrig	ation type is	drip	74	irrigation type is drip			74	irrigation type is drip			79
Walnuts 76 Trees/A	1215	1072	929	857	1251	1072	929	857	1286	1108	965	857
Stonefruit 172 Trees/A	521	458	410	363	537	458	410	363	537	474	410	379
For further information concerning all counties receiving this report, conta	ct the Fresno C	Co. Farm Advi	isor's office at	t (559) 241-7	7526.							

For further information concerning all counties receiving this report, contact the Fresho Co. Farm Advisor's office at (559) 241-/526

# UCCE/DWR Weekly Crop Water Use Report